



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:58 PM UTC

PDB ID : 4XYL / pdb_00004xyl
Title : Ca. Korarchaeum cryptofilum ACD1 in complex with coenzyme A
Authors : Weisse, R.H.-J.; Scheidig, A.J.
Deposited on : 2015-02-02
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	3% 95% .
1	C	464	4% 95% .
2	B	230	25% 86% 11% ..
2	D	230	25% 82% 13% ..

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 21560 atoms, of which 10674 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha subunit of Acyl-CoA synthetase (NDP forming).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	463	Total	C	H	N	O	S	0	0	0
			6963	2210	3504	577	658	14			
1	C	463	Total	C	H	N	O	S	0	0	0
			6935	2204	3486	574	657	14			

- Molecule 2 is a protein called beta subunit of Acyl-CoA synthetase (NDP forming).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	225	Total	C	H	N	O	S	0	0	0
			3593	1130	1829	305	325	4			
2	D	226	Total	C	H	N	O	S	0	0	0
			3543	1122	1791	298	328	4			

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	C	1	Total Cl 1 1	0	0

- Molecule 4 is COENZYME A (CCD ID: COA) (formula: C₂₁H₃₆N₇O₁₆P₃S).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	P	S	0	0
			80	21	32	7	16	3	1		
4	C	1	Total	C	H	N	O	P	S	0	0
			80	21	32	7	16	3	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	177	Total	O	0	0
			177	177		
5	B	30	Total	O	0	0
			30	30		
5	C	126	Total	O	0	0
			126	126		
5	D	31	Total	O	0	0
			31	31		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: alpha subunit of Acyl-CoA synthetase (NDP forming)




- Molecule 1: alpha subunit of Acyl-CoA synthetase (NDP forming)




- Molecule 2: beta subunit of Acyl-CoA synthetase (NDP forming)



- Molecule 2: beta subunit of Acyl-CoA synthetase (NDP forming)





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	102.46Å 112.18Å 124.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.46 – 1.95 83.46 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (83.46-1.95) 99.8 (83.46-1.95)	Depositor EDS
R_{merge}	0.25	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 1.95Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.206 , 0.246 0.211 , 0.249	Depositor DCC
R_{free} test set	5288 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	27.4	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 28.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	21560	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, COA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/3527	0.69	0/4789
1	C	0.40	0/3516	0.69	0/4776
2	B	0.41	0/1791	0.70	1/2412 (0.0%)
2	D	0.39	0/1778	0.73	0/2398
All	All	0.40	0/10612	0.70	1/14375 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	66	VAL	N-CA-C	5.62	114.69	106.55

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3459	3504	3506	8	0
1	C	3449	3486	3487	11	0
2	B	1764	1829	1831	15	0
2	D	1752	1791	1793	20	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	48	32	32	1	0
4	C	48	32	32	0	0
5	A	177	0	0	0	0
5	B	30	0	0	0	0
5	C	126	0	0	1	0
5	D	31	0	0	0	0
All	All	10886	10674	10681	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:144:GLU:OE2	2:D:178:ARG:NH1	2.17	0.78
1:A:200:ASP:OD1	1:A:233:ARG:NH2	2.20	0.74
1:C:24:LYS:NZ	5:C:699:HOH:O	2.21	0.73
2:B:79:LEU:HD11	2:B:85:VAL:HA	1.75	0.68
2:B:68:HIS:O	2:B:68:HIS:ND1	2.28	0.66
2:B:44:GLU:OE2	2:B:86:ARG:NH2	2.34	0.61
2:D:27:ILE:HD12	2:D:32:ILE:HB	1.81	0.61
2:B:20:GLU:OE1	2:B:69:LYS:NZ	2.35	0.58
1:C:200:ASP:OD1	1:C:233:ARG:NH2	2.38	0.56
2:B:82:GLU:H	2:B:82:GLU:CD	2.13	0.56
2:D:209:ASP:OD2	2:D:226:ARG:NH2	2.40	0.55
2:D:27:ILE:HD11	2:D:34:VAL:HG22	1.88	0.54
2:D:15:ARG:HH11	2:D:15:ARG:HG2	1.73	0.53
2:B:82:GLU:O	2:B:85:VAL:HG22	2.08	0.52
1:A:418:GLU:O	1:A:421:GLU:HG3	2.09	0.52
2:D:87:LYS:O	2:D:91:GLU:HG2	2.11	0.51
2:D:121:ILE:CD1	2:D:143:VAL:HG21	2.42	0.50
1:C:113:GLU:HA	1:C:116:ARG:HG2	1.96	0.48
2:D:121:ILE:HD13	2:D:143:VAL:HG21	1.96	0.47
2:D:34:VAL:HG12	2:D:35:THR:N	2.28	0.47
2:B:68:HIS:CE1	2:B:71:ASP:HB2	2.49	0.47
1:C:116:ARG:HG3	1:C:117:ARG:N	2.29	0.46
2:B:21:HIS:HB3	2:B:109:LEU:HD13	1.98	0.46
1:A:348:ASN:HA	1:A:349:PRO:C	2.42	0.45
1:C:109:GLU:H	1:C:109:GLU:CD	2.25	0.45
2:B:121:ILE:CD1	2:B:143:VAL:HG21	2.47	0.44
1:A:161:GLY:HA3	4:A:502:COA:S1P	2.58	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:215:ILE:HD11	1:C:220:GLY:HA2	1.99	0.44
2:D:43:GLU:HG3	2:D:89:TYR:CD1	2.53	0.43
1:A:103:LYS:NZ	1:A:256:GLY:O	2.51	0.43
1:A:16:VAL:HA	1:A:45:ILE:HB	2.01	0.43
1:A:165:ILE:HG21	1:C:384:THR:HG22	2.01	0.42
2:D:65:GLN:HG2	2:D:103:ALA:HA	2.01	0.42
2:D:180:MET:HA	2:D:180:MET:HE2	2.01	0.42
2:B:121:ILE:HD13	2:B:143:VAL:HG21	2.01	0.42
2:D:203:GLU:O	2:D:230:ARG:HD2	2.19	0.42
2:B:89:TYR:CE1	2:B:93:ILE:HD11	2.55	0.42
2:D:121:ILE:O	2:D:121:ILE:HG13	2.19	0.42
1:A:219:ARG:HD2	1:A:222:MET:HE3	2.02	0.42
2:D:112:GLU:OE2	2:D:221:LYS:NZ	2.53	0.42
2:B:57:VAL:O	2:B:79:LEU:HB3	2.20	0.41
1:C:140:LEU:C	1:C:140:LEU:HD23	2.46	0.41
2:B:20:GLU:O	2:B:24:LYS:HE2	2.20	0.41
1:C:92:VAL:HG22	1:C:123:MET:HE2	2.02	0.41
2:D:105:ILE:HG22	2:D:107:GLY:N	2.35	0.41
2:B:43:GLU:HB2	2:B:89:TYR:CE1	2.56	0.41
2:D:44:GLU:O	2:D:47:VAL:HG22	2.21	0.41
2:D:67:VAL:HG13	2:D:68:HIS:CD2	2.55	0.41
1:C:362:TYR:OH	1:C:402:GLU:OE2	2.36	0.40
2:D:143:VAL:HG13	2:D:144:GLU:N	2.36	0.40
1:C:88:GLY:HA3	1:C:121:TYR:CG	2.57	0.40
2:D:27:ILE:HD11	2:D:34:VAL:CG2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	461/464 (99%)	450 (98%)	10 (2%)	1 (0%)	43 36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	461/464 (99%)	451 (98%)	9 (2%)	1 (0%)	43	36
2	B	221/230 (96%)	218 (99%)	3 (1%)	0	100	100
2	D	222/230 (96%)	218 (98%)	4 (2%)	0	100	100
All	All	1365/1388 (98%)	1337 (98%)	26 (2%)	2 (0%)	48	41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	348	ASN
1	C	348	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/370 (100%)	361 (98%)	7 (2%)	50	45
1	C	365/370 (99%)	360 (99%)	5 (1%)	59	56
2	B	192/197 (98%)	183 (95%)	9 (5%)	23	12
2	D	188/197 (95%)	171 (91%)	17 (9%)	9	2
All	All	1113/1134 (98%)	1075 (97%)	38 (3%)	32	23

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	LEU
1	A	117	ARG
1	A	249	ARG
1	A	340	VAL
1	A	359	ASP
1	A	386	VAL
1	A	421	GLU
2	B	67	VAL
2	B	77	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	79	LEU
2	B	82	GLU
2	B	83	GLU
2	B	87	LYS
2	B	96	VAL
2	B	110	VAL
2	B	204	GLU
1	C	2	ASN
1	C	34	LEU
1	C	92	VAL
1	C	140	LEU
1	C	340	VAL
2	D	15	ARG
2	D	27	ILE
2	D	42	SER
2	D	45	GLU
2	D	51	ARG
2	D	67	VAL
2	D	69	LYS
2	D	91	GLU
2	D	92	ILE
2	D	93	ILE
2	D	95	ASN
2	D	104	GLU
2	D	110	VAL
2	D	180	MET
2	D	204	GLU
2	D	221	LYS
2	D	230	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	313	GLN
1	C	28	GLN
1	C	313	GLN
2	D	78	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	COA	C	502	-	47,50,50	0.25	0	69,75,75	0.46	0
4	COA	A	502	-	47,50,50	0.29	0	69,75,75	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	COA	C	502	-	-	4/48/64/64	0/3/3/3
4	COA	A	502	-	-	5/48/64/64	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	502	COA	P2A-O3A-P1A-O1A
4	C	502	COA	C3B-O3B-P3B-O8A

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	502	COA	C2B-C1B-N9A-C8A
4	C	502	COA	C2B-C1B-N9A-C8A
4	C	502	COA	S1P-C2P-C3P-N4P
4	A	502	COA	C3B-O3B-P3B-O7A
4	A	502	COA	O4B-C1B-N9A-C8A
4	A	502	COA	P2A-O3A-P1A-O2A
4	C	502	COA	O4B-C1B-N9A-C8A


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	502	COA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	463/464 (99%)	0.02	16 (3%)	47 53	22, 31, 54, 76	0
1	C	463/464 (99%)	0.10	17 (3%)	45 52	22, 32, 54, 91	0
2	B	225/230 (97%)	1.06	57 (25%)	1 1	25, 40, 94, 107	0
2	D	226/230 (98%)	1.06	57 (25%)	1 1	27, 43, 106, 121	0
All	All	1377/1388 (99%)	0.39	147 (10%)	11 12	22, 34, 85, 121	0

All (147) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	404	GLY	7.0
2	B	96	VAL	6.9
2	B	92	ILE	6.3
1	C	255	THR	6.3
2	B	93	ILE	6.2
2	B	85	VAL	5.9
2	B	61	ILE	5.9
2	B	101	PRO	5.6
2	B	72	VAL	5.3
2	B	46	ALA	5.3
2	D	92	ILE	5.2
2	D	67	VAL	5.2
2	D	93	ILE	5.0
1	C	254	HIS	4.9
2	B	47	VAL	4.9
2	B	75	VAL	4.9
2	D	66	VAL	4.7
1	C	256	GLY	4.6
2	B	87	LYS	4.6
2	B	77	VAL	4.6
1	C	258	ILE	4.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	252	ALA	4.5
2	D	77	VAL	4.5
2	B	95	ASN	4.5
2	B	66	VAL	4.5
2	B	67	VAL	4.5
2	D	103	ALA	4.5
2	D	105	ILE	4.4
1	A	255	THR	4.4
2	D	71	ASP	4.3
2	B	88	ALA	4.3
2	B	73	GLY	4.3
2	B	100	VAL	4.2
2	D	64	PRO	4.0
1	C	343	PHE	4.0
2	B	89	TYR	4.0
2	D	96	VAL	4.0
2	B	74	GLY	3.9
2	B	68	HIS	3.9
2	D	40	ALA	3.8
2	D	70	SER	3.7
2	B	108	ILE	3.7
2	D	73	GLY	3.7
2	B	82	GLU	3.7
2	D	72	VAL	3.7
1	C	259	ALA	3.6
2	D	90	ARG	3.6
2	D	98	ARG	3.6
2	D	75	VAL	3.6
2	D	85	VAL	3.6
1	A	252	ALA	3.6
2	B	79	LEU	3.6
2	B	94	GLU	3.5
2	D	47	VAL	3.5
2	D	46	ALA	3.5
2	D	42	SER	3.5
2	B	40	ALA	3.5
2	D	55	PHE	3.4
2	D	62	VAL	3.4
2	D	61	ILE	3.3
1	A	404	GLY	3.3
1	C	257	SER	3.3
2	D	89	TYR	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	59	LEU	3.3
2	D	68	HIS	3.3
2	B	81	SER	3.3
2	D	97	LYS	3.3
2	B	90	ARG	3.2
2	D	88	ALA	3.2
2	B	99	ASN	3.1
1	C	253	SER	3.1
2	B	65	GLN	3.1
2	B	57	VAL	3.1
2	D	59	LEU	3.1
2	D	87	LYS	3.1
2	D	49	PHE	3.0
1	C	250	ALA	3.0
2	D	41	LYS	3.0
1	A	253	SER	2.9
2	B	63	SER	2.9
1	C	251	ALA	2.9
1	A	258	ILE	2.9
2	B	78	ASN	2.9
1	A	343	PHE	2.9
2	D	69	LYS	2.9
2	B	98	ARG	2.9
2	D	81	SER	2.9
2	D	79	LEU	2.9
2	B	58	VAL	2.9
2	B	62	VAL	2.9
2	B	49	PHE	2.9
1	C	462	ALA	2.8
2	D	99	ASN	2.8
2	B	43	GLU	2.8
2	B	45	GLU	2.8
1	A	406	SER	2.8
2	B	70	SER	2.8
2	D	57	VAL	2.8
2	D	63	SER	2.7
1	C	342	PRO	2.7
2	D	104	GLU	2.7
2	D	177	PHE	2.7
2	B	64	PRO	2.7
2	D	108	ILE	2.7
1	C	2	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	42	SER	2.6
1	A	256	GLY	2.5
2	D	74	GLY	2.5
2	B	91	GLU	2.5
1	A	342	PRO	2.5
2	B	97	LYS	2.4
2	B	178	ARG	2.4
2	D	107	GLY	2.4
2	D	86	ARG	2.4
2	D	83	GLU	2.4
2	D	80	ARG	2.4
2	D	113	PHE	2.3
2	D	45	GLU	2.3
1	A	254	HIS	2.3
2	D	110	VAL	2.3
2	B	48	ASN	2.3
2	D	65	GLN	2.2
1	A	251	ALA	2.2
2	D	78	ASN	2.2
2	D	95	ASN	2.2
2	B	44	GLU	2.2
1	A	384	THR	2.2
2	D	27	ILE	2.2
1	C	341	PRO	2.2
2	B	113	PHE	2.1
2	B	109	LEU	2.1
2	D	43	GLU	2.1
1	A	246	VAL	2.1
1	A	259	ALA	2.1
1	A	339	PHE	2.1
2	D	53	ILE	2.1
2	B	69	LYS	2.1
1	A	2	ASN	2.1
2	D	56	PRO	2.1
1	C	249	ARG	2.1
2	B	106	GLU	2.0
2	B	50	SER	2.0
2	B	86	ARG	2.0
2	B	84	GLU	2.0
2	B	76	LYS	2.0
2	B	55	PHE	2.0
2	D	58	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	COA	C	502	48/48	0.95	0.08	32,39,47,58	0
4	COA	A	502	48/48	0.96	0.07	25,31,41,50	0
3	CL	A	501	1/1	0.98	0.10	40,40,40,40	0
3	CL	C	501	1/1	0.98	0.07	39,39,39,39	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around COA C 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around COA A 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.