



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:28 PM UTC

PDB ID : 4Y06 / pdb_00004y06
Title : Crystal structure of the DAP BII (G675R) dipeptide complex
Authors : Sakamoto, Y.; Iizuka, I.; Tateoka, C.; Roppongi, S.; Fujimoto, M.; Nonaka, T.; Ogasawara, W.; Tanaka, N.
Deposited on : 2015-02-05
Resolution : 2.18 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8975 (2.20-2.16)
Clashscore	190562	9786 (2.20-2.16)
Ramachandran outliers	187476	9664 (2.20-2.16)
Sidechain outliers	187428	9664 (2.20-2.16)
RSRZ outliers	180081	8979 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	722	% 89% 7% .
1	B	722	% 89% 7% .

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 11405 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dipeptidyl aminopeptidase BII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	697	Total	C	N	O	S	0	0	0
			5378	3402	940	1017	19			
1	B	697	Total	C	N	O	S	0	0	0
			5378	3402	940	1017	19			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	675	ARG	GLY	engineered mutation	UNP V5YM14
B	675	ARG	GLY	engineered mutation	UNP V5YM14

- Molecule 2 is LEUCINE (CCD ID: LEU) (formula: C₆H₁₃NO₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			8	6	1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			8	6	1	1		

- Molecule 3 is GLUTAMIC ACID (CCD ID: GLU) (formula: $C_5H_9NO_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			10	5	1	4		
3	B	1	Total	C	N	O	0	0
			10	5	1	4		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total 3	Zn 3	0	0
5	B	2	Total 2	Zn 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	237	Total 237	O 237	0	0
6	B	287	Total 287	O 287	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dipeptidyl aminopeptidase BII



• Molecule 1: Dipeptidyl aminopeptidase BII



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	122.42Å 122.42Å 222.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.00 – 2.18 36.00 – 2.18	Depositor EDS
% Data completeness (in resolution range)	99.9 (36.00-2.18) 99.9 (36.00-2.18)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.20 (at 2.18Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.176 , 0.235 0.185 , 0.240	Depositor DCC
R_{free} test set	4361 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	36.3	Xtrriage
Anisotropy	0.015	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 33.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11405	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.27 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.8948e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	3/5496 (0.1%)	1.09	3/7449 (0.0%)
1	B	1.19	6/5496 (0.1%)	1.08	3/7449 (0.0%)
All	All	1.17	9/10992 (0.1%)	1.08	6/14898 (0.0%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	291	TRP	CA-C	8.00	1.60	1.52
1	A	284	GLU	N-CA	5.97	1.53	1.46
1	B	639	ILE	C-O	-5.49	1.18	1.24
1	B	82	VAL	N-CA	5.34	1.52	1.46
1	B	89	TYR	CA-C	5.25	1.59	1.52
1	B	389	THR	CA-C	5.12	1.59	1.52
1	B	202	PRO	C-O	-5.07	1.18	1.24
1	A	355	GLU	CA-C	5.02	1.59	1.52
1	B	624	ASP	C-O	-5.01	1.18	1.24

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	707	GLU	CA-C-N	-5.69	117.12	123.16
1	A	707	GLU	C-N-CA	-5.69	117.12	123.16
1	B	673	PHE	N-CA-C	5.39	121.54	114.75
1	B	569	GLU	CA-C-N	5.36	128.46	120.95
1	B	569	GLU	C-N-CA	5.36	128.46	120.95
1	A	571	VAL	N-CA-C	5.03	115.22	107.77

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5378	0	5294	18	0
1	B	5378	0	5294	22	0
2	A	8	0	10	0	0
2	B	8	0	10	0	0
3	A	10	0	6	1	0
3	B	10	0	6	1	0
4	A	36	0	48	1	0
4	B	48	0	64	3	0
5	A	3	0	0	0	0
5	B	2	0	0	0	0
6	A	237	0	0	1	0
6	B	287	0	0	3	0
All	All	11405	0	10732	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:675:ARG:NH1	6:B:901:HOH:O	2.32	0.63
1:A:306:MET:HE3	1:A:455:GLN:HB3	1.83	0.61
1:A:299:HIS:HD2	1:A:459:TYR:OH	1.87	0.58
1:B:442:LYS:HE3	1:B:507:LEU:HD21	1.85	0.57
1:A:287:ASN:OD1	1:A:390:ARG:NH1	2.37	0.57
1:B:141:ILE:HG23	1:B:151:ARG:HG2	1.87	0.56
1:A:635:GLU:OE1	6:A:901:HOH:O	2.17	0.56
1:B:48:LEU:HD12	1:B:53:ILE:HD11	1.89	0.55
1:B:657:SER:HB3	3:B:802:GLU:C	2.32	0.54
1:B:609:LYS:CE	4:B:807:GOL:H11	2.38	0.54
1:B:592:LYS:HD2	1:B:595:MET:HE3	1.90	0.53
1:B:222:THR:H	1:B:645:ASN:HD21	1.56	0.53
1:A:657:SER:OG	3:A:802:GLU:C	2.53	0.51
1:B:303:GLN:HA	1:B:306:MET:HE3	1.95	0.49
1:A:218:TRP:CG	1:A:219:PRO:HA	2.49	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:222:THR:H	1:A:645:ASN:HD21	1.61	0.47
1:B:78:ASN:HB2	6:B:1058:HOH:O	2.14	0.47
1:B:287:ASN:OD1	1:B:390:ARG:NH1	2.47	0.46
1:A:472:ARG:NH2	1:A:483:ASN:OD1	2.36	0.46
1:A:376:HIS:CE1	1:A:380:LEU:HD21	2.52	0.45
1:A:303:GLN:NE2	1:A:460:TRP:HE1	2.15	0.45
1:B:84:ASN:ND2	1:B:657:SER:HB2	2.32	0.45
1:A:389:THR:HA	1:A:471:GLN:HE22	1.82	0.44
1:A:549:ASN:HD21	1:A:553:ARG:HE	1.66	0.43
1:B:218:TRP:CG	1:B:219:PRO:HA	2.54	0.43
1:A:448:TYR:OH	4:A:803:GOL:H2	2.19	0.42
1:B:622:LEU:C	1:B:622:LEU:HD23	2.45	0.42
1:B:93:GLN:NE2	1:B:447:ARG:HE	2.18	0.42
1:A:261:LEU:HD22	1:A:669:VAL:HB	2.02	0.42
1:B:28:MET:HE3	1:B:28:MET:HB2	1.93	0.41
1:A:127:LEU:HA	1:A:190:ASN:HD22	1.84	0.41
1:A:473:VAL:CG1	1:A:534:THR:HG21	2.49	0.41
1:A:553:ARG:N	1:A:554:PRO:HD2	2.35	0.41
1:B:592:LYS:CD	1:B:595:MET:HE3	2.50	0.41
1:B:281:LEU:HD23	1:B:380:LEU:HD13	2.03	0.41
1:B:219:PRO:HD2	4:B:807:GOL:H12	2.02	0.41
1:B:687:ASP:OD1	1:B:687:ASP:C	2.63	0.41
1:A:592:LYS:NZ	1:A:595:MET:HE1	2.36	0.41
4:B:803:GOL:H31	6:B:952:HOH:O	2.21	0.40
1:B:93:GLN:HE22	1:B:447:ARG:HE	1.68	0.40
1:B:292:THR:HG22	1:B:296:ILE:HD12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	695/722 (96%)	677 (97%)	17 (2%)	1 (0%)	48	55
1	B	695/722 (96%)	680 (98%)	14 (2%)	1 (0%)	48	55
All	All	1390/1444 (96%)	1357 (98%)	31 (2%)	2 (0%)	48	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	366	GLY
1	A	360	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	542/559 (97%)	527 (97%)	15 (3%)	38	49
1	B	542/559 (97%)	528 (97%)	14 (3%)	40	51
All	All	1084/1118 (97%)	1055 (97%)	29 (3%)	39	50

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	LYS
1	A	51	GLN
1	A	59	ASP
1	A	99	GLU
1	A	234	LYS
1	A	243	LYS
1	A	318	GLN
1	A	368	LYS
1	A	398	LEU
1	A	492	LEU
1	A	513	ASP
1	A	535	LEU
1	A	548	GLU
1	A	613	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	689	LYS
1	B	41	LEU
1	B	43	LYS
1	B	51	GLN
1	B	104	LYS
1	B	112	THR
1	B	146	ASP
1	B	352	LEU
1	B	358	VAL
1	B	365	GLN
1	B	380	LEU
1	B	550	LEU
1	B	557	LEU
1	B	565	LYS
1	B	613	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (32) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	GLN
1	A	84	ASN
1	A	93	GLN
1	A	105	ASN
1	A	190	ASN
1	A	215	ASN
1	A	299	HIS
1	A	303	GLN
1	A	330	ASN
1	A	334	ASN
1	A	350	GLN
1	A	365	GLN
1	A	376	HIS
1	A	471	GLN
1	A	549	ASN
1	A	558	GLN
1	A	585	ASN
1	A	645	ASN
1	A	683	ASN
1	B	84	ASN
1	B	93	GLN
1	B	249	GLN
1	B	303	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	334	ASN
1	B	350	GLN
1	B	400	ASN
1	B	462	ASN
1	B	558	GLN
1	B	585	ASN
1	B	613	GLN
1	B	645	ASN
1	B	683	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 5 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LEU	A	801	3	5,7,8	0.78	0	6,8,10	0.63	0
4	GOL	B	807	-	5,5,5	1.00	0	5,5,5	2.22	3 (60%)
2	LEU	B	801	3	5,7,8	0.65	0	6,8,10	0.87	1 (16%)
4	GOL	B	804	-	5,5,5	0.82	0	5,5,5	1.63	2 (40%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	805	-	5,5,5	0.71	0	5,5,5	0.84	0
4	GOL	A	803	-	5,5,5	1.02	1 (20%)	5,5,5	1.34	1 (20%)
4	GOL	B	803	-	5,5,5	1.32	1 (20%)	5,5,5	2.16	2 (40%)
4	GOL	B	805	-	5,5,5	0.63	0	5,5,5	0.37	0
3	GLU	B	802	2	8,9,9	1.13	0	8,11,11	1.13	1 (12%)
4	GOL	A	806	-	5,5,5	0.58	0	5,5,5	0.38	0
4	GOL	A	808	-	5,5,5	0.35	0	5,5,5	0.22	0
4	GOL	B	808	-	5,5,5	0.74	0	5,5,5	0.55	0
4	GOL	B	810	-	5,5,5	0.34	0	5,5,5	0.72	0
4	GOL	A	804	-	5,5,5	0.46	0	5,5,5	0.57	0
4	GOL	A	807	-	5,5,5	0.51	0	5,5,5	0.77	0
3	GLU	A	802	2	8,9,9	1.18	1 (12%)	8,11,11	0.73	0
4	GOL	B	809	-	5,5,5	0.33	0	5,5,5	0.58	0
4	GOL	B	806	-	5,5,5	0.22	0	5,5,5	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LEU	A	801	3	-	0/5/6/8	-
4	GOL	B	807	-	-	1/4/4/4	-
2	LEU	B	801	3	-	0/5/6/8	-
4	GOL	B	804	-	-	2/4/4/4	-
4	GOL	A	805	-	-	2/4/4/4	-
4	GOL	A	803	-	-	3/4/4/4	-
4	GOL	B	803	-	-	1/4/4/4	-
4	GOL	B	805	-	-	0/4/4/4	-
3	GLU	B	802	2	-	2/9/9/9	-
4	GOL	A	806	-	-	2/4/4/4	-
4	GOL	A	808	-	-	0/4/4/4	-
4	GOL	B	808	-	-	2/4/4/4	-
4	GOL	B	810	-	-	4/4/4/4	-
4	GOL	A	804	-	-	2/4/4/4	-
4	GOL	A	807	-	-	0/4/4/4	-
3	GLU	A	802	2	-	2/9/9/9	-
4	GOL	B	809	-	-	2/4/4/4	-
4	GOL	B	806	-	-	1/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	803	GOL	O2-C2	2.37	1.50	1.43
3	A	802	GLU	OE2-CD	-2.15	1.23	1.30
4	A	803	GOL	O2-C2	2.09	1.49	1.43

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	807	GOL	C3-C2-C1	3.20	123.54	111.80
4	B	803	GOL	O2-C2-C1	3.11	122.07	109.18
4	B	803	GOL	C3-C2-C1	-2.81	101.51	111.80
4	B	807	GOL	O1-C1-C2	2.66	122.36	110.38
4	B	804	GOL	O3-C3-C2	2.63	122.23	110.38
4	B	807	GOL	O2-C2-C1	-2.53	98.72	109.18
3	B	802	GLU	OXT-C-O	-2.49	118.42	124.08
2	B	801	LEU	CB-CA-C	-2.06	107.81	110.99
4	B	804	GOL	O2-C2-C3	2.05	117.65	109.18
4	A	803	GOL	O2-C2-C3	2.00	117.48	109.18

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	805	GOL	C1-C2-C3-O3
4	B	804	GOL	O1-C1-C2-O2
4	B	804	GOL	O1-C1-C2-C3
4	B	808	GOL	C1-C2-C3-O3
4	B	809	GOL	O1-C1-C2-C3
4	B	808	GOL	O2-C2-C3-O3
4	A	803	GOL	O1-C1-C2-C3
4	A	804	GOL	O1-C1-C2-C3
4	A	806	GOL	C1-C2-C3-O3
4	B	806	GOL	O1-C1-C2-C3
4	B	807	GOL	O1-C1-C2-C3
4	B	810	GOL	O1-C1-C2-C3
4	B	810	GOL	C1-C2-C3-O3
4	A	806	GOL	O2-C2-C3-O3
4	A	805	GOL	O2-C2-C3-O3
4	B	803	GOL	O1-C1-C2-O2
4	B	809	GOL	O1-C1-C2-O2
4	B	810	GOL	O1-C1-C2-O2
4	A	803	GOL	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	803	GOL	O2-C2-C3-O3
3	B	802	GLU	OXT-C-CA-CB
3	A	802	GLU	OXT-C-CA-CB
4	A	804	GOL	O1-C1-C2-O2
3	A	802	GLU	OE1-CD-CG-CB
4	B	810	GOL	O2-C2-C3-O3
3	B	802	GLU	OE2-CD-CG-CB

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	807	GOL	2	0
4	A	803	GOL	1	0
4	B	803	GOL	1	0
3	B	802	GLU	1	0
3	A	802	GLU	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	697/722 (96%)	0.00	8 (1%) 78 77	23, 40, 64, 97	0
1	B	697/722 (96%)	-0.18	5 (0%) 84 83	22, 36, 59, 81	0
All	All	1394/1444 (96%)	-0.09	13 (0%) 81 80	22, 38, 62, 97	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	570	PHE	6.4
1	B	570	PHE	4.5
1	A	721	GLY	3.2
1	A	468	PRO	3.2
1	A	469	ALA	3.1
1	A	367	ALA	2.8
1	B	145	GLY	2.6
1	A	366	GLY	2.5
1	B	721	GLY	2.3
1	A	360	GLY	2.2
1	B	47	LYS	2.2
1	A	467	LEU	2.1
1	B	135	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	804	6/6	0.79	0.19	65,70,72,78	0
4	GOL	B	803	6/6	0.84	0.13	39,42,55,56	0
4	GOL	B	805	6/6	0.85	0.18	37,59,60,65	0
4	GOL	B	807	6/6	0.85	0.16	36,52,56,67	0
4	GOL	B	808	6/6	0.85	0.14	49,62,67,68	0
4	GOL	A	806	6/6	0.89	0.12	56,64,67,68	0
4	GOL	A	805	6/6	0.90	0.12	58,62,65,69	0
4	GOL	B	806	6/6	0.91	0.13	63,63,66,72	0
3	GLU	B	802	10/10	0.92	0.09	28,33,40,48	0
4	GOL	A	807	6/6	0.93	0.11	42,56,58,65	0
4	GOL	A	803	6/6	0.93	0.10	43,48,54,56	0
5	ZN	A	810	1/1	0.94	0.10	96,96,96,96	0
2	LEU	A	801	8/9	0.95	0.09	28,30,34,39	0
4	GOL	B	809	6/6	0.95	0.09	31,46,47,49	0
3	GLU	A	802	10/10	0.95	0.08	30,32,39,51	0
2	LEU	B	801	8/9	0.96	0.09	21,26,29,33	0
4	GOL	B	810	6/6	0.96	0.07	36,44,47,55	0
4	GOL	B	804	6/6	0.96	0.08	31,45,51,57	0
4	GOL	A	808	6/6	0.97	0.06	37,39,43,53	0
5	ZN	A	809	1/1	0.98	0.03	60,60,60,60	0
5	ZN	A	811	1/1	0.98	0.07	47,47,47,47	0
5	ZN	B	812	1/1	0.98	0.04	57,57,57,57	0
5	ZN	B	811	1/1	0.99	0.09	50,50,50,50	0

6.5 Other polymers

There are no such residues in this entry.