



Full wwPDB NMR Structure Validation Report ⓘ

Mar 8, 2026 – 11:43 AM UTC

PDB ID : 1Y15 / pdb_00001y15
Title : Mouse Prion Protein with mutation N174T
Authors : Gossert, A.D.; Bonjour, S.; Lysek, D.A.; Fiorito, F.; Wuthrich, K.
Deposited on : 2004-11-17

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	112	71% 11% 18%

2 Ensemble composition and analysis

This entry contains 20 models. Model 16 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:129-A:166, A:172-A:225 (92)	0.66	16

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 5, 9, 13, 19
2	3, 6, 10, 14, 15, 16, 20
3	8, 11, 18
Single-model clusters	7; 12; 17

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1791 atoms, of which 865 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Major prion protein.

Mol	Chain	Residues	Atoms						Trace
1	A	112	Total	C	H	N	O	S	0
			1791	575	865	162	180	9	

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	174	THR	ASN	engineered mutation	UNP P04925

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Major prion protein




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Major prion protein




4.2.2 Score per residue for model 2

- Molecule 1: Major prion protein




4.2.3 Score per residue for model 3

- Molecule 1: Major prion protein




4.2.4 Score per residue for model 4

- Molecule 1: Major prion protein




4.2.5 Score per residue for model 5

- Molecule 1: Major prion protein



4.2.6 Score per residue for model 6


- Molecule 1: Major prion protein



4.2.7 Score per residue for model 7


- Molecule 1: Major prion protein






4.2.8 Score per residue for model 8

- Molecule 1: Major prion protein




4.2.9 Score per residue for model 9

- Molecule 1: Major prion protein




4.2.10 Score per residue for model 10

- Molecule 1: Major prion protein




4.2.11 Score per residue for model 11

- Molecule 1: Major prion protein




4.2.12 Score per residue for model 12

- Molecule 1: Major prion protein




4.2.13 Score per residue for model 13

- Molecule 1: Major prion protein




4.2.14 Score per residue for model 14

- Molecule 1: Major prion protein




4.2.15 Score per residue for model 15

- Molecule 1: Major prion protein




4.2.16 Score per residue for model 16 (medoid)

- Molecule 1: Major prion protein




4.2.17 Score per residue for model 17

- Molecule 1: Major prion protein




4.2.18 Score per residue for model 18

- Molecule 1: Major prion protein




4.2.19 Score per residue for model 19

- Molecule 1: Major prion protein



4.2.20 Score per residue for model 20

- Molecule 1: Major prion protein



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	6.2
CANDID	refinement	1.0

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.70±0.01	0±0/794 (0.0± 0.0%)	1.49±0.04	4±2/1075 (0.4± 0.2%)
All	All	0.70	0/15880 (0.0%)	1.49	81/21500 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	2.4±1.6
All	All	0	48

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	132	SER	CA-C-N	6.77	134.47	121.54	3	10
1	A	132	SER	C-N-CA	6.77	134.47	121.54	3	10
1	A	197	ASN	OD1-CG-ND2	-6.74	115.86	122.60	13	1
1	A	223	GLN	CA-C-N	6.57	129.08	120.28	13	8
1	A	223	GLN	C-N-CA	6.57	129.08	120.28	13	8
1	A	202	ASP	CA-CB-CG	6.34	118.94	112.60	9	1
1	A	161	VAL	CB-CA-C	6.34	117.63	110.41	19	2
1	A	175	PHE	CA-CB-CG	6.14	119.94	113.80	17	4
1	A	140	HIS	CB-CG-CD2	-6.10	123.27	131.20	12	3
1	A	225	TYR	CA-C-N	5.97	128.55	120.38	4	5
1	A	225	TYR	C-N-CA	5.97	128.55	120.38	4	5
1	A	147	ASP	CA-CB-CG	-5.83	106.77	112.60	13	1
1	A	209	VAL	CB-CA-C	5.79	119.11	111.70	8	2
1	A	173	ASN	CA-CB-CG	5.54	118.14	112.60	4	1
1	A	187	HIS	CB-CG-CD2	-5.37	124.22	131.20	12	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	177	HIS	CB-CG-CD2	-5.32	124.28	131.20	13	3
1	A	190	THR	CA-CB-CG2	5.30	119.50	110.50	16	1
1	A	188	THR	CA-C-O	-5.26	115.29	120.82	16	1
1	A	132	SER	N-CA-C	5.26	117.46	108.67	20	1
1	A	166	VAL	CA-CB-CG1	5.25	119.33	110.40	18	1
1	A	221	GLU	N-CA-C	-5.18	105.52	111.07	19	1
1	A	175	PHE	CA-C-N	5.16	127.86	122.14	19	1
1	A	175	PHE	C-N-CA	5.16	127.86	122.14	19	1
1	A	174	THR	CA-C-N	5.15	127.45	120.65	14	1
1	A	174	THR	C-N-CA	5.15	127.45	120.65	14	1
1	A	154	MET	CA-C-N	5.15	130.57	122.49	6	1
1	A	154	MET	C-N-CA	5.15	130.57	122.49	6	1
1	A	210	VAL	CA-C-O	-5.12	115.62	120.95	14	1
1	A	191	THR	CB-CA-C	5.08	120.53	110.42	2	1
1	A	208	ARG	CA-CB-CG	5.06	124.22	114.10	6	1
1	A	191	THR	N-CA-C	-5.04	107.19	113.55	14	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	164	ARG	Sidechain	7
1	A	157	TYR	Sidechain	6
1	A	148	ARG	Sidechain	5
1	A	218	TYR	Sidechain	4
1	A	137	PRO	Peptide	4
1	A	163	TYR	Sidechain	3
1	A	149	TYR	Sidechain	3
1	A	162	TYR	Sidechain	3
1	A	136	ARG	Sidechain	3
1	A	156	ARG	Sidechain	3
1	A	155	TYR	Sidechain	1
1	A	175	PHE	Sidechain	1
1	A	208	ARG	Sidechain	1
1	A	131	GLY	Peptide	1
1	A	225	TYR	Sidechain	1
1	A	198	PHE	Sidechain	1
1	A	219	GLN	Peptide	1

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	775	731	731	1±1
All	All	15500	14620	14620	18

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:130:LEU:HD23	1:A:162:TYR:CE1	0.52	2.39	5	1
1:A:139:ILE:CD1	1:A:209:VAL:HG23	0.51	2.35	17	1
1:A:199:THR:O	1:A:203:VAL:HG23	0.48	2.09	19	2
1:A:198:PHE:CE1	1:A:206:MET:HE2	0.47	2.44	7	1
1:A:139:ILE:HG22	1:A:140:HIS:H	0.47	1.70	4	1
1:A:139:ILE:HD11	1:A:209:VAL:HA	0.45	1.87	11	4
1:A:134:MET:HE1	1:A:217:GLN:CG	0.44	2.42	18	1
1:A:139:ILE:HG21	1:A:141:PHE:CE1	0.44	2.46	4	1
1:A:131:GLY:C	1:A:160:GLN:HE22	0.44	2.21	15	1
1:A:139:ILE:HD13	1:A:212:GLN:NE2	0.43	2.28	15	1
1:A:201:THR:HG22	1:A:205:MET:SD	0.42	2.54	6	1
1:A:166:VAL:HG22	1:A:221:GLU:HB3	0.41	1.92	7	1
1:A:153:ASN:HA	1:A:155:TYR:CE2	0.41	2.49	12	1
1:A:158:PRO:O	1:A:213:MET:HE1	0.40	2.16	2	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	92/112 (82%)	78±3 (85±4%)	11±3 (12±3%)	2±1 (2±1%)	6	43

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1840/2240 (82%)	1569 (85%)	225 (12%)	46 (2%)	6	43

All 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	133	ALA	19
1	A	131	GLY	7
1	A	188	THR	4
1	A	225	TYR	4
1	A	137	PRO	3
1	A	166	VAL	2
1	A	176	VAL	1
1	A	195	GLY	1
1	A	154	MET	1
1	A	132	SER	1
1	A	136	ARG	1
1	A	197	ASN	1
1	A	192	THR	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	87/102 (85%)	73±3 (84±3%)	14±3 (16±3%)	4	41
All	All	1740/2040 (85%)	1467 (84%)	273 (16%)	4	41

All 55 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	193	THR	16
1	A	138	MET	13
1	A	209	VAL	12
1	A	187	HIS	11
1	A	207	GLU	11
1	A	191	THR	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	190	THR	10
1	A	160	GLN	9
1	A	148	ARG	9
1	A	143	ASN	8
1	A	147	ASP	8
1	A	154	MET	8
1	A	155	TYR	8
1	A	221	GLU	8
1	A	197	ASN	8
1	A	135	SER	7
1	A	166	VAL	7
1	A	199	THR	7
1	A	132	SER	7
1	A	129	MET	6
1	A	220	LYS	5
1	A	177	HIS	5
1	A	136	ARG	5
1	A	144	ASP	5
1	A	130	LEU	5
1	A	173	ASN	4
1	A	196	GLU	4
1	A	140	HIS	3
1	A	194	LYS	3
1	A	198	PHE	3
1	A	164	ARG	3
1	A	175	PHE	3
1	A	204	LYS	3
1	A	219	GLN	3
1	A	185	LYS	3
1	A	217	GLN	3
1	A	159	ASN	2
1	A	200	GLU	2
1	A	206	MET	2
1	A	214	CYS	2
1	A	201	THR	2
1	A	156	ARG	2
1	A	172	GLN	2
1	A	182	ILE	2
1	A	186	GLN	2
1	A	210	VAL	2
1	A	181	ASN	2
1	A	208	ARG	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	223	GLN	1
1	A	152	GLU	1
1	A	216	THR	1
1	A	179	CYS	1
1	A	178	ASP	1
1	A	222	SER	1
1	A	212	GLN	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided