



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 06:05 AM UTC

PDB ID : 1Y19 / pdb_00001y19
Title : Structural basis for phosphatidylinositol phosphate kinase type I-gamma binding to talin at focal adhesions
Authors : de Pereda, J.M.; Wegener, K.; Santelli, E.; Bate, N.; Ginsberg, M.H.; Critchley, D.R.; Campbell, I.D.; Liddington, R.C.
Deposited on : 2004-11-17
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	14	
1	C	14	
1	E	14	
1	G	14	
1	I	14	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	14	50%29%21%
2	B	202	3% 62%30%5%
2	D	202	4% 62%30%5%
2	F	202	4% 63%30%5%
2	H	202	2% 62%31%5%
2	J	202	2% 68%25%5%
2	L	202	% 63%28%5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10348 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol-4-phosphate 5-kinase, type 1 gamma.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	11	Total	C	N	O	0	0	0
			93	63	14	16			
1	C	11	Total	C	N	O	0	0	0
			93	63	14	16			
1	E	11	Total	C	N	O	0	0	0
			93	63	14	16			
1	G	11	Total	C	N	O	0	0	0
			93	63	14	16			
1	I	11	Total	C	N	O	0	0	0
			93	63	14	16			
1	K	11	Total	C	N	O	0	0	0
			93	63	14	16			

- Molecule 2 is a protein called Talin 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	192	Total	C	N	O	S	0	0	0
			1554	1000	265	282	7			
2	D	192	Total	C	N	O	S	0	0	0
			1554	1000	265	282	7			
2	F	192	Total	C	N	O	S	0	0	0
			1554	1000	265	282	7			
2	H	192	Total	C	N	O	S	0	0	0
			1554	1000	265	282	7			
2	J	192	Total	C	N	O	S	0	0	0
			1554	1000	265	282	7			
2	L	192	Total	C	N	O	S	0	0	0
			1554	1000	265	282	7			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	14	Total 14	O 14	0	0
3	B	72	Total 72	O 72	0	0
3	C	5	Total 5	O 5	0	0
3	D	81	Total 81	O 81	0	0
3	E	7	Total 7	O 7	0	0
3	F	71	Total 71	O 71	0	0
3	G	8	Total 8	O 8	0	0
3	H	74	Total 74	O 74	0	0
3	I	6	Total 6	O 6	0	0
3	J	73	Total 73	O 73	0	0
3	K	10	Total 10	O 10	0	0
3	L	45	Total 45	O 45	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol-4-phosphate 5-kinase, type 1 gamma

Chain A: 




- Molecule 1: Phosphatidylinositol-4-phosphate 5-kinase, type 1 gamma

Chain C: 



- Molecule 1: Phosphatidylinositol-4-phosphate 5-kinase, type 1 gamma

Chain E: 




- Molecule 1: Phosphatidylinositol-4-phosphate 5-kinase, type 1 gamma

Chain G: 




- Molecule 1: Phosphatidylinositol-4-phosphate 5-kinase, type 1 gamma

Chain I: 




- Molecule 1: Phosphatidylinositol-4-phosphate 5-kinase, type 1 gamma


Chain K: 




• Molecule 2: Talin 1




• Molecule 2: Talin 1



• Molecule 2: Talin 1




• Molecule 2: Talin 1




• Molecule 2: Talin 1

Chain J:  2% 68% 25% • 5%



• Molecule 2: Talin 1

Chain L:  63% 28% • 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	118.10Å 118.10Å 93.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.60 30.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	96.5 (30.00-2.60) 96.4 (30.00-2.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 2.61Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.253 , 0.286 0.254 , 0.288	Depositor DCC
R_{free} test set	2175 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	48.0	Xtriage
Anisotropy	0.337	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 65.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.21$	Xtriage
Estimated twinning fraction	0.106 for -h,-k,l 0.106 for h,-h-k,-l 0.296 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	10348	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/98	0.97	1/135 (0.7%)
1	C	0.34	0/98	0.88	1/135 (0.7%)
1	E	0.32	0/98	0.88	1/135 (0.7%)
1	G	0.34	0/98	0.95	1/135 (0.7%)
1	I	0.34	0/98	0.93	0/135
1	K	0.35	0/98	0.86	0/135
2	B	0.37	0/1586	0.88	4/2129 (0.2%)
2	D	0.37	0/1586	0.87	2/2129 (0.1%)
2	F	0.34	0/1586	0.83	1/2129 (0.0%)
2	H	0.34	0/1586	0.84	2/2129 (0.1%)
2	J	0.36	0/1586	0.84	2/2129 (0.1%)
2	L	0.35	0/1586	0.85	1/2129 (0.0%)
All	All	0.36	0/10104	0.86	16/13584 (0.1%)

There are no bond length outliers.

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	399	ILE	N-CA-C	6.43	116.59	110.42
2	F	321	GLY	N-CA-C	-6.09	107.83	115.08
1	G	642	TRP	N-CA-C	5.92	117.94	110.24
1	A	642	TRP	N-CA-C	5.90	118.56	110.24
2	B	326	VAL	N-CA-C	5.89	112.92	107.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	93	0	81	6	0
1	C	93	0	81	9	0
1	E	93	0	81	6	0
1	G	93	0	81	6	0
1	I	93	0	81	2	0
1	K	93	0	81	6	0
2	B	1554	0	1575	63	0
2	D	1554	0	1575	56	0
2	F	1554	0	1575	45	0
2	H	1554	0	1575	51	0
2	J	1554	0	1575	43	0
2	L	1554	0	1575	48	0
3	A	14	0	0	0	0
3	B	72	0	0	7	0
3	C	5	0	0	0	0
3	D	81	0	0	5	0
3	E	7	0	0	0	0
3	F	71	0	0	1	0
3	G	8	0	0	0	0
3	H	74	0	0	3	0
3	I	6	0	0	0	0
3	J	73	0	0	7	0
3	K	10	0	0	0	0
3	L	45	0	0	4	0
All	All	10348	0	9936	299	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 299 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:306:LYS:HG2	2:D:347:VAL:HG21	1.35	1.07
2:B:249:PRO:CG	2:B:288:GLN:HE22	1.72	1.00
2:B:288:GLN:HA	2:B:288:GLN:NE2	1.77	0.97
2:B:288:GLN:HA	2:B:288:GLN:HE21	1.29	0.94
2:B:288:GLN:HE21	2:B:288:GLN:CA	1.76	0.92

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	9/14 (64%)	9 (100%)	0	0	100	100
1	C	9/14 (64%)	9 (100%)	0	0	100	100
1	E	9/14 (64%)	8 (89%)	0	1 (11%)	0	0
1	G	9/14 (64%)	9 (100%)	0	0	100	100
1	I	9/14 (64%)	8 (89%)	1 (11%)	0	100	100
1	K	9/14 (64%)	8 (89%)	1 (11%)	0	100	100
2	B	190/202 (94%)	177 (93%)	11 (6%)	2 (1%)	11	25
2	D	190/202 (94%)	171 (90%)	16 (8%)	3 (2%)	7	16
2	F	190/202 (94%)	171 (90%)	17 (9%)	2 (1%)	11	25
2	H	190/202 (94%)	174 (92%)	14 (7%)	2 (1%)	11	25
2	J	190/202 (94%)	176 (93%)	13 (7%)	1 (0%)	24	46
2	L	190/202 (94%)	177 (93%)	11 (6%)	2 (1%)	11	25
All	All	1194/1296 (92%)	1097 (92%)	84 (7%)	13 (1%)	11	25

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	320	LYS
1	E	642	TRP
2	F	262	LEU
2	J	373	TYR
2	L	371	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	10/13 (77%)	10 (100%)	0	100	100
1	C	10/13 (77%)	10 (100%)	0	100	100
1	E	10/13 (77%)	10 (100%)	0	100	100
1	G	10/13 (77%)	10 (100%)	0	100	100
1	I	10/13 (77%)	10 (100%)	0	100	100
1	K	10/13 (77%)	10 (100%)	0	100	100
2	B	168/177 (95%)	160 (95%)	8 (5%)	23	47
2	D	168/177 (95%)	162 (96%)	6 (4%)	31	58
2	F	168/177 (95%)	161 (96%)	7 (4%)	26	52
2	H	168/177 (95%)	163 (97%)	5 (3%)	36	64
2	J	168/177 (95%)	160 (95%)	8 (5%)	23	47
2	L	168/177 (95%)	159 (95%)	9 (5%)	20	42
All	All	1068/1140 (94%)	1025 (96%)	43 (4%)	28	55

5 of 43 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	J	253	GLN
2	L	225	ASN
2	J	274	LYS
2	J	387	GLN
2	L	320	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	228	HIS
2	L	250	HIS
2	D	246	GLN
2	D	242	GLN
2	L	288	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	11/14 (78%)	0.18	0 100 100	35, 41, 54, 56	0
1	C	11/14 (78%)	0.31	0 100 100	29, 49, 67, 69	0
1	E	11/14 (78%)	0.54	0 100 100	46, 57, 70, 75	0
1	G	11/14 (78%)	0.66	0 100 100	44, 54, 71, 74	0
1	I	11/14 (78%)	0.42	0 100 100	39, 52, 71, 74	0
1	K	11/14 (78%)	0.40	0 100 100	40, 53, 67, 71	0
2	B	192/202 (95%)	0.23	7 (3%) 46 40	22, 47, 87, 99	0
2	D	192/202 (95%)	0.16	8 (4%) 40 35	26, 46, 84, 96	0
2	F	192/202 (95%)	0.30	8 (4%) 40 35	32, 56, 88, 98	0
2	H	192/202 (95%)	0.40	5 (2%) 57 51	33, 56, 90, 100	0
2	J	192/202 (95%)	0.30	4 (2%) 63 58	32, 53, 89, 100	0
2	L	192/202 (95%)	0.41	3 (1%) 70 66	32, 56, 91, 98	0
All	All	1218/1296 (93%)	0.31	35 (2%) 53 48	22, 53, 89, 100	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	321	GLY	3.2
2	B	372	ASP	3.1
2	L	370	PHE	3.1
2	D	376	GLY	3.0
2	D	371	GLY	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.