



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:22 PM UTC

PDB ID : 4Y1A / pdb_00004y1a
Title : immune complex
Authors : Beringer, D.X.; Vivian, J.P.; Reid, H.H.; Rossjohn, J.
Deposited on : 2015-02-07
Resolution : 4.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1082 (4.20-3.80)
Clashscore	190562	1129 (4.20-3.80)
Ramachandran outliers	187476	1064 (4.20-3.80)
Sidechain outliers	187428	1055 (4.20-3.80)
RSRZ outliers	180081	1082 (4.20-3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	181	
2	B	200	
3	C	16	
4	D	210	
5	E	243	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	2	 100%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6616 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	178	Total	C	N	O	S	0	0	0
			1464	949	238	272	5			

- Molecule 2 is a protein called HLA class II histocompatibility antigen, DRB1-4 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	184	Total	C	N	O	S	0	0	0
			1517	958	266	288	5			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	expression tag	UNP P13760
B	0	SER	-	expression tag	UNP P13760
B	191	THR	-	expression tag	UNP P13760
B	192	GLY	-	expression tag	UNP P13760
B	193	GLY	-	expression tag	UNP P13760
B	194	ASP	-	expression tag	UNP P13760
B	195	ASP	-	expression tag	UNP P13760
B	196	ASP	-	expression tag	UNP P13760
B	197	ASP	-	expression tag	UNP P13760
B	198	LYS	-	expression tag	UNP P13760

- Molecule 3 is a protein called Insulin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	14	Total	C	N	O	0	0	0
			106	66	20	20			

- Molecule 4 is a protein called FS17_alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	205	Total	C	N	O	S	0	0	0
			1594	1005	262	319	8			

- Molecule 5 is a protein called FS17_ beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	240	Total	C	N	O	S	0	0	0
			1907	1207	334	357	9			

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	F	2	Total	C	N	O	0	0	0
			28	16	2	10			


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain




- Molecule 2: HLA class II histocompatibility antigen, DRB1-4 beta chain



- Molecule 3: Insulin



- Molecule 4: FS17_alpha




● Molecule 5: FS17_beta

Chain E:

76%

22%

..



● Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:

100%



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	121.91Å 153.93Å 164.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.29 – 4.00 47.29 – 4.00	Depositor EDS
% Data completeness (in resolution range)	98.0 (47.29-4.00) 98.1 (47.29-4.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 4.00Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.225 , 0.283 0.231 , 0.285	Depositor DCC
R_{free} test set	652 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	81.2	Xtriage
Anisotropy	1.211	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 78.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	6616	wwPDB-VP
Average B, all atoms (Å ²)	94.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/1509	0.72	0/2058
2	B	0.24	0/1557	0.71	1/2115 (0.0%)
3	C	0.22	0/106	0.56	0/139
4	D	0.30	0/1632	0.76	1/2213 (0.0%)
5	E	0.26	0/1960	0.68	0/2668
All	All	0.27	0/6764	0.72	2/9193 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	148	SER	N-CA-C	7.29	121.48	112.58
2	B	33	HIS	CB-CA-C	-5.19	110.17	117.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1464	0	1400	19	0
2	B	1517	0	1427	28	0
3	C	106	0	117	2	0
4	D	1594	0	1519	27	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	E	1907	0	1835	31	0
6	F	28	0	25	0	0
All	All	6616	0	6323	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:VAL:HB	1:A:176:LYS:HE3	1.71	0.71
4:D:203:CYS:SG	4:D:204:ALA:N	2.67	0.68
4:D:205:ASN:O	4:D:208:ASN:ND2	2.26	0.68
5:E:99:THR:HG23	5:E:123:THR:HA	1.76	0.68
4:D:208:ASN:O	4:D:209:ASN:ND2	2.27	0.67
2:B:97:PRO:HB3	2:B:122:PHE:HB3	1.75	0.66
4:D:148:SER:N	4:D:149:ASP:HA	2.11	0.66
1:A:140:ARG:HG2	1:A:142:ASP:H	1.62	0.65
5:E:40:TYR:OH	5:E:107:ARG:NH1	2.30	0.64
5:E:11:ARG:HG2	5:E:19:MET:HE1	1.80	0.64
5:E:146:ILE:HG23	5:E:209:ALA:HB1	1.80	0.64
2:B:4:ARG:HD2	2:B:5:PRO:HD2	1.79	0.64
4:D:115:TYR:HB3	4:D:116:GLY:HA2	1.79	0.63
1:A:107:CYS:HB3	1:A:149:HIS:HB2	1.79	0.62
2:B:180:LEU:HD13	2:B:184:LEU:HD13	1.81	0.62
5:E:133:PRO:HD3	5:E:241:PRO:HB3	1.83	0.61
1:A:123:ARG:NH1	1:A:124:ASN:OD1	2.32	0.61
2:B:2:ASP:OD1	2:B:6:ARG:NH2	2.34	0.61
1:A:118:ASN:HB2	1:A:166:GLU:HG2	1.83	0.60
1:A:87:PRO:HB3	1:A:112:PHE:HB3	1.85	0.59
4:D:185:MET:HE1	5:E:208:SER:HB3	1.86	0.57
1:A:16:PRO:HD2	2:B:6:ARG:HH11	1.69	0.57
4:D:19:SER:HB2	4:D:91:ILE:HB	1.87	0.56
2:B:145:THR:HG22	2:B:158:LEU:H	1.71	0.56
4:D:101:VAL:HG12	4:D:125:ILE:HG12	1.87	0.56
2:B:10:GLN:HB2	2:B:31:PHE:HB2	1.90	0.54
2:B:149:GLN:HG2	2:B:155:PHE:HE2	1.72	0.54
5:E:184:ASP:OD2	5:E:202:SER:OG	2.23	0.54
2:B:19:ASN:HB3	2:B:22:GLU:HB2	1.91	0.53
5:E:141:PRO:HB3	5:E:153:THR:H	1.74	0.53
4:D:203:CYS:O	4:D:205:ASN:N	2.40	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:E:141:PRO:HD3	5:E:154:LEU:HG	1.90	0.52
4:D:201:PHE:HA	4:D:202:ALA:HB3	1.91	0.52
1:A:70:LEU:HD13	2:B:9:GLU:HB2	1.93	0.51
5:E:40:TYR:HB2	5:E:105:ALA:HB3	1.92	0.51
1:A:17:ASP:OD1	2:B:6:ARG:NH1	2.43	0.51
2:B:128:GLU:HB3	2:B:176:GLU:HB3	1.92	0.50
2:B:143:VAL:HG13	2:B:160:MET:HB2	1.93	0.49
2:B:28:ASP:HB3	2:B:40:PHE:HB3	1.95	0.49
1:A:156:SER:OG	1:A:159:ASP:OD2	2.29	0.49
5:E:138:VAL:HG23	5:E:248:ALA:HB3	1.93	0.49
1:A:12:PHE:CE2	1:A:21:GLU:HB2	2.48	0.49
5:E:44:GLN:HB3	5:E:101:VAL:HG13	1.95	0.49
2:B:46:GLU:OE1	2:B:48:ARG:NH1	2.47	0.48
2:B:86:GLY:O	2:B:90:THR:OG1	2.25	0.48
5:E:43:ARG:HB2	5:E:53:ILE:HD11	1.95	0.48
2:B:62:ASN:HA	2:B:68:LEU:HD11	1.94	0.48
5:E:110:ASP:HB3	5:E:111:PRO:HD3	1.96	0.48
5:E:141:PRO:HG2	5:E:152:ALA:HB1	1.96	0.48
2:B:99:VAL:HG21	2:B:175:VAL:HG21	1.95	0.47
5:E:38:TYR:CE1	5:E:57:VAL:HG23	2.49	0.47
5:E:3:GLY:HA2	5:E:27:MET:HE1	1.94	0.47
5:E:136:VAL:HG13	5:E:248:ALA:HB2	1.95	0.47
4:D:208:ASN:C	4:D:210:SER:H	2.21	0.47
2:B:168:GLY:HA2	2:B:190:ALA:HB3	1.97	0.47
1:A:26:PHE:HB2	1:A:31:ILE:HD11	1.96	0.47
5:E:6:GLN:HG3	5:E:118:PRO:HD2	1.97	0.47
5:E:234:TRP:NE1	5:E:236:GLN:HB2	2.30	0.46
2:B:68:LEU:O	2:B:72:ARG:HB2	2.16	0.46
4:D:207:PHE:HB3	4:D:212:ILE:HD11	1.99	0.44
4:D:96:PRO:HA	4:D:128:VAL:HB	1.99	0.44
5:E:191:GLN:HB3	5:E:194:LEU:HD13	1.99	0.44
5:E:134:PRO:HB3	5:E:161:PHE:HB3	1.99	0.44
4:D:180:LEU:HB3	5:E:182:CYS:HB2	1.98	0.44
5:E:43:ARG:NH1	5:E:76:TYR:OH	2.51	0.44
4:D:43:LYS:HB3	4:D:53:LEU:HD11	1.99	0.44
4:D:44:LYS:O	4:D:100:ALA:HB1	2.18	0.44
5:E:54:HIS:HB2	5:E:78:VAL:HG11	1.99	0.44
1:A:124:ASN:ND2	1:A:160:VAL:H	2.16	0.44
4:D:92:VAL:HA	4:D:93:PRO:HA	1.74	0.43
4:D:147:SER:C	4:D:149:ASP:HA	2.43	0.43
1:A:99:LEU:HA	1:A:155:PRO:HB2	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:8:LEU:H	2:B:33:HIS:CE1	2.37	0.43
2:B:127:ILE:HG22	2:B:177:HIS:HD2	1.83	0.43
2:B:102:TYR:HA	2:B:103:PRO:HD3	1.89	0.43
4:D:107:SER:HB2	4:D:118:LEU:HD23	2.00	0.43
4:D:117:LYS:HZ3	5:E:52:LEU:HD22	1.84	0.43
2:B:67:LEU:HD11	3:C:7:LEU:HD22	2.01	0.42
2:B:116:VAL:HG22	2:B:160:MET:HG2	2.01	0.42
2:B:117:CYS:HB2	2:B:131:TRP:CZ2	2.54	0.42
2:B:162:GLU:OE1	2:B:162:GLU:N	2.48	0.42
1:A:126:LYS:HA	1:A:127:PRO:HD3	1.88	0.42
4:D:175:THR:HA	5:E:188:LEU:HD11	2.02	0.42
4:D:139:VAL:HG11	4:D:204:ALA:HA	2.02	0.41
4:D:18:ILE:HG12	4:D:92:VAL:HG22	2.02	0.41
1:A:167:HIS:CD2	1:A:168:TRP:H	2.38	0.41
5:E:59:GLU:HA	5:E:60:GLY:HA2	1.65	0.41
4:D:178:CYS:HB3	5:E:204:ARG:HH11	1.86	0.41
5:E:79:SER:OG	5:E:88:LEU:HB2	2.20	0.41
1:A:119:VAL:HG22	1:A:165:VAL:HG22	2.02	0.41
5:E:42:TYR:HD2	5:E:50:LEU:HD23	1.86	0.41
1:A:89:VAL:HG22	1:A:109:ILE:HA	2.03	0.41
2:B:85:VAL:HG13	3:C:-1:GLN:HB2	2.02	0.41
4:D:77:THR:HB	4:D:90:HIS:HB2	2.04	0.41
1:A:64:ALA:HB1	4:D:113:THR:HG21	2.02	0.40
4:D:114:SER:O	4:D:115:TYR:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	176/181 (97%)	168 (96%)	8 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	180/200 (90%)	172 (96%)	8 (4%)	0	100	100
3	C	12/16 (75%)	11 (92%)	1 (8%)	0	100	100
4	D	203/210 (97%)	183 (90%)	19 (9%)	1 (0%)	24	60
5	E	238/243 (98%)	224 (94%)	14 (6%)	0	100	100
All	All	809/850 (95%)	758 (94%)	50 (6%)	1 (0%)	48	81

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	74	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/166 (98%)	158 (97%)	5 (3%)	35	56
2	B	165/178 (93%)	161 (98%)	4 (2%)	43	63
3	C	11/12 (92%)	11 (100%)	0	100	100
4	D	183/189 (97%)	176 (96%)	7 (4%)	29	51
5	E	207/210 (99%)	202 (98%)	5 (2%)	43	63
All	All	729/755 (97%)	708 (97%)	21 (3%)	37	58

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	GLU
1	A	116	VAL
1	A	122	LEU
1	A	158	GLU
1	A	160	VAL
2	B	113	ASN
2	B	143	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	170	VAL
2	B	181	THR
4	D	44	LYS
4	D	64	ASP
4	D	78	VAL
4	D	113	THR
4	D	142	LEU
4	D	205	ASN
4	D	209	ASN
5	E	59	GLU
5	E	71	VAL
5	E	78	VAL
5	E	112	VAL
5	E	166	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
4	D	131	ASN
4	D	205	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	F	1	1,6	14,14,15	0.34	0	17,19,21	0.59	0
6	NAG	F	2	6	14,14,15	0.21	0	17,19,21	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	F	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	F	2	6	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	F	1	NAG	O5-C5-C6-O6
6	F	1	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	178/181 (98%)	0.39	0 100 100	60, 108, 154, 180	0
2	B	184/200 (92%)	0.38	3 (1%) 70 52	61, 85, 136, 171	0
3	C	14/16 (87%)	0.51	0 100 100	70, 79, 110, 119	0
4	D	205/210 (97%)	0.50	5 (2%) 59 44	43, 95, 148, 212	0
5	E	240/243 (98%)	0.51	1 (0%) 88 75	55, 80, 108, 132	0
All	All	821/850 (96%)	0.45	9 (1%) 78 60	43, 90, 141, 212	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	114	SER	2.7
2	B	104	ALA	2.5
4	D	84	ALA	2.5
4	D	108	SER	2.5
2	B	191	THR	2.4
4	D	15	GLU	2.4
2	B	59	GLU	2.3
4	D	36	PHE	2.3
5	E	152	ALA	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	F	2	14/15	0.69	0.14	138,147,154,155	0
6	NAG	F	1	14/15	0.85	0.11	125,134,140,145	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.