



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:45 AM UTC

PDB ID : 2Y2I / pdb_00002y2i
Title : PENICILLIN-BINDING PROTEIN 1B (PBP-1B) IN COMPLEX WITH AN ALKYL BORONATE (ZA3)
Authors : Contreras-Martel, C.; Amoroso, A.; Woon, E.C.; Zervosen, A.; Inglis, S.; Martins, A.; Verlaine, O.; Rydzik, A.; Job, V.; Luxen, A.; Joris, B.; Schofield, C.J.; Dessen, A.
Deposited on : 2010-12-15
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	494	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3983 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PENICILLIN-BINDING PROTEIN 1B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	467	3596	2252	609	720	15	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	656	GLY	ASN	engineered mutation	UNP Q7CRA4
A	686	GLN	ARG	engineered mutation	UNP Q7CRA4
A	687	GLN	ARG	engineered mutation	UNP Q7CRA4

- Molecule 2 is TRIHYDROXY-[[[(2-NITROPHENYL)CARBONYLAMINO]METHYL]BORON (CCD ID: ZA3) (formula: C₈H₁₀BN₂O₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	B	C	N	O		
2	A	1	16	1	8	2	5	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	11	Total	Cl	0	0
			11	11		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	354	Total	O	0	0
			354	354		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PENICILLIN-BINDING PROTEIN 1B



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	97.64Å 148.77Å 97.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.82 – 1.78 48.82 – 1.78	Depositor EDS
% Data completeness (in resolution range)	96.6 (48.82-1.78) 96.6 (48.82-1.78)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 1.78Å)	Xtriage
Refinement program	REFMAC 5.6.0095	Depositor
R, R_{free}	0.179 , 0.213 0.179 , 0.213	Depositor DCC
R_{free} test set	6669 reflections (10.14%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3983	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZA3, NA, CL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.02	2/3669 (0.1%)	0.97	2/4980 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	758	ALA	CA-CB	5.15	1.60	1.53
1	A	655	THR	CA-CB	5.09	1.61	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	748	VAL	N-CA-C	5.16	115.78	107.98
1	A	625	SER	CA-CB-OG	-5.10	100.91	111.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3596	0	3465	61	0
2	A	16	0	9	5	0
3	A	5	0	0	0	0
4	A	11	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	1	0	0	0	0
6	A	354	0	0	11	0
All	All	3983	0	3474	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:444:GLN:HG2	6:A:2096:HOH:O	1.43	1.14
1:A:497:MET:HE1	1:A:502:LYS:HG2	1.22	1.07
1:A:495:PRO:HB2	1:A:497:MET:HE3	1.36	1.02
1:A:659:GLU:HG2	6:A:2267:HOH:O	1.63	0.96
1:A:657:GLN:HB2	1:A:659:GLU:OE2	1.65	0.95
1:A:376:ARG:HD3	6:A:2041:HOH:O	1.68	0.92
1:A:497:MET:HE1	1:A:502:LYS:CG	2.01	0.91
1:A:362:SER:HG	1:A:365:GLU:HG3	1.38	0.89
1:A:397:LYS:HD3	6:A:2300:HOH:O	1.71	0.87
1:A:362:SER:OG	1:A:365:GLU:HG3	1.75	0.86
1:A:486:TYR:CE1	1:A:506:MET:HE2	2.11	0.85
1:A:108:ILE:CD1	1:A:390:ILE:HD12	2.12	0.78
1:A:633:LYS:HG2	4:A:1202:CL:CL	2.24	0.74
1:A:657:GLN:HB2	1:A:659:GLU:CD	2.14	0.72
1:A:486:TYR:HE1	1:A:506:MET:HE2	1.57	0.68
1:A:645:ASN:O	1:A:716:ARG:NH2	2.27	0.67
1:A:108:ILE:N	1:A:108:ILE:HD13	2.09	0.67
1:A:486:TYR:CD1	1:A:506:MET:HE2	2.31	0.65
1:A:741:VAL:O	1:A:742:GLU:C	2.42	0.62
1:A:687:GLN:NE2	6:A:2284:HOH:O	2.30	0.62
1:A:780:GLN:HE21	1:A:780:GLN:C	2.09	0.60
1:A:108:ILE:HD11	1:A:390:ILE:HD12	1.83	0.59
1:A:497:MET:CE	1:A:502:LYS:HG2	2.15	0.59
1:A:456:ARG:HD3	6:A:2271:HOH:O	2.01	0.59
1:A:489:ASN:HD22	1:A:495:PRO:HA	1.68	0.59
1:A:783:TRP:O	1:A:787:VAL:HG23	2.02	0.58
1:A:657:GLN:O	1:A:658:ASP:HB2	2.02	0.58
1:A:108:ILE:HD12	1:A:390:ILE:HD12	1.82	0.58
1:A:364:LYS:HA	1:A:367:LYS:HD2	1.85	0.58
1:A:735:LYS:HE2	4:A:1208:CL:CL	2.42	0.56
1:A:430:GLN:OE1	6:A:2082:HOH:O	2.18	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:495:PRO:HB2	1:A:497:MET:CE	2.25	0.55
1:A:107:GLU:C	1:A:108:ILE:HD13	2.31	0.55
1:A:659:GLU:CG	6:A:2267:HOH:O	2.36	0.55
1:A:108:ILE:HD12	1:A:390:ILE:HB	1.88	0.54
1:A:556:MET:HE2	2:A:1000:ZA3:O16	2.08	0.54
1:A:108:ILE:HG12	1:A:119:ILE:HD11	1.89	0.54
1:A:495:PRO:CB	1:A:497:MET:HE3	2.24	0.53
1:A:518:ASN:OD1	2:A:1000:ZA3:H32C	2.08	0.52
1:A:654:THR:O	2:A:1000:ZA3:O1	2.23	0.51
1:A:105:ILE:HG12	1:A:105:ILE:O	2.10	0.50
1:A:744:LYS:HB3	1:A:746:VAL:HG23	1.94	0.49
1:A:554:LEU:N	1:A:555:PRO:HD2	2.27	0.49
1:A:610:LYS:HE2	6:A:2231:HOH:O	2.12	0.48
1:A:456:ARG:HD2	1:A:660:ASN:OD1	2.15	0.47
1:A:743:GLY:O	1:A:744:LYS:C	2.57	0.46
1:A:362:SER:HG	1:A:365:GLU:H	1.64	0.46
1:A:556:MET:CE	2:A:1000:ZA3:O16	2.64	0.46
1:A:768:TYR:CE1	1:A:783:TRP:CD1	3.04	0.45
1:A:622:GLU:HA	1:A:625:SER:HB2	1.98	0.45
1:A:730:LYS:HE3	1:A:730:LYS:HB2	1.65	0.45
1:A:362:SER:OG	1:A:364:LYS:HG3	2.17	0.44
1:A:354:TYR:CG	1:A:597:VAL:HG13	2.53	0.44
1:A:556:MET:HB3	2:A:1000:ZA3:C7	2.48	0.44
1:A:444:GLN:CG	6:A:2096:HOH:O	2.28	0.43
1:A:738:LYS:HA	1:A:746:VAL:O	2.20	0.42
1:A:382:GLU:HG3	1:A:388:TYR:CE2	2.54	0.42
1:A:692:ASN:HD22	1:A:692:ASN:H	1.67	0.41
1:A:444:GLN:NE2	6:A:2096:HOH:O	2.32	0.41
1:A:720:ASP:CG	1:A:721:PRO:HD2	2.46	0.41
1:A:108:ILE:HD12	1:A:390:ILE:CD1	2.48	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	463/494 (94%)	451 (97%)	10 (2%)	2 (0%)	30	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	742	GLU
1	A	744	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	382/408 (94%)	364 (95%)	18 (5%)	23	6

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	ILE
1	A	108	ILE
1	A	364	LYS
1	A	389	LYS
1	A	500	ASN
1	A	569	ASN
1	A	610	LYS
1	A	692	ASN
1	A	701	VAL
1	A	738	LYS
1	A	740	SER
1	A	742	GLU
1	A	744	LYS
1	A	745	GLU
1	A	746	VAL
1	A	749	THR
1	A	769	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	780	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	424	ASN
1	A	444	GLN
1	A	448	ASN
1	A	489	ASN
1	A	514	ASN
1	A	635	ASN
1	A	692	ASN
1	A	695	ASN
1	A	780	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 12 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ZA3	A	1000	5,1	16,16,17	4.12	3 (18%)	13,21,24	1.28	1 (7%)
3	SO4	A	1100	-	4,4,4	0.82	0	6,6,6	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ZA3	A	1000	5,1	-	0/8/13/14	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	ZA3	O17-N15	14.30	1.47	1.22
2	A	1000	ZA3	C3-N4	-7.08	1.36	1.45
2	A	1000	ZA3	C5-N4	2.04	1.38	1.33

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	ZA3	O17-N15-C14	3.09	124.32	119.03

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1000	ZA3	5	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	467/494 (94%)	0.15	21 (4%) 38 44	19, 33, 70, 127	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	105	ILE	7.1
1	A	788	GLY	6.4
1	A	744	LYS	4.3
1	A	741	VAL	3.7
1	A	787	VAL	3.6
1	A	746	VAL	3.2
1	A	387	GLY	3.1
1	A	363	ALA	2.9
1	A	743	GLY	2.9
1	A	739	VAL	2.8
1	A	117	ALA	2.8
1	A	786	ILE	2.8
1	A	106	SER	2.7
1	A	108	ILE	2.7
1	A	749	THR	2.5
1	A	366	LEU	2.4
1	A	119	ILE	2.3
1	A	118	SER	2.3
1	A	745	GLU	2.1
1	A	499	ALA	2.0
1	A	364	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZA3	A	1000	16/17	0.81	0.16	29,50,69,82	0
4	CL	A	1208	1/1	0.89	0.17	60,60,60,60	0
4	CL	A	1203	1/1	0.91	0.17	57,57,57,57	0
4	CL	A	1202	1/1	0.93	0.14	49,49,49,49	0
4	CL	A	1792	1/1	0.95	0.10	49,49,49,49	1
5	NA	A	1300	1/1	0.95	0.09	32,32,32,32	0
4	CL	A	1206	1/1	0.97	0.16	41,41,41,41	0
4	CL	A	1201	1/1	0.97	0.09	33,33,33,33	0
3	SO4	A	1100	5/5	0.97	0.16	33,35,37,40	0
4	CL	A	1200	1/1	0.97	0.09	35,35,35,35	0
4	CL	A	1213	1/1	0.98	0.20	40,40,40,40	0
4	CL	A	1791	1/1	0.99	0.03	25,25,25,25	1
4	CL	A	1212	1/1	0.99	0.09	32,32,32,32	0
4	CL	A	1211	1/1	0.99	0.08	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.