



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:41 AM UTC

PDB ID : 5Y39 / pdb_00005y39
Title : Crystal structure of Ragulator complex (p18 76-145)
Authors : Zhang, T.; Ding, J.
Deposited on : 2017-07-28
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	70	
1	F	70	
2	B	125	
2	G	125	
3	C	124	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	124	6% 78% 14% 6%
4	D	101	4% 75% 19%
4	I	101	35% 52% 7% 41%
5	E	90	2% 82% 14%
5	J	90	7% 82% 10% 7%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6808 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ragulator complex protein LAMTOR1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	67	Total	C	N	O	S	0	0	0
			534	340	94	99	1			
1	F	32	Total	C	N	O		0	0	0
			227	146	37	44				

- Molecule 2 is a protein called Ragulator complex protein LAMTOR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	116	Total	C	N	O	S	0	0	0
			878	552	149	171	6			
2	G	117	Total	C	N	O	S	0	0	0
			872	547	149	170	6			

- Molecule 3 is a protein called Ragulator complex protein LAMTOR3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	121	Total	C	N	O	S	0	0	0
			941	606	158	176	1			
3	H	117	Total	C	N	O	S	0	0	0
			893	575	149	168	1			

- Molecule 4 is a protein called Ragulator complex protein LAMTOR4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	99	Total	C	N	O	S	0	0	0
			747	465	136	143	3			
4	I	60	Total	C	N	O	S	0	0	0
			380	239	70	70	1			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-1	MET	-	expression tag	UNP Q0VGL1
D	0	GLY	-	expression tag	UNP Q0VGL1
I	-1	MET	-	expression tag	UNP Q0VGL1
I	0	GLY	-	expression tag	UNP Q0VGL1

- Molecule 5 is a protein called Regulator complex protein LAMTOR5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	90	Total	C	N	O	S	0	0	0
			660	403	114	136	7			
5	J	84	Total	C	N	O	S	0	0	0
			618	379	106	126	7			


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	9	Total	O	0	0
			9	9		
6	B	15	Total	O	0	0
			15	15		
6	C	14	Total	O	0	0
			14	14		
6	D	7	Total	O	0	0
			7	7		
6	E	6	Total	O	0	0
			6	6		
6	G	2	Total	O	0	0
			2	2		
6	I	1	Total	O	0	0
			1	1		
6	J	4	Total	O	0	0
			4	4		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Regulator complex protein LAMTOR1





- Molecule 1: Regulator complex protein LAMTOR1




- Molecule 2: Regulator complex protein LAMTOR2




- Molecule 2: Regulator complex protein LAMTOR2




- Molecule 3: Regulator complex protein LAMTOR3




- Molecule 3: Regulator complex protein LAMTOR3




- Molecule 4: Regulator complex protein LAMTOR4





- Molecule 4: Regulator complex protein LAMTOR4



- Molecule 5: Regulator complex protein LAMTOR5



- Molecule 5: Regulator complex protein LAMTOR5



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	110.66Å 117.85Å 157.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.01 – 2.65 50.01 – 2.65	Depositor EDS
% Data completeness (in resolution range)	88.1 (50.01-2.65) 88.1 (50.01-2.65)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.83 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, R_{free}	0.216 , 0.263 0.216 , 0.263	Depositor DCC
R_{free} test set	1365 reflections (4.50%)	wwPDB-VP
Wilson B-factor (Å ²)	38.4	Xtriage
Anisotropy	0.257	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 67.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6808	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.45	0/549	0.98	2/749 (0.3%)
1	F	0.46	0/233	1.01	0/321
2	B	0.42	0/888	0.89	0/1202
2	G	0.43	0/881	0.95	2/1195 (0.2%)
3	C	0.49	0/958	0.86	0/1300
3	H	0.44	0/910	0.89	1/1238 (0.1%)
4	D	0.48	0/756	0.87	0/1020
4	I	0.46	0/376	0.86	0/502
5	E	0.44	0/666	0.90	0/903
5	J	0.43	0/621	0.87	0/839
All	All	0.45	0/6838	0.90	5/9269 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	65	LEU	N-CA-C	-6.63	98.89	109.76
1	A	106	PRO	CA-C-N	5.93	127.25	119.84
1	A	106	PRO	C-N-CA	5.93	127.25	119.84
2	G	79	GLY	N-CA-C	5.79	117.85	110.96
2	G	36	GLY	N-CA-C	5.48	118.76	110.75

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	534	0	531	9	0
1	F	227	0	204	3	0
2	B	878	0	880	7	0
2	G	872	0	869	9	0
3	C	941	0	967	8	0
3	H	893	0	897	7	0
4	D	747	0	756	12	0
4	I	380	0	333	2	0
5	E	660	0	658	9	0
5	J	618	0	615	4	0
6	A	9	0	0	0	0
6	B	15	0	0	0	0
6	C	14	0	0	0	0
6	D	7	0	0	0	0
6	E	6	0	0	0	0
6	G	2	0	0	0	0
6	I	1	0	0	0	0
6	J	4	0	0	0	0
All	All	6808	0	6710	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:5:LEU:HD11	3:C:113:LEU:HD11	1.52	0.90
3:C:9:LEU:HB3	3:C:35:VAL:HG21	1.61	0.81
1:A:102:TRP:H	1:A:102:TRP:CD1	2.12	0.66
5:E:139:SER:O	5:E:141:PRO:HD3	1.98	0.64
2:B:77:MET:HB2	3:C:66:SER:HB3	1.82	0.60
2:B:93:MET:HE1	2:B:107:LYS:CB	2.32	0.59
4:I:74:THR:HG22	4:I:88:LYS:HD2	1.87	0.56
1:A:143:LEU:HD13	5:E:164:ILE:HG21	1.88	0.56
2:B:93:MET:HE1	2:B:107:LYS:HB2	1.90	0.54
2:G:93:MET:HE1	2:G:107:LYS:HB3	1.88	0.53
5:J:88:GLU:HG3	5:J:116:ARG:HH12	1.73	0.53
2:B:93:MET:HE2	2:B:104:LEU:HD12	1.90	0.53
4:D:95:GLU:CD	4:D:95:GLU:H	2.18	0.51
1:A:124:ILE:HD11	5:E:150:GLU:HG2	1.93	0.51
2:G:69:LEU:HD21	2:G:72:ILE:HD11	1.92	0.51
1:F:131:GLN:O	1:F:135:ILE:HG22	2.12	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:81:VAL:HG21	3:C:63:LEU:HD13	1.94	0.50
2:B:93:MET:HE1	2:B:107:LYS:HB3	1.92	0.50
5:E:152:ASP:N	5:E:152:ASP:OD1	2.44	0.49
4:D:43:ILE:HG13	5:E:137:LEU:HD21	1.93	0.49
1:F:143:LEU:C	1:F:145:GLN:H	2.20	0.49
3:H:16:VAL:HG22	3:H:102:LEU:HD13	1.95	0.49
4:D:65:ARG:HD2	5:E:150:GLU:OE2	2.14	0.48
1:F:143:LEU:HD22	5:J:164:ILE:HD12	1.94	0.48
5:J:160:LYS:HA	5:J:165:THR:HA	1.96	0.48
4:D:2:THR:O	4:D:6:THR:HB	2.14	0.48
4:D:43:ILE:HD13	4:D:86:VAL:HG11	1.96	0.47
2:G:67:ASP:OD1	3:H:76:ASN:HB2	2.15	0.46
1:A:103:LYS:HG3	3:C:4:ASP:HB3	1.97	0.46
3:H:66:SER:O	3:H:67:LYS:C	2.59	0.45
2:G:93:MET:HE2	2:G:104:LEU:HD12	1.98	0.45
1:A:134:ARG:O	1:A:138:TYR:HD1	2.00	0.44
3:H:87:PRO:HD2	3:H:118:ARG:HD2	2.00	0.44
3:C:30:VAL:HA	3:C:31:PRO:HD3	1.85	0.44
2:G:53:TRP:HE1	2:G:85:ARG:NH2	2.15	0.44
4:D:19:LEU:HB3	4:D:86:VAL:HB	1.99	0.44
1:A:126:PHE:HA	1:A:129:LEU:HD12	2.01	0.43
4:D:19:LEU:HB2	4:D:33:LEU:HD23	2.01	0.43
4:I:68:VAL:HB	4:I:75:LEU:HB2	2.01	0.43
2:G:118:PRO:O	2:G:121:GLN:HB2	2.19	0.43
3:H:110:LEU:HB3	3:H:114:PHE:CE2	2.53	0.43
1:A:102:TRP:CE2	3:C:113:LEU:HD13	2.54	0.43
2:G:93:MET:HE1	2:G:107:LYS:CB	2.49	0.42
5:E:160:LYS:HG2	5:E:161:HIS:O	2.19	0.42
4:D:16:LEU:HB2	4:D:88:LYS:HD3	2.01	0.42
5:J:95:MET:HG3	5:J:116:ARG:HB3	2.02	0.42
1:A:111:LEU:HG	4:D:63:PHE:O	2.19	0.42
4:D:79:VAL:HG22	4:D:84:VAL:HG22	2.01	0.41
5:E:97:ASN:HA	5:E:98:PRO:HD3	1.94	0.41
2:G:24:LEU:HD11	2:G:89:LEU:HD13	2.01	0.41
4:D:9:LEU:O	4:D:12:ILE:HG12	2.20	0.41
4:D:69:VAL:HA	4:D:74:THR:HB	2.02	0.41
3:H:13:LEU:HA	3:H:19:LEU:HD23	2.02	0.41
1:A:143:LEU:HD11	5:E:113:LEU:HD13	2.03	0.41
2:G:63:ALA:HB1	3:H:44:ALA:HB2	2.03	0.41
2:B:86:VAL:HB	2:B:112:VAL:HG22	2.03	0.40
3:C:5:LEU:HD11	3:C:113:LEU:CD1	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	65/70 (93%)	61 (94%)	4 (6%)	0	100	100
1	F	30/70 (43%)	28 (93%)	2 (7%)	0	100	100
2	B	114/125 (91%)	110 (96%)	4 (4%)	0	100	100
2	G	115/125 (92%)	108 (94%)	7 (6%)	0	100	100
3	C	119/124 (96%)	117 (98%)	2 (2%)	0	100	100
3	H	115/124 (93%)	111 (96%)	4 (4%)	0	100	100
4	D	97/101 (96%)	93 (96%)	4 (4%)	0	100	100
4	I	46/101 (46%)	42 (91%)	4 (9%)	0	100	100
5	E	88/90 (98%)	85 (97%)	3 (3%)	0	100	100
5	J	78/90 (87%)	74 (95%)	4 (5%)	0	100	100
All	All	867/1020 (85%)	829 (96%)	38 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	60/63 (95%)	54 (90%)	6 (10%)	7	11
1	F	22/63 (35%)	20 (91%)	2 (9%)	9	14

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	91/98 (93%)	81 (89%)	10 (11%)	6	9
2	G	89/98 (91%)	80 (90%)	9 (10%)	7	11
3	C	106/108 (98%)	99 (93%)	7 (7%)	15	26
3	H	98/108 (91%)	88 (90%)	10 (10%)	7	10
4	D	81/84 (96%)	71 (88%)	10 (12%)	4	7
4	I	29/84 (34%)	26 (90%)	3 (10%)	7	10
5	E	76/76 (100%)	68 (90%)	8 (10%)	6	10
5	J	71/76 (93%)	66 (93%)	5 (7%)	14	24
All	All	723/858 (84%)	653 (90%)	70 (10%)	8	12

All (70) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	THR
1	A	102	TRP
1	A	111	LEU
1	A	124	ILE
1	A	143	LEU
1	A	144	SER
2	B	7	LEU
2	B	25	LEU
2	B	31	LEU
2	B	40	THR
2	B	58	ARG
2	B	91	LEU
2	B	97	GLU
2	B	109	GLN
2	B	113	GLN
2	B	116	GLU
3	C	9	LEU
3	C	11	LYS
3	C	33	ILE
3	C	67	LYS
3	C	102	LEU
3	C	110	LEU
3	C	116	GLU
4	D	6	THR
4	D	20	VAL
4	D	37	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	38	GLN
4	D	46	LEU
4	D	55	LEU
4	D	65	ARG
4	D	69	VAL
4	D	74	THR
4	D	94	ARG
5	E	91	LEU
5	E	92	GLU
5	E	103	VAL
5	E	152	ASP
5	E	153	ASN
5	E	161	HIS
5	E	164	ILE
5	E	166	VAL
1	F	118	VAL
1	F	135	ILE
2	G	7	LEU
2	G	9	GLN
2	G	24	LEU
2	G	31	LEU
2	G	68	ASN
2	G	90	LEU
2	G	91	LEU
2	G	97	GLU
2	G	120	THR
3	H	33	ILE
3	H	57	THR
3	H	91	SER
3	H	93	ILE
3	H	102	LEU
3	H	108	LYS
3	H	110	LEU
3	H	113	LEU
3	H	117	LEU
3	H	118	ARG
4	I	11	ARG
4	I	69	VAL
4	I	76	LEU
5	J	86	THR
5	J	88	GLU
5	J	91	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
5	J	143	ASP
5	J	147	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	20	GLN
3	C	68	ASN
3	C	76	ASN
2	G	13	GLN
2	G	51	ASN
3	H	59	GLN
3	H	79	GLN
3	H	119	GLN
5	J	109	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	67/70 (95%)	0.80	13 (19%) 3 2	28, 60, 118, 134	0
1	F	32/70 (45%)	1.98	13 (40%) 0 0	68, 103, 198, 206	0
2	B	116/125 (92%)	-0.06	1 (0%) 81 78	22, 30, 56, 86	0
2	G	117/125 (93%)	0.15	3 (2%) 57 51	27, 40, 111, 133	0
3	C	121/124 (97%)	-0.03	2 (1%) 69 64	24, 36, 70, 107	0
3	H	117/124 (94%)	0.58	8 (6%) 23 17	34, 76, 112, 135	0
4	D	99/101 (98%)	0.29	4 (4%) 42 34	25, 45, 109, 151	0
4	I	60/101 (59%)	2.43	35 (58%) 0 0	98, 133, 217, 229	0
5	E	90/90 (100%)	0.01	2 (2%) 62 56	24, 37, 74, 87	0
5	J	84/90 (93%)	0.72	6 (7%) 22 16	39, 73, 119, 136	0
All	All	903/1020 (88%)	0.47	87 (9%) 13 11	22, 50, 133, 229	0

All (87) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	I	7	GLN	6.0
4	I	86	VAL	5.7
4	I	18	TYR	5.3
4	I	10	GLU	4.9
4	I	27	LEU	4.8
4	I	4	ALA	4.4
4	I	52	GLY	4.4
4	I	9	LEU	4.2
1	A	102	TRP	4.1
4	I	11	ARG	4.0
2	G	65	ASN	3.9
4	I	30	SER	3.8
1	F	115	PRO	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	I	81	GLY	3.6
4	I	39	ALA	3.6
1	A	107	PRO	3.6
4	I	79	VAL	3.5
1	A	99	LEU	3.4
1	A	97	SER	3.4
5	J	151	SER	3.4
1	F	145	GLN	3.3
4	D	93	GLY	3.3
5	J	121	ASP	3.2
4	I	72	GLU	3.2
1	F	135	ILE	3.2
1	A	104	LYS	3.1
1	F	114	GLN	3.1
1	F	129	LEU	3.0
3	H	7	ARG	3.0
1	A	106	PRO	3.0
4	I	29	SER	3.0
4	I	85	PHE	3.0
4	I	40	ALA	3.0
4	I	16	LEU	2.9
4	D	59	MET	2.9
3	H	5	LEU	2.9
4	I	64	LYS	2.9
1	A	101	HIS	2.9
1	F	126	PHE	2.8
2	G	63	ALA	2.8
1	F	118	VAL	2.8
4	I	8	GLY	2.8
4	I	5	LEU	2.7
3	H	66	SER	2.7
1	F	122	GLU	2.6
2	G	62	GLN	2.6
1	A	105	LEU	2.6
3	H	86	LEU	2.6
4	D	58	GLY	2.6
3	H	27	ARG	2.6
4	I	20	VAL	2.6
5	J	162	ASP	2.5
1	A	93	ALA	2.5
3	C	3	ASP	2.5
1	A	98	SER	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
5	E	161	HIS	2.5
4	I	38	GLN	2.5
4	I	21	LEU	2.5
5	J	141	PRO	2.5
4	I	42	ALA	2.4
4	I	23	GLU	2.4
3	H	119	GLN	2.4
4	I	84	VAL	2.4
1	F	120	ALA	2.4
4	I	6	THR	2.4
4	I	87	VAL	2.4
1	F	121	SER	2.4
3	H	117	LEU	2.3
1	A	80	GLU	2.3
4	I	88	LYS	2.3
1	F	119	LEU	2.3
1	A	95	LEU	2.3
1	F	116	HIS	2.3
4	I	22	SER	2.3
4	I	28	ALA	2.2
5	E	152	ASP	2.2
1	A	87	GLN	2.2
1	F	117	GLN	2.2
3	C	123	VAL	2.2
3	H	87	PRO	2.1
5	J	108	SER	2.1
2	B	37	TYR	2.1
4	I	19	LEU	2.1
4	I	78	THR	2.1
4	I	53	PHE	2.1
5	J	161	HIS	2.0
4	D	96	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.