



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:45 PM UTC

PDB ID : 6Y39 / pdb_00006y39
Title : HapE-P88L mutant CCAAT-binding complex from *Aspergillus nidulans* with cycA DNA
Authors : Groll, M.; Huber, E.M.
Deposited on : 2020-02-17
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	64	3% 41% 56%
1	D	64	3% 44% 55%
1	G	64	3% 39% 8% 52%
2	B	92	% 90% 8%
2	E	92	90% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	92	2% 91% 5%
3	C	119	2% 95%
3	F	119	3% 92% 6%
3	I	119	% 94% 5%
4	J	25	56% 40%
5	K	25	88% 12%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6811 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HapB.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	28	Total	C	N	O	0	0	0
			243	154	49	40			
1	D	29	Total	C	N	O	0	0	0
			254	160	53	41			
1	G	31	Total	C	N	O	0	0	0
			269	170	55	44			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	230	MET	-	initiating methionine	UNP G5EAZ0
D	230	MET	-	initiating methionine	UNP G5EAZ0
G	230	MET	-	initiating methionine	UNP G5EAZ0

- Molecule 2 is a protein called Transcription factor HapC (Eurofung).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	90	Total	C	N	O	S	0	0	0
			725	458	120	141	6			
2	E	87	Total	C	N	O	S	0	0	0
			695	442	114	133	6			
2	H	87	Total	C	N	O	S	0	0	0
			695	442	114	133	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	41	MET	-	initiating methionine	UNP Q5B5Z6
E	41	MET	-	initiating methionine	UNP Q5B5Z6
H	41	MET	-	initiating methionine	UNP Q5B5Z6

- Molecule 3 is a protein called CBFD_NFYB_HMF domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	118	Total	C	N	O	S	0	0	0
			952	604	170	173	5			
3	F	117	Total	C	N	O	S	0	0	0
			940	598	166	171	5			
3	I	118	Total	C	N	O	S	0	0	0
			952	604	170	173	5			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	46	MET	-	initiating methionine	UNP Q5AYY8
C	88	LEU	PRO	engineered mutation	UNP Q5AYY8
F	46	MET	-	initiating methionine	UNP Q5AYY8
F	88	LEU	PRO	engineered mutation	UNP Q5AYY8
I	46	MET	-	initiating methionine	UNP Q5AYY8
I	88	LEU	PRO	engineered mutation	UNP Q5AYY8

- Molecule 4 is a DNA chain called DNA (25 mer).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	J	25	Total	C	N	O	P	0	0	0
			504	243	81	155	25			

- Molecule 5 is a DNA chain called DNA (25 mer).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	K	25	Total	C	N	O	P	0	0	0
			521	247	104	145	25			

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	F	1	Total Cl 1 1	0	0
6	H	1	Total Cl 1 1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	5	Total O 5 5	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	4	Total 4	O 4	0	0
7	C	10	Total 10	O 10	0	0
7	D	2	Total 2	O 2	0	0
7	E	9	Total 9	O 9	0	0
7	F	11	Total 11	O 11	0	0
7	G	1	Total 1	O 1	0	0
7	H	6	Total 6	O 6	0	0
7	I	5	Total 5	O 5	0	0
7	J	2	Total 2	O 2	0	0
7	K	4	Total 4	O 4	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: HapB





- Molecule 1: HapB



- Molecule 1: HapB




- Molecule 2: Transcription factor HapC (Eurofung)




- Molecule 2: Transcription factor HapC (Eurofung)





- Molecule 2: Transcription factor HapC (Eurofung)



- Molecule 3: CBFD_NFYB_HMF domain-containing protein




- Molecule 3: CBFD_NFYB_HMF domain-containing protein





- Molecule 3: CBFD_NFYB_HMF domain-containing protein



- Molecule 4: DNA (25 mer)



- Molecule 5: DNA (25 mer)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.15Å 103.50Å 159.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.30 30.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.0 (30.00-2.30) 97.9 (30.00-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
R, R_{free}	0.221 , 0.254 0.223 , 0.263	Depositor DCC
R_{free} test set	2643 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	55.3	Xtriage
Anisotropy	0.591	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 34.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6811	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	0/246	1.43	0/327
1	D	1.03	0/257	1.44	0/341
1	G	1.01	0/272	1.45	0/362
2	B	1.01	0/735	1.56	0/983
2	E	1.05	0/705	1.57	0/945
2	H	0.99	0/705	1.57	2/945 (0.2%)
3	C	1.04	0/971	1.44	0/1312
3	F	1.04	0/959	1.41	0/1298
3	I	1.04	0/971	1.41	0/1312
4	J	0.52	2/561 (0.4%)	0.95	2/862 (0.2%)
5	K	0.38	0/587	0.78	0/905
All	All	0.96	2/6969 (0.0%)	1.39	4/9592 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	J	21	DT	O3'-P	-5.36	1.53	1.61
4	J	16	DA	O3'-P	-5.03	1.53	1.61

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	53	VAL	CA-C-N	5.55	127.72	120.28
2	H	53	VAL	C-N-CA	5.55	127.72	120.28
4	J	24	DA	C2'-C3'-O3'	5.44	119.66	111.50
4	J	5	DT	C4'-C3'-O3'	5.25	117.88	110.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	243	0	258	2	0
1	D	254	0	271	1	0
1	G	269	0	289	5	0
2	B	725	0	727	4	0
2	E	695	0	702	2	0
2	H	695	0	702	1	0
3	C	952	0	957	4	0
3	F	940	0	944	4	0
3	I	952	0	957	4	0
4	J	504	0	286	6	0
5	K	521	0	281	2	0
6	F	1	0	0	0	0
6	H	1	0	0	0	0
7	A	5	0	0	0	0
7	B	4	0	0	0	0
7	C	10	0	0	0	0
7	D	2	0	0	0	0
7	E	9	0	0	0	0
7	F	11	0	0	0	0
7	G	1	0	0	0	0
7	H	6	0	0	0	0
7	I	5	0	0	0	0
7	J	2	0	0	0	0
7	K	4	0	0	0	0
All	All	6811	0	6374	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:258:LEU:HD23	1:G:259:ARG:N	1.82	0.94
1:G:258:LEU:HD23	1:G:258:LEU:C	2.18	0.68
3:I:109:ALA:HB3	3:I:110:PRO:HD3	1.82	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:64:ASN:OD1	3:C:138:ARG:NH1	2.36	0.59
3:C:109:ALA:HB3	3:C:110:PRO:HD3	1.86	0.58
4:J:15:DG:N2	5:K:14:DA:C2	2.75	0.55
4:J:5:DT:H2''	4:J:6:DC:O5'	2.08	0.53
3:F:109:ALA:HB3	3:F:110:PRO:HD3	1.89	0.53
4:J:3:DC:H2'	4:J:4:DA:C8	2.43	0.52
2:B:45:ASP:HB3	3:C:94:LYS:HB3	1.93	0.51
3:I:162:VAL:O	3:I:164:ARG:NH1	2.44	0.51
1:G:258:LEU:HD23	1:G:259:ARG:CA	2.41	0.51
4:J:6:DC:H2''	4:J:7:DC:H5'	1.94	0.48
3:F:53:ASN:HB2	3:F:61:ARG:HG3	1.96	0.48
2:E:99:ARG:NH2	2:E:106:ASP:OD1	2.47	0.48
1:D:258:LEU:C	1:D:258:LEU:HD23	2.39	0.47
2:B:97:GLU:OE2	2:B:99:ARG:NH1	2.48	0.47
4:J:1:DT:H2'	4:J:2:DT:C6	2.50	0.47
1:A:251:ARG:HG3	2:B:46:ARG:HH22	1.82	0.45
2:E:103:ASN:HA	3:F:106:SER:HB3	1.97	0.45
1:A:258:LEU:HD12	1:A:258:LEU:N	2.32	0.45
3:C:137:ARG:CG	3:I:86:GLN:HG3	2.47	0.44
1:G:260:LEU:C	1:G:260:LEU:HD23	2.43	0.44
2:H:64:ASN:HB2	3:I:138:ARG:HH11	1.84	0.43
5:K:20:DA:H2''	5:K:21:DG:C8	2.54	0.43
3:F:112:LEU:HD23	3:F:112:LEU:HA	1.93	0.42
1:G:236:VAL:O	1:G:237:ASN:C	2.63	0.41
4:J:3:DC:C2'	4:J:4:DA:C8	3.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	26/64 (41%)	24 (92%)	2 (8%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	27/64 (42%)	25 (93%)	2 (7%)	0	100	100
1	G	29/64 (45%)	28 (97%)	1 (3%)	0	100	100
2	B	88/92 (96%)	86 (98%)	2 (2%)	0	100	100
2	E	85/92 (92%)	84 (99%)	1 (1%)	0	100	100
2	H	85/92 (92%)	84 (99%)	1 (1%)	0	100	100
3	C	116/119 (98%)	112 (97%)	4 (3%)	0	100	100
3	F	115/119 (97%)	112 (97%)	3 (3%)	0	100	100
3	I	116/119 (98%)	113 (97%)	3 (3%)	0	100	100
All	All	687/825 (83%)	668 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	26/56 (46%)	26 (100%)	0	100	100
1	D	27/56 (48%)	27 (100%)	0	100	100
1	G	29/56 (52%)	27 (93%)	2 (7%)	14	20
2	B	78/80 (98%)	76 (97%)	2 (3%)	40	59
2	E	75/80 (94%)	74 (99%)	1 (1%)	61	77
2	H	75/80 (94%)	74 (99%)	1 (1%)	61	77
3	C	102/103 (99%)	102 (100%)	0	100	100
3	F	101/103 (98%)	100 (99%)	1 (1%)	68	82
3	I	102/103 (99%)	102 (100%)	0	100	100
All	All	615/717 (86%)	608 (99%)	7 (1%)	65	81

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	44	GLN
2	B	94	CYS
2	E	102	VAL
3	F	87	LEU
1	G	258	LEU
1	G	261	THR
2	H	55	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	E	95	GLN
3	F	74	HIS
3	I	57	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	28/64 (43%)	0.39	2 (7%) 22 24	50, 61, 126, 130	0
1	D	29/64 (45%)	0.51	2 (6%) 23 25	52, 63, 125, 131	0
1	G	31/64 (48%)	0.70	2 (6%) 25 27	56, 66, 135, 141	0
2	B	90/92 (97%)	0.18	1 (1%) 78 79	45, 64, 111, 139	0
2	E	87/92 (94%)	0.31	0 100 100	46, 62, 103, 112	0
2	H	87/92 (94%)	0.33	2 (2%) 61 63	49, 68, 101, 109	0
3	C	118/119 (99%)	0.35	2 (1%) 69 70	43, 66, 101, 124	0
3	F	117/119 (98%)	0.32	4 (3%) 48 50	42, 65, 106, 130	0
3	I	118/119 (99%)	0.38	1 (0%) 82 83	50, 70, 95, 110	0
4	J	25/25 (100%)	0.29	0 100 100	59, 78, 105, 108	0
5	K	25/25 (100%)	0.24	0 100 100	68, 83, 102, 108	0
All	All	755/875 (86%)	0.34	16 (2%) 63 65	42, 67, 107, 141	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	84	ILE	4.3
1	G	258	LEU	3.3
1	A	258	LEU	3.2
3	F	159	ILE	2.9
1	D	258	LEU	2.7
3	C	84	ILE	2.5
2	B	46	ARG	2.4
1	A	254	LEU	2.2
3	F	47	GLY	2.2
2	H	47	TRP	2.2
3	F	89	LEU	2.1
3	C	164	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	231	GLU	2.1
1	D	233	PRO	2.0
3	I	164	ARG	2.0
2	H	44	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	CL	H	201	1/1	0.85	0.15	86,86,86,86	0
6	CL	F	201	1/1	0.92	0.08	89,89,89,89	0

6.5 Other polymers [i](#)

There are no such residues in this entry.