



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 11:49 PM UTC

PDB ID : 9Y31 / pdb_00009y31
EMDB ID : EMD-72430
Title : Symmetry relaxed asymmetric structure of an expansion intermediate of Turnip crinkle virus
Authors : Venkatakrisnan, V.; Braet, S.; Ramesh, R.; Clawson, M.A.; Laremore, T.N.; Wong, S.M.; Anand, G.S.
Deposited on : 2025-09-01
Resolution : 3.62 Å (reported)
Based on initial model : 9Y2Z

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY



The reported resolution of this entry is 3.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	11773 (3.12 - 4.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	351	
1	E	351	
1	H	351	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 433 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Capsid protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	C	20	Total	C	N	O	0	0
			153	97	28	28		
1	E	18	Total	C	N	O	0	0
			136	86	25	25		
1	H	19	Total	C	N	O	0	0
			144	91	26	27		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Capsid protein



- Molecule 1: Capsid protein



LEU
GLY
GLU
ALA
ASP
ALA
ALA
GLY
SER
VAL
GLN
TRP
ALA
GLY
VAL
GLN
GLY
GLU
ARG
GLY
GLN
VAL
LYS
MET
VAL
GLU
GLN
PRO
LYS
GLY
LYS
TRP
GLN
ALA
LEU
ARG
ILE

● Molecule 1: Capsid protein

Chain H: 5% 95%

MET
GLU
ASN
ASP
PRO
ARG
VAL
ARG
LYS
PHE
GLY
SER
ALA
SER
ASP
GLY
ALA
TRP
TRP
ILE
LYS
LYS
TRP
SER
THR
THR
LEU
THR
SER
ARG
GLN
PRO
VAL
GLN
LYS
V65
P73
ARG
VAL
SER

THR
ALA
ARG
ASP
GLY
ILE
THR
ARG
SER
GLY
SER
GLU
LEU
ILE
THR
THR
LEU
LYS
LYS
ASN
THR
ASP
THR
GLU
GLY
PRO
LYS
THR
THR
THR
ALA
VAL
LEU
ASN
PRO
SER
SER
GLN
PRO
GLY
THR
PHE
ASN
GLN
LEU
ILE
LYS
LYS
GLU
ALA
PRO
VAL
ALA
GLN
TYR
GLU
LYS
TYR
ARG
PHE
THR
SER
LEU
ARG
PHE

ARG
TYR
SER
PRO
MET
SER
PRO
SER
THR
THR
GLY
GLY
LYS
VAL
PHE
LEU
ALA
ASP
ARG
ASP
ALA
ALA
THR
GLY
LYS
PRO
PRO
ASN
ASN
ASP
LEU
ALA
SER
LEU
SER
LEU
TYR
ILE
GLY
GLY
CYS
THR
PHE
VAL
SER
SER
VAL
PRO
TRP
THR
GLU
ALA
PHE
ILE
LEU
THR
VAL
VAL
PRO
THR
ASP
SER
THR
ASP
THR
PHE

VAL
ALA
ASP
GLY
ILE
SER
ASP
PRO
LYS
LEU
VAL
GLY
PHE
LYS
MET
ILE
ALA
THR
ALA
THR
TYR
GLY
ALA
LEU
ILE
GLY
TRP
PHE
GLU
HIS
ASP
THR
GLY
PHE
GLN
CYS
HIS
GLY
GLY
ALA
TRP
GLY
ASN
LEU
GLY
VAL
GLY
ASP
LEU
GLY
VAL
GLY
ASP
PHE
VAL
GLY
GLY
THR
LYS
ALA
VAL
GLY
GLY
PHE
GLY
ASP
PHE
GLY
PHE
ALA
GLY
VAL

LYS
ASP
GLY
PRO
ARG
LEU
VAL
SER
TRP
SER
LYS
VAL
GLU
THR
LYS
GLY
THR
ALA
GLY
TRP
GLY
TRP
GLU
HIS
ASP
THR
GLY
PHE
GLN
CYS
HIS
GLY
GLY
ALA
VAL
GLY
GLY
ASP
PHE
GLY
PHE
ALA
GLY
VAL

GLY
SER
VAL
GLN
TRP
ALA
GLY
VAL
LYS
VAL
ALA
GLU
ARG
GLY
GLN
GLY
VAL
LYS
MET
VAL
THR
THR
GLY
GLU
GLU
GLN
PRO
LYS
GLY
LYS
TRP
GLN
ALA
LEU
ARG
ILE

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	60938	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	25	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	40000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON II (4k x 4k)	Depositor
Maximum map value	3.433	Depositor
Minimum map value	-1.687	Depositor
Average map value	-0.005	Depositor
Map value standard deviation	0.177	Depositor
Recommended contour level	0.45	Depositor
Map size (Å)	638.63995, 638.63995, 638.63995	wwPDB
Map dimensions	600, 600, 600	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0644, 1.0644, 1.0644	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.98	0/155	1.14	2/211 (0.9%)
1	E	1.23	0/138	1.08	0/188
1	H	1.05	0/146	0.97	0/200
All	All	1.09	0/439	1.07	2/599 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	58	LEU	CA-C-N	5.14	131.35	121.54
1	C	58	LEU	C-N-CA	5.14	131.35	121.54

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	67	ARG	Peptide
1	E	68	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	153	0	160	0	0
1	E	136	0	141	1	0
1	H	144	0	150	0	0
All	All	433	0	451	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:64:LEU:HD12	1:E:64:LEU:HA	1.88	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	18/351 (5%)	14 (78%)	4 (22%)	0	100	100
1	E	16/351 (5%)	13 (81%)	2 (12%)	1 (6%)	1	11
1	H	17/351 (5%)	11 (65%)	6 (35%)	0	100	100
All	All	51/1053 (5%)	38 (74%)	12 (24%)	1 (2%)	8	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	68	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	17/285 (6%)	17 (100%)	0	100	100
1	E	15/285 (5%)	15 (100%)	0	100	100
1	H	16/285 (6%)	16 (100%)	0	100	100
All	All	48/855 (6%)	48 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-72430. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 300




Y Index: 300




Z Index: 300


6.2.2 Raw map



X Index: 300



Y Index: 300




Z Index: 300


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 347




Y Index: 253




Z Index: 253


6.3.2 Raw map



X Index: 347



Y Index: 252




Z Index: 253


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.45. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 4648 nm³; this corresponds to an approximate mass of 4199 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.276 $\text{\AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.276 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.62	-	-
Author-provided FSC curve	3.62	4.25	3.70
Unmasked-calculated*	6.09	9.43	6.67

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.09 differs from the reported value 3.62 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-72430 and PDB model 9Y31. Per-residue inclusion information can be found in [section 3](#) on [page 4](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.45 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.45).

9.4 Atom inclusion [i](#)



At the recommended contour level, 94% of all backbone atoms, 89% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.45) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8880	0.5020
C	0.8780	0.4980
E	0.9240	0.5060
H	0.8630	0.5020

