



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:30 AM UTC

PDB ID : 1Y5U / pdb_00001y5u
Title : Dianhydrosugar-based benzamidine, factor Xa specific inhibitor in complex with bovine trypsin mutant
Authors : Di Fenza, A.; Heine, A.; Klebe, G.
Deposited on : 2004-12-03
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	T	223	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	T	2010	-	X	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	PEG	T	501	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 1930 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Trypsin, cationic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	T	223	1646	1025	279	327	15	0	6	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
T	97	GLU	ASN	engineered mutation	UNP P00760
T	99	TYR	LEU	engineered mutation	UNP P00760

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	T	1	Total	Ca	0	0
			1	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	T	1	Total 5	O 4	S 1	0	0
3	T	1	Total 5	O 4	S 1	0	0

- Molecule 4 is 2-O-{3-[AMINO(IMINO)METHYL]PHENYL}-5-O-{4-[AMINO(IMINO)METHYL]PHENYL}-1,4:3,6-DIANHYDRO-D-GLUCITOL (CCD ID: TL4) (formula: $C_{20}H_{22}N_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	T	1	Total	C	N	O	0	0
			28	20	4	4		

- Molecule 5 is IMIDAZOLE (CCD ID: IMD) (formula: $\text{C}_3\text{H}_5\text{N}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	T	1	Total	C	N	0	0
			5	3	2		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	T	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	T	1	Total	C	O	0	0
			7	4	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	T	227	Total	O	0	0
			227	227		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Trypsin, cationic



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	54.79Å 54.79Å 108.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 1.60 47.45 – 1.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-1.60) 97.0 (47.45-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 1.60Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.157 , 0.215 0.167 , 0.213	Depositor DCC
R_{free} test set	2476 reflections (9.96%)	wwPDB-VP
Wilson B-factor (Å ²)	12.9	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 62.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.036 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1930	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GOL, SO4, TL4, IMD, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	T	0.54	0/1702	1.44	15/2308 (0.6%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	T	99	TYR	CA-C-O	7.60	129.29	120.32
1	T	152	PRO	CA-C-N	-6.24	111.13	121.92
1	T	152	PRO	C-N-CA	-6.24	111.13	121.92
1	T	189	ASP	CA-CB-CG	6.04	118.64	112.60
1	T	210	GLN	CA-C-N	-5.80	115.37	121.35
1	T	210	GLN	C-N-CA	-5.80	115.37	121.35
1	T	57	HIS	CA-CB-CG	5.79	119.59	113.80
1	T	40	HIS	CB-CG-CD2	5.74	138.67	131.20
1	T	166	SER	CA-C-O	5.58	126.33	120.42
1	T	40	HIS	ND1-CG-CD2	-5.52	100.58	106.10
1	T	92	PRO	CA-C-N	5.44	131.09	122.60
1	T	92	PRO	C-N-CA	5.44	131.09	122.60
1	T	91	HIS	CA-CB-CG	-5.39	108.41	113.80
1	T	178	SER	CA-C-N	5.28	131.64	122.56
1	T	178	SER	C-N-CA	5.28	131.64	122.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	T	1646	0	1591	21	0
2	T	1	0	0	0	0
3	T	10	0	0	0	0
4	T	28	0	20	3	0
5	T	5	0	5	1	0
6	T	6	0	4	0	0
7	T	7	0	10	4	0
8	T	227	0	0	7	0
All	All	1930	0	1630	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:T:501:PEG:H31	8:T:9210:HOH:O	2.00	0.61
1:T:76:VAL:HG23	7:T:501:PEG:H42	1.83	0.60
1:T:20:TYR:HE2	5:T:12:IMD:H2	1.67	0.58
1:T:192[A]:GLN:HG3	8:T:9185:HOH:O	2.08	0.53
1:T:151:TYR:OH	1:T:192[A]:GLN:HG2	2.11	0.51
1:T:154:VAL:HG22	8:T:9258:HOH:O	2.10	0.51
1:T:31:VAL:HG22	1:T:44:GLY:C	2.38	0.48
1:T:151:TYR:OH	1:T:192[B]:GLN:HG3	2.14	0.47
1:T:192[B]:GLN:HG2	8:T:9145:HOH:O	2.16	0.46
1:T:98:THR:O	4:T:360:TL4:N1	2.49	0.46
1:T:73:ILE:O	7:T:501:PEG:H11	2.16	0.45
1:T:175:GLN:OE1	4:T:360:TL4:N3	2.49	0.44
1:T:135:GLN:OE1	1:T:159:LYS:HD3	2.16	0.44
1:T:187:GLY:HA2	8:T:9119:HOH:O	2.17	0.44
1:T:91:HIS:CG	1:T:92:PRO:HD2	2.53	0.44
1:T:91:HIS:CD2	1:T:92:PRO:HD2	2.53	0.42
1:T:17:VAL:O	1:T:188(A):LYS:HA	2.19	0.42
1:T:175:GLN:HG2	4:T:360:TL4:HN32	1.84	0.42
1:T:27:VAL:HG13	1:T:29:TYR:CZ	2.55	0.42
1:T:130:SER:HB2	8:T:9206:HOH:O	2.19	0.42
7:T:501:PEG:H21	7:T:501:PEG:H41	1.69	0.41
1:T:211:GLY:HA2	1:T:229:THR:O	2.21	0.41
1:T:230[B]:LYS:HE2	8:T:9123:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	T	227/223 (102%)	225 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	T	187/184 (102%)	183 (98%)	4 (2%)	47	23

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	T	32	SER
1	T	146	SER
1	T	147	SER
1	T	170	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	T	50	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	T	74	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	T	2010	-	5,5,5	4.63	5 (100%)	5,5,5	4.52	3 (60%)
5	IMD	T	12	-	5,5,5	0.96	0	5,5,5	0.82	0
3	SO4	T	3022	-	4,4,4	0.30	0	6,6,6	0.29	0
4	TL4	T	360	-	31,31,31	3.74	21 (67%)	35,44,44	2.50	14 (40%)
3	SO4	T	3023	-	4,4,4	0.36	0	6,6,6	0.15	0
7	PEG	T	501	-	6,6,6	0.55	0	5,5,5	2.16	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	T	2010	-	-	2/4/4/4	-
7	PEG	T	501	-	-	3/4/4/4	-
4	TL4	T	360	-	-	1/16/38/38	0/4/4/4
5	IMD	T	12	-	-	-	0/1/1/1

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	T	2010	GOL	C3-C2	-7.57	1.22	1.51
4	T	360	TL4	O8-C9	7.02	1.58	1.44
4	T	360	TL4	C27-C7	6.55	1.51	1.38
4	T	360	TL4	O8-C7	6.12	1.50	1.38
4	T	360	TL4	C23-C19	5.49	1.47	1.39
4	T	360	TL4	C23-C15	5.15	1.47	1.39
4	T	360	TL4	C18-C19	4.81	1.46	1.39
4	T	360	TL4	C16-C15	4.79	1.47	1.38
4	T	360	TL4	C6-C7	4.69	1.47	1.38
4	T	360	TL4	C4-C2	4.63	1.56	1.47
6	T	2010	GOL	O1-C1	4.61	1.61	1.42
4	T	360	TL4	C6-C5	4.44	1.46	1.38
4	T	360	TL4	C17-C16	4.39	1.46	1.38
4	T	360	TL4	C5-C4	4.26	1.45	1.39
4	T	360	TL4	C28-C4	4.17	1.45	1.39
4	T	360	TL4	O14-C13	3.95	1.52	1.44
6	T	2010	GOL	O3-C3	3.70	1.58	1.42
4	T	360	TL4	O14-C15	3.44	1.44	1.38
4	T	360	TL4	C17-C18	3.36	1.44	1.38
4	T	360	TL4	C10-C9	3.31	1.58	1.52
6	T	2010	GOL	C1-C2	-2.95	1.40	1.51
4	T	360	TL4	C24-C13	2.89	1.57	1.52
4	T	360	TL4	C28-C27	2.61	1.43	1.38
6	T	2010	GOL	O2-C2	-2.54	1.36	1.43
4	T	360	TL4	C19-C20	2.36	1.51	1.47
4	T	360	TL4	C12-C13	2.16	1.58	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	T	2010	GOL	O3-C3-C2	7.81	145.54	110.38
4	T	360	TL4	C28-C27-C7	5.67	126.20	119.73
6	T	2010	GOL	O2-C2-C3	5.52	132.03	109.18
4	T	360	TL4	C24-C13-C12	-4.99	93.94	102.40
4	T	360	TL4	C27-C7-C6	-4.95	112.94	120.16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	T	360	TL4	C15-O14-C13	-4.90	110.91	118.93
4	T	360	TL4	O14-C13-C24	4.18	120.29	109.17
4	T	360	TL4	C10-O11-C12	-3.91	100.03	107.84
7	T	501	PEG	O2-C2-C1	3.83	126.98	110.11
4	T	360	TL4	C7-O8-C9	3.56	124.75	118.93
4	T	360	TL4	C4-C2-N3	3.44	123.23	118.01
6	T	2010	GOL	O1-C1-C2	3.03	124.02	110.38
4	T	360	TL4	O25-C26-C9	2.94	120.18	111.65
4	T	360	TL4	O8-C9-C10	-2.89	101.48	109.17
4	T	360	TL4	O8-C7-C27	2.86	133.93	119.90
7	T	501	PEG	C3-O2-C2	2.68	124.97	113.26
4	T	360	TL4	C27-C28-C4	-2.55	118.08	120.80
4	T	360	TL4	C10-C9-C26	2.22	106.17	102.40
4	T	360	TL4	C12-C26-C9	-2.13	98.84	103.78

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	T	501	PEG	C4-C3-O2-C2
7	T	501	PEG	O1-C1-C2-O2
6	T	2010	GOL	O1-C1-C2-O2
7	T	501	PEG	O2-C3-C4-O4
6	T	2010	GOL	O2-C2-C3-O3
4	T	360	TL4	C23-C15-O14-C13


There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	T	12	IMD	1	0
4	T	360	TL4	3	0
7	T	501	PEG	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	T	223/223 (100%)	-0.09	2 (0%) 81 84	8, 14, 28, 38	6 (2%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	T	186	GLU	2.2
1	T	147	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	PEG	T	501	7/7	0.74	0.23	37,40,57,70	0
6	GOL	T	2010	6/6	0.80	0.17	21,40,41,61	0
5	IMD	T	12	5/5	0.81	0.16	37,39,41,44	0
3	SO4	T	3023	5/5	0.89	0.12	36,39,58,79	0
4	TL4	T	360	28/28	0.91	0.10	11,26,33,51	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	T	3022	5/5	0.95	0.10	20,24,30,54	0
2	CA	T	9	1/1	0.99	0.04	12,12,12,12	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.