



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 01:28 AM UTC

PDB ID : 4Y7M / pdb_00004y7m
Title : T6SS protein TssM C-terminal domain (835-1129) from EAEC
Authors : Nguyen, V.S.; Spinelli, S.; Durand, E.; Roussel, A.; Cambillau, C.
Deposited on : 2015-02-15
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	129	5% 94% 6%
1	B	129	3% 95% ..
2	C	295	2% 73% 8% 18%
2	D	295	16% 64% 12% 23%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6337 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hi113 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	129	Total	C	N	O	S	0	0	0
			959	604	158	190	7			
1	B	129	Total	C	N	O	S	0	0	0
			963	607	159	190	7			

- Molecule 2 is a protein called Type VI secretion protein IcmF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	241	Total	C	N	O	S	0	0	0
			1856	1185	326	341	4			
2	D	226	Total	C	N	O	S	0	0	0
			1740	1111	311	314	4			

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	164	Total	O	0	0
			164	164		
4	B	163	Total	O	0	0
			163	163		
4	C	300	Total	O	0	0
			300	300		
4	D	177	Total	O	0	0
			177	177		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hi113 protein




- Molecule 1: Hi113 protein




- Molecule 2: Type VI secretion protein IcmF



- Molecule 2: Type VI secretion protein IcmF





4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α , β , γ	95.23Å 95.23Å 172.95Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.61 – 1.92 47.61 – 1.92	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.61-1.92) 100.0 (47.61-1.92)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 1.92Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.183 , 0.210 0.184 , 0.212	Depositor DCC
R_{free} test set	3356 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	33.4	Xtriage
Anisotropy	0.381	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 55.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.047 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6337	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/980	0.99	1/1330 (0.1%)
1	B	0.80	1/984 (0.1%)	1.00	2/1334 (0.1%)
2	C	0.78	0/1903	1.04	2/2600 (0.1%)
2	D	0.75	0/1784	1.16	5/2433 (0.2%)
All	All	0.76	1/5651 (0.0%)	1.06	10/7697 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	116	MET	SD-CE	-6.17	1.64	1.79

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	68	PHE	CA-CB-CG	7.08	120.88	113.80
2	D	880	ARG	N-CA-C	-7.04	101.88	113.50
1	A	68	PHE	CA-CB-CG	6.87	120.67	113.80
2	C	1105	VAL	N-CA-C	-6.67	105.25	111.45
1	B	18	LEU	CD1-CG-CD2	6.39	124.85	110.80
2	D	1055	ARG	N-CA-C	-5.50	100.83	109.25
2	D	878	ARG	N-CA-C	-5.39	106.64	113.43
2	C	1055	ARG	N-CA-C	-5.22	101.26	109.25
2	D	879	TRP	CA-C-N	5.12	128.05	120.17
2	D	879	TRP	C-N-CA	5.12	128.05	120.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	959	0	898	3	0
1	B	963	0	909	2	0
2	C	1856	0	1821	10	0
2	D	1740	0	1706	18	0
3	B	5	0	0	0	0
3	C	10	0	0	0	0
4	A	164	0	0	0	0
4	B	163	0	0	0	0
4	C	300	0	0	1	0
4	D	177	0	0	0	0
All	All	6337	0	5334	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:999:GLN:HE22	2:C:1014:THR:H	1.27	0.79
2:D:999:GLN:HE22	2:D:1014:THR:H	1.26	0.78
1:A:11:LEU:HD11	1:A:128:SER:HB3	1.72	0.70
2:D:1004:VAL:HG12	2:D:1005:ASN:N	2.07	0.69
2:D:980:GLU:OE1	2:D:1083:ARG:HD2	1.98	0.63
2:C:995:ILE:HG12	2:C:1000:LYS:HG2	1.85	0.59
2:D:995:ILE:HG12	2:D:1000:LYS:HG2	1.85	0.58
2:D:1056:ARG:HD2	2:D:1091:LEU:HD22	1.86	0.57
2:D:932:LEU:HD13	2:D:941:PRO:HB3	1.87	0.57
2:C:1056:ARG:HD2	2:C:1091:LEU:HD22	1.88	0.54
2:D:883:VAL:HG23	2:D:884:VAL:H	1.72	0.54
2:D:951:PHE:HB3	2:D:956:LEU:HD11	1.90	0.53
2:C:896:PRO:O	2:C:1097:ARG:HD2	2.09	0.52
2:D:876:ASN:C	2:D:878:ARG:H	2.17	0.52
2:C:1037:GLN:HE21	2:C:1040:ASP:HB2	1.75	0.52
2:D:1004:VAL:CG1	2:D:1005:ASN:N	2.73	0.50
2:D:1054:ALA:HB2	2:D:1070:TRP:CE2	2.48	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:896:PRO:O	2:D:1097:ARG:HD2	2.13	0.48
2:D:884:VAL:HG11	2:D:1103:GLU:HG2	1.95	0.48
2:D:879:TRP:CD1	2:D:1105:VAL:HG23	2.47	0.48
2:D:879:TRP:O	2:D:879:TRP:CG	2.67	0.48
2:D:1004:VAL:HG12	2:D:1005:ASN:CG	2.41	0.45
2:D:980:GLU:OE1	2:D:1083:ARG:CD	2.65	0.44
2:C:1054:ALA:HB2	2:C:1070:TRP:CE2	2.53	0.44
2:C:1071:GLN:HE21	2:C:1077:MET:HE1	1.82	0.43
1:A:12:VAL:HG11	1:A:86:LEU:HD13	2.01	0.43
1:A:20:LEU:HG	1:A:83:MET:HE2	2.02	0.42
2:C:870:PRO:HD2	4:C:1597:HOH:O	2.19	0.42
1:B:78:MET:HE2	1:B:78:MET:HB2	1.90	0.42
2:C:1023:GLY:HA2	2:C:1046:GLY:HA3	2.02	0.42
2:C:879:TRP:HZ2	2:C:1101:LEU:HD22	1.85	0.41
2:D:881:THR:HA	2:D:882:ALA:HA	1.81	0.41
1:B:12:VAL:HG11	1:B:86:LEU:HD13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/129 (98%)	125 (98%)	2 (2%)	0	100	100
1	B	127/129 (98%)	125 (98%)	2 (2%)	0	100	100
2	C	239/295 (81%)	233 (98%)	6 (2%)	0	100	100
2	D	222/295 (75%)	212 (96%)	10 (4%)	0	100	100
All	All	715/848 (84%)	695 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/102 (96%)	97 (99%)	1 (1%)	68	64
1	B	99/102 (97%)	98 (99%)	1 (1%)	68	64
2	C	190/237 (80%)	185 (97%)	5 (3%)	40	24
2	D	176/237 (74%)	168 (96%)	8 (4%)	24	9
All	All	563/678 (83%)	548 (97%)	15 (3%)	39	24

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	MET
1	B	78	MET
2	C	911	LYS
2	C	946	THR
2	C	950	THR
2	C	1092	VAL
2	C	1101	LEU
2	D	878	ARG
2	D	911	LYS
2	D	924	LEU
2	D	927	ASN
2	D	963	SER
2	D	1029	VAL
2	D	1092	VAL
2	D	1101	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	82	GLN
1	A	84	ASN
1	B	77	ASN
2	C	999	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	C	1037	GLN
2	C	1071	GLN
2	D	927	ASN
2	D	999	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	C	1202	-	4,4,4	0.23	0	6,6,6	0.10	0
3	SO4	B	201	-	4,4,4	0.38	0	6,6,6	0.22	0
3	SO4	C	1201	-	4,4,4	0.37	0	6,6,6	0.37	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	129/129 (100%)	0.15	6 (4%) 36 41	26, 37, 55, 72	0
1	B	129/129 (100%)	0.02	4 (3%) 51 56	27, 33, 51, 75	0
2	C	241/295 (81%)	0.04	7 (2%) 53 60	26, 34, 58, 75	0
2	D	226/295 (76%)	1.04	48 (21%) 2 3	27, 45, 100, 126	0
All	All	725/848 (85%)	0.37	65 (8%) 15 18	26, 36, 81, 126	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	879	TRP	6.9
2	D	951	PHE	6.3
2	D	882	ALA	6.3
2	D	955	PHE	5.6
2	D	1105	VAL	5.2
2	D	932	LEU	5.2
1	A	42	GLY	5.1
2	C	949	LEU	4.9
2	D	872	ALA	4.5
2	D	1101	LEU	4.4
2	D	887	TRP	4.2
2	D	877	ALA	4.1
2	D	928	LEU	4.1
2	D	1102	PRO	4.0
2	D	881	THR	3.9
2	D	923	PHE	3.9
2	D	884	VAL	3.9
1	B	42	GLY	3.7
1	B	41	PRO	3.7
2	C	1108	LEU	3.6
2	D	940	VAL	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	1035	THR	3.5
2	D	883	VAL	3.5
2	D	959	ILE	3.4
2	D	931	VAL	3.4
2	D	924	LEU	3.4
2	D	880	ARG	3.4
2	C	943	THR	3.4
2	D	1100	VAL	3.3
2	D	885	ASP	3.2
2	D	956	LEU	3.2
2	D	1034	GLY	3.2
1	A	11	LEU	3.1
2	C	948	GLY	3.1
2	C	868	LEU	3.1
1	B	129	SER	3.1
2	D	891	PHE	3.0
2	D	875	LEU	2.9
2	D	892	SER	2.9
2	D	942	ASP	2.9
2	D	952	ASN	2.7
1	A	43	LYS	2.7
2	D	893	GLY	2.7
2	D	954	ALA	2.6
2	D	1103	GLU	2.6
2	D	890	ALA	2.6
1	A	129	SER	2.6
2	D	939	TRP	2.6
2	D	886	GLY	2.6
2	D	1104	THR	2.6
2	D	1075	GLY	2.5
2	D	927	ASN	2.4
2	D	941	PRO	2.4
2	D	957	LYS	2.4
2	C	944	ILE	2.3
2	C	946	THR	2.3
2	D	1033	ALA	2.2
1	B	76	LYS	2.2
2	D	873	GLU	2.2
2	D	888	ASN	2.2
2	D	889	ASN	2.2
2	D	953	PRO	2.2
2	D	874	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	88	PRO	2.1
1	A	14	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	C	1202	5/5	0.71	0.13	103,104,104,106	0
3	SO4	B	201	5/5	0.81	0.14	105,105,106,107	0
3	SO4	C	1201	5/5	0.89	0.15	93,93,94,94	0

6.5 Other polymers [i](#)

There are no such residues in this entry.