



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 04:30 AM UTC

PDB ID : 4Y7P / pdb_00004y7p
Title : Structure of alkaline D-peptidase from *Bacillus cereus*
Authors : Nakano, S.; Okazaki, S.; Ishitsubo, E.; Kawahara, N.; Komeda, H.; Tokiwa, H.; Asano, Y.
Deposited on : 2015-02-15
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2865 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alkaline D-peptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	5	0
			2653	1681	448	517	7			

- Molecule 2 is THIOCYANATE ION (CCD ID: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	S	0	0
			3	1	1	1		
2	A	1	Total	C	N	S	0	0
			3	1	1	1		
2	A	1	Total	C	N	S	0	0
			3	1	1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	203	Total 203	O 203	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alkaline D-peptidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.02Å 104.02Å 98.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.22 – 2.10 24.22 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (24.22-2.10) 99.8 (24.22-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.37 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.174 , 0.219 0.183 , 0.224	Depositor DCC
R_{free} test set	1813 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	30.4	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2865	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SCN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.52	21/2706 (0.8%)	1.28	14/3660 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	201	GLU	CD-OE1	10.20	1.44	1.25
1	A	224	ARG	CD-NE	-7.95	1.35	1.46
1	A	185	THR	C-O	-7.02	1.15	1.24
1	A	298	ILE	CA-CB	-6.49	1.48	1.55
1	A	40	ALA	CA-C	-6.25	1.44	1.52
1	A	202	PRO	C-O	-6.25	1.17	1.24
1	A	84	VAL	CA-CB	-5.94	1.47	1.54
1	A	42	THR	CA-C	-5.91	1.45	1.52
1	A	350	ASN	CA-C	-5.88	1.44	1.52
1	A	158	SER	C-O	-5.80	1.16	1.24
1	A	327	LEU	CA-C	-5.58	1.46	1.52
1	A	21	ASN	CA-C	-5.41	1.45	1.52
1	A	326	THR	C-O	5.40	1.30	1.23
1	A	66	THR	C-O	-5.36	1.17	1.24
1	A	116	GLN	CA-C	5.33	1.59	1.52
1	A	274	GLU	N-CA	5.24	1.52	1.46
1	A	122	LEU	C-O	-5.18	1.18	1.24
1	A	282	THR	CA-C	-5.05	1.46	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	146	SER	C-O	-5.04	1.17	1.24
1	A	227	ILE	N-CA	-5.02	1.40	1.46
1	A	124	ASN	N-CA	5.01	1.52	1.46

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	345	PRO	CA-C-N	-8.19	113.72	122.89
1	A	345	PRO	C-N-CA	-8.19	113.72	122.89
1	A	109	GLY	N-CA-C	-8.00	103.33	112.29
1	A	224	ARG	CB-CG-CD	-6.91	95.40	111.30
1	A	134	ARG	CG-CD-NE	-6.61	97.45	112.00
1	A	293	ARG	N-CA-C	6.42	120.09	109.88
1	A	201	GLU	CG-CD-OE2	-6.05	104.49	118.40
1	A	346	ASP	CA-C-N	-5.90	113.67	120.04
1	A	346	ASP	C-N-CA	-5.90	113.67	120.04
1	A	224	ARG	NE-CZ-NH1	5.70	127.20	121.50
1	A	224	ARG	NE-CZ-NH2	-5.58	114.18	119.20
1	A	328	GLY	N-CA-C	-5.45	108.20	114.69
1	A	122	LEU	N-CA-C	-5.35	105.44	111.28
1	A	350	ASN	N-CA-C	5.14	117.78	111.82

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	229	LEU	Peptide
1	A	287	GLY	Peptide
1	A	292[B]	GLY	Peptide
1	A	357	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2653	0	2644	31	0
2	A	9	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	203	0	0	1	0
All	All	2865	0	2644	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:289[B]:ALA:H	1:A:291[B]:LEU:HD22	1.38	0.87
1:A:289[B]:ALA:N	1:A:291[B]:LEU:HD22	1.93	0.83
1:A:291[B]:LEU:H	1:A:291[B]:LEU:HD13	1.54	0.73
1:A:136:LYS:HA	1:A:136:LYS:HE3	1.77	0.67
1:A:286:THR:H	1:A:292[B]:GLY:HA2	1.62	0.65
1:A:124:ASN:C	1:A:124:ASN:HD22	2.08	0.62
1:A:326:THR:CG2	3:A:585:HOH:O	2.50	0.59
1:A:242:ASN:HD22	1:A:244:SER:H	1.50	0.58
1:A:287:GLY:O	1:A:288[A]:GLU:HG2	2.05	0.57
1:A:230:ASP:HB3	1:A:233:SER:OG	2.09	0.52
1:A:230:ASP:O	1:A:343:GLU:OE1	2.28	0.51
1:A:242:ASN:HD22	1:A:242:ASN:C	2.18	0.51
1:A:289[B]:ALA:N	1:A:291[B]:LEU:CD2	2.71	0.49
1:A:288[B]:GLU:HB2	1:A:291[B]:LEU:HD23	1.95	0.48
1:A:340:LEU:C	1:A:340:LEU:HD23	2.39	0.47
1:A:145:LYS:HD3	1:A:147:TYR:CZ	2.50	0.47
1:A:291[B]:LEU:HD13	1:A:291[B]:LEU:N	2.20	0.46
1:A:148:ARG:HH11	1:A:148:ARG:HG2	1.80	0.46
1:A:290[B]:ALA:O	1:A:291[B]:LEU:C	2.59	0.46
1:A:291[A]:LEU:N	1:A:291[A]:LEU:CD1	2.78	0.45
1:A:316:ILE:O	1:A:317:PRO:C	2.58	0.45
1:A:224:ARG:HD2	1:A:237:ASP:OD2	2.17	0.45
1:A:303:LEU:HD11	1:A:309:ILE:HD11	1.98	0.44
1:A:288[B]:GLU:H	1:A:291[B]:LEU:CD2	2.30	0.44
1:A:290[B]:ALA:O	1:A:292[B]:GLY:N	2.51	0.43
1:A:230:ASP:CB	1:A:233:SER:OG	2.66	0.43
1:A:31:LEU:HD21	1:A:38:ILE:HG22	1.99	0.43
1:A:242:ASN:ND2	1:A:244:SER:H	2.14	0.43
1:A:288[A]:GLU:C	1:A:290[A]:ALA:N	2.73	0.42
1:A:62:LYS:HA	1:A:63:PRO:HD3	1.76	0.42
1:A:224:ARG:CD	1:A:237:ASP:OD2	2.67	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	345/388 (89%)	327 (95%)	15 (4%)	3 (1%)	14 10

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	291[A]	LEU
1	A	291[B]	LEU
1	A	250	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	287/326 (88%)	268 (93%)	19 (7%)	15 13

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	ARG
1	A	20	ARG
1	A	49	TRP
1	A	59	SER
1	A	115	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	124	ASN
1	A	136	LYS
1	A	145	LYS
1	A	178	LEU
1	A	230	ASP
1	A	242	ASN
1	A	267	LEU
1	A	291[A]	LEU
1	A	291[B]	LEU
1	A	326	THR
1	A	327	LEU
1	A	330	LYS
1	A	349	LYS
1	A	358	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	124	ASN
1	A	189	ASN
1	A	242	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SCN	A	402	-	1,2,2	0.40	0	0,1,1	-	-
2	SCN	A	401	-	1,2,2	0.48	0	0,1,1	-	-
2	SCN	A	403	-	1,2,2	1.56	0	0,1,1	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	342/388 (88%)	-0.01	13 (3%)	44 46	18, 30, 60, 102	5 (1%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	291[A]	LEU	11.1
1	A	289[A]	ALA	10.4
1	A	290[A]	ALA	7.3
1	A	288[A]	GLU	6.4
1	A	292[A]	GLY	6.0
1	A	49	TRP	4.6
1	A	231	GLY	4.0
1	A	232	ALA	3.9
1	A	229	LEU	3.9
1	A	358	LYS	3.3
1	A	357	SER	2.6
1	A	327	LEU	2.3
1	A	47	LYS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SCN	A	401	3/3	0.96	0.18	36,36,38,43	0
2	SCN	A	403	3/3	0.97	0.06	27,27,31,36	0
2	SCN	A	402	3/3	0.99	0.07	28,28,33,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.