



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 27, 2026 – 05:51 PM EDT

PDB ID : 5Y7O / pdb_00005y7o
Title : Crystal structure of folding sensor region of UGGT from *Thermomyces dubon-
tii*
Authors : Satoh, T.; Song, C.; Zhu, T.; Toshimori, T.; Murata, K.; Hayashi, Y.;
Kamikubo, H.; Uchihashi, T.; Kato, K.
Deposited on : 2017-08-17
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 14318 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UGGT.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	913	Total	C	N	O	S	Se	0	0	0
			7159	4556	1224	1362	2	15			
1	B	913	Total	C	N	O	S	Se	0	0	0
			7159	4556	1224	1362	2	15			

SEQUENCE-PLOTS INFOmissingINFO

3 Data and refinement statistics

Property	Value	Source
Space group	P 32 1 2	Depositor
Cell constants a, b, c, α , β , γ	195.09Å 195.09Å 142.26Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.97 – 3.10 19.97 – 3.10	Depositor EDS
% Data completeness (in resolution range)	98.0 (19.97-3.10) 98.0 (19.97-3.10)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 3.09Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.232 , 0.278 0.232 , 0.288	Depositor DCC
R_{free} test set	2744 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	110.4	Xtriage
Anisotropy	0.256	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 162.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.55$, $\langle L^2 \rangle = 0.40$	Xtriage
Estimated twinning fraction	0.478 for -h,-k,l	Xtriage
Reported twinning fraction	0.500 for -h,-k,l	Depositor
Outliers	0 of 54601 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14318	wwPDB-VP
Average B, all atoms (Å ²)	123.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality

4.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	3/7286 (0.0%)	1.51	117/9871 (1.2%)
1	B	0.82	4/7286 (0.1%)	1.44	103/9871 (1.0%)
All	All	0.84	7/14572 (0.0%)	1.47	220/19742 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	14
1	B	0	9
All	All	0	23

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	209	ALA	CA-CB	-9.38	1.37	1.53
1	A	100	ALA	CA-CB	-6.32	1.45	1.54
1	A	292	VAL	CA-CB	-6.11	1.48	1.55
1	B	292	VAL	CA-CB	-5.94	1.48	1.55
1	B	855	VAL	CA-CB	5.52	1.59	1.53

The worst 5 of 220 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	664	ASP	CA-C-N	-15.82	100.06	119.84
1	B	664	ASP	C-N-CA	-15.82	100.06	119.84
1	A	804	ARG	N-CA-C	12.97	128.78	109.07
1	A	854	LYS	N-CA-C	-11.74	98.46	113.72
1	A	664	ASP	CA-C-N	-11.31	105.70	119.84

There are no chirality outliers.

5 of 23 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	150	MSE	Peptide
1	A	151	GLN	Peptide
1	A	204	LYS	Peptide
1	A	391	LEU	Peptide
1	A	456	TYR	Peptide

4.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7159	0	7156	704	0
1	B	7159	0	7156	600	1
All	All	14318	0	14312	1297	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 45.

The worst 5 of 1297 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:703:VAL:HG22	1:A:804:ARG:HH11	1.15	1.10
1:A:380:ARG:HH11	1:A:912:ILE:HD12	1.02	1.07
1:B:210:LYS:HB2	1:B:211:GLU:HG2	1.11	1.06
1:A:433:ASN:HD22	1:A:500:ARG:HA	1.21	1.04
1:B:420:ARG:HE	1:B:649:LEU:HD21	1.22	1.00

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:456:TYR:OH	1:B:456:TYR:OH[5_554]	2.09	0.11

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	893/1130 (79%)	685 (77%)	155 (17%)	53 (6%)	1	8
1	B	893/1130 (79%)	696 (78%)	155 (17%)	42 (5%)	2	11
All	All	1786/2260 (79%)	1381 (77%)	310 (17%)	95 (5%)	1	10

5 of 95 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	56	GLU
1	A	61	TYR
1	A	148	ASP
1	A	348	LEU
1	A	524	MSE

4.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	782/948 (82%)	713 (91%)	69 (9%)	9	33
1	B	782/948 (82%)	722 (92%)	60 (8%)	12	38
All	All	1564/1896 (82%)	1435 (92%)	129 (8%)	10	36

5 of 129 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	824	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	866	SER
1	A	763	LEU
1	A	762	ILE
1	B	914	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	616	GLN
1	B	896	ASN
1	B	1037	HIS
1	B	913	ASN
1	B	635	GLN

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

4.6 Ligand geometry [i](#)

There are no ligands in this entry.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	898/1130 (79%)	-1.13	0	100	100	70, 117, 187, 345	0
1	B	898/1130 (79%)	-1.11	1 (0%)	92	86	76, 115, 181, 279	0
All	All	1796/2260 (79%)	-1.12	1 (0%)	92	86	70, 116, 185, 345	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	94	GLY	3.2

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.4 Ligands [i](#)

There are no ligands in this entry.

5.5 Other polymers [i](#)

There are no such residues in this entry.