



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 06:05 AM UTC

PDB ID : 7Y7P / pdb_00007y7p
Title : QDE-1 in complex with RNA template, RNA primer and AMPNPP
Authors : Cui, R.X.; Gan, J.H.; Ma, J.B.
Deposited on : 2022-06-22
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)
RNA backbone	3983	1044 (2.90-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1026	75% 15% 9%
1	B	1026	73% 15% 12%
2	C	14	21% 43% 14% 21%
2	E	14	7% 29% 50% 21%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	7	86% 14%
3	J	7	29% 43% 14% 14%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 15042 atoms, of which 18 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-dependent RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	934	Total	C	N	O	S	0	0	0
			7155	4577	1226	1320	32			
1	B	908	Total	C	N	O	S	0	0	0
			6905	4410	1187	1275	33			

- Molecule 2 is a RNA chain called RNA (5'-R(*GP*AP*A*CP*UP*AP*UP*GP*GP*UP*C P*GP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	11	Total	C	N	O	P	0	0	0
			231	104	42	75	10			
2	E	11	Total	C	N	O	P	0	0	0
			231	104	42	75	10			

- Molecule 3 is a RNA chain called RNA (5'-R(*UP*CP*CP*GP*AP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	7	Total	C	N	O	P	0	0	0
			142	65	24	47	6			
3	J	6	Total	C	N	O	P	0	0	0
			125	56	22	41	6			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Ca	0	0
			3	3		
4	B	3	Total	Ca	0	0
			3	3		
4	C	1	Total	Ca	0	0
			1	1		


- # ZAN
-
- The chemical structure of ZAN (Zinc Nucleoside Analog) is shown. It consists of a pyrimidine base (labeled N1, N3, N7, N9, C2, C4, C6, C8) attached to a ribose sugar (labeled C1', C2', C3', C4', C5') via a glycosidic bond. The ribose sugar is further attached to a phosphate group (labeled O1, O2, O3, O4, O5, O6, O7, O8, O9, O10, O11, O12, O13, O14, O15, O16, O17, O18, O19, O20, O21, O22, O23, O24, O25, O26, O27, O28, O29, O30, O31, O32, O33, O34, O35, O36, O37, O38, O39, O40, O41, O42, O43, O44, O45, O46, O47, O48, O49, O50, O51, O52, O53, O54, O55, O56, O57, O58, O59, O60, O61, O62, O63, O64, O65, O66, O67, O68, O69, O70, O71, O72, O73, O74, O75, O76, O77, O78, O79, O80, O81, O82, O83, O84, O85, O86, O87, O88, O89, O90, O91, O92, O93, O94, O95, O96, O97, O98, O99, O100, O101, O102, O103, O104, O105, O106, O107, O108, O109, O110, O111, O112, O113, O114, O115, O116, O117, O118, O119, O120, O121, O122, O123, O124, O125, O126, O127, O128, O129, O130, O131, O132, O133, O134, O135, O136, O137, O138, O139, O140, O141, O142, O143, O144, O145, O146, O147, O148, O149, O150, O151, O152, O153, O154, O155, O156, O157, O158, O159, O160, O161, O162, O163, O164, O165, O166, O167, O168, O169, O170, O171, O172, O173, O174, O175, O176, O177, O178, O179, O180, O181, O182, O183, O184, O185, O186, O187, O188, O189, O190, O191, O192, O193, O194, O195, O196, O197, O198, O199, O200, O201, O202, O203, O204, O205, O206, O207, O208, O209, O210, O211, O212, O213, O214, O215, O216, O217, O218, O219, O220, O221, O222, O223, O224, O225, O226, O227, O228, O229, O230, O231, O232, O233, O234, O235, O236, O237, O238, O239, O240, O241, O242, O243, O244, O245, O246, O247, O248, O249, O250, O251, O252, O253, O254, O255, O256, O257, O258, O259, O260, O261, O262, O263, O264, O265, O266, O267, O268, O269, O270, O271, O272, O273, O274, O275, O276, O277, O278, O279, O280, O281, O282, O283, O284, O285, O286, O287, O288, O289, O290, O291, O292, O293, O294, O295, O296, O297, O298, O299, O300, O301, O302, O303, O304, O305, O306, O307, O308, O309, O310, O311, O312, O313, O314, O315, O316, O317, O318, O319, O320, O321, O322, O323, O324, O325, O326, O327, O328, O329, O330, O331, O332, O333, O334, O335, O336, O337, O338, O339, O340, O341, O342, O343, O344, O345, O346, O347, O348, O349, O350, O351, O352, O353, O354, O355, O356, O357, O358, O359, O360, O361, O362, O363, O364, O365, O366, O367, O368, O369, O370, O371, O372, O373, O374, O375, O376, O377, O378, O379, O380, O381, O382, O383, O384, O385, O386, O387, O388, O389, O390, O391, O392, O393, O394, O395, O396, O397, O398, O399, O400, O401, O402, O403, O404, O405, O406, O407, O408, O409, O410, O411, O412, O413, O414, O415, O416, O417, O418, O419, O420, O421, O422, O423, O424, O425, O426, O427, O428, O429, O430, O431, O432, O433, O434, O435, O436, O437, O438, O439, O440, O441, O442, O443, O444, O445, O446, O447, O448, O449, O450, O451, O452, O453, O454, O455, O456, O457, O458, O459, O460, O461, O462, O463, O464, O465, O466, O467, O468, O469, O470, O471, O472, O473, O474, O475, O476, O477, O478, O479, O480, O481, O482, O483, O484, O485, O486, O487, O488, O489, O490, O491, O492, O493, O494, O495, O496, O497, O498, O499, O500, O501, O502, O503, O504, O505, O506, O507, O508, O509, O510, O511, O512, O513, O514, O515, O516, O517, O518, O519, O520, O521, O522, O523, O524, O525, O526, O527, O528, O529, O530, O531, O532, O533, O534, O535, O536, O537, O538, O539, O540, O541, O542, O543, O544, O545, O546, O547, O548, O549, O550, O551, O552, O553, O554, O555, O556, O557, O558, O559, O560, O561, O562, O563, O564, O565, O566, O567, O568, O569, O570, O571, O572, O573, O574, O575, O576, O577, O578, O579, O580, O581, O582, O583, O584, O585, O586, O587, O588, O589, O590, O591, O592, O593, O594, O595, O596, O597, O598, O599, O600, O601, O602, O603, O604, O605, O606, O607, O608, O609, O610, O611, O612, O613, O614, O615, O616, O617, O618, O619, O620, O621, O622, O623, O624, O625, O626, O627, O628, O629, O630, O631, O632, O633, O634, O635, O636, O637, O638, O639, O640, O641, O642, O643, O644, O645, O646, O647, O648, O649, O650, O651, O652, O653, O654, O655, O656, O657, O658, O659, O660, O661, O662, O663, O664, O665, O666, O667, O668, O669, O670, O671, O672, O673, O674, O675, O676, O677, O678, O679, O680, O681, O682, O683, O684, O685, O686, O687, O688, O689, O690, O691, O692, O693, O694, O695, O696, O697, O698, O699, O700, O701, O702, O703, O704, O705, O706, O707, O708, O709, O710, O711, O712, O713, O714, O715, O716, O717, O718, O719, O720, O721, O722, O723, O724, O725, O726, O727, O728, O729, O730, O731, O732, O733, O734, O735, O736, O737, O738, O739, O740, O741, O742, O743, O744, O745, O746, O747, O748, O749, O750, O751, O752, O753, O754, O755, O756, O757, O758, O759, O760, O761, O762, O763, O764, O765, O766, O767, O768, O769, O770, O771, O772, O773, O774, O775, O776, O777, O778, O779, O780, O781, O782, O783, O784, O785, O786, O787, O788, O789, O790, O791, O792, O793, O794, O795, O796, O797, O798, O799, O800, O801, O802, O803, O804, O805, O806, O807, O808, O809, O810, O811, O81

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
5	B	1	Total 31	C 10	N 6	O 12	P 3	0	0

- PEG
-
- The diagram illustrates the chemical structure of PEG (polyethylene glycol). It shows a zigzag chain of carbon atoms (C1, C2, C3, C4) and oxygen atoms (O1, O2). The chain is terminated by hydroxyl groups (HO and OH). The atoms are labeled with green text, and the bonds are shown as black lines. The hydroxyl groups are shown in red.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	70	Total	O	0	0
			70	70		
8	B	67	Total	O	0	0
			67	67		
8	C	6	Total	O	0	0
			6	6		
8	D	2	Total	O	0	0
			2	2		
8	E	6	Total	O	0	0
			6	6		
8	J	2	Total	O	0	0
			2	2		


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

[illegible]



Chain B:

2% 73% 15% 12%

ALA ARG SER LEU GLU THR ALA ARG SER VAL GLN VAL HIS PRO VAL V394 F405 P406 K407 W408 L414 A415 V416 E432 ASP GLU SER LEU GLY L438 V450 I453 L460 ASP ALA PHE ARG GLY LYS PRO PHE PRO E470 K471 P472 P473 A474 D475 V493 M496 V497 L498 S505 P506 THR A508 V513 M519 Q522 G523 C524 I538 S542 P543 S547 P548 S549 V550 S555 K556 Q557 P558 A560 V564 M570 L575 V576 F583 A588 G589 Y590 R596 GLU PHE GLN LEU LEU ARG ALA GLU ASP PRO LYS P606 I607 P608




- Molecule 2: RNA (5'-R(*GP*AP*A*CP*UP*AP*UP*GP*GP*UP*CP*GP*GP*A)-3')



- Molecule 2: RNA (5'-R(*GP*AP*A*CP*UP*AP*UP*GP*GP*UP*CP*GP*GP*A)-3')



- Molecule 3: RNA (5'-R(*UP*CP*CP*GP*AP*CP*C)-3')



- Molecule 3: RNA (5'-R(*UP*CP*CP*GP*AP*CP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	101.95Å 120.20Å 114.86Å 90.00° 109.62° 90.00°	Depositor
Resolution (Å)	29.62 – 2.70 29.62 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.0 (29.62-2.70) 97.9 (29.62-2.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.01 (at 2.72Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.223 , 0.254 0.224 , 0.255	Depositor DCC
R_{free} test set	3462 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	56.3	Xtriage
Anisotropy	0.384	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 57.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	15042	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.28 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.6473e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, ZAN, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.70	1/7338 (0.0%)	1.02	5/9983 (0.1%)
1	B	0.76	0/7080	1.10	6/9641 (0.1%)
2	C	0.47	0/258	0.70	0/401
2	E	0.09	0/258	0.30	0/401
3	D	0.08	0/157	0.17	0/242
3	J	0.11	0/138	0.33	0/212
All	All	0.71	1/15229 (0.0%)	1.04	11/20880 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	930	LEU	C-N	-6.18	1.25	1.33

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1099	VAL	N-CA-C	-7.01	105.51	112.17
1	A	995	GLY	CA-C-O	-6.01	118.31	122.22
1	B	1007	ASP	CA-CB-CG	5.99	118.59	112.60
1	B	995	GLY	CA-C-O	-5.80	118.45	122.22
1	B	1099	VAL	N-CA-C	-5.80	107.33	112.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7155	0	6706	97	0
1	B	6905	0	6419	96	0
2	C	231	0	116	3	0
2	E	231	0	116	2	0
3	D	142	0	77	1	0
3	J	125	0	66	2	0
4	A	3	0	0	0	0
4	B	3	0	0	1	0
4	C	1	0	0	0	0
5	A	31	0	17	1	0
5	B	31	0	16	0	0
6	B	7	10	10	0	0
7	B	6	8	8	0	0
8	A	70	0	0	3	0
8	B	67	0	0	1	0
8	C	6	0	0	0	0
8	D	2	0	0	0	0
8	E	6	0	0	0	0
8	J	2	0	0	0	0
All	All	15024	18	13551	194	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 194 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1088:TYR:CD2	1:A:1088:TYR:CZ	2.37	1.06
1:B:460:LEU:C	1:B:460:LEU:HD13	2.04	0.82
1:A:1329:LEU:HA	1:A:1332:ILE:HD12	1.66	0.78
1:A:1227:LEU:HD11	1:A:1305:ARG:HG3	1.66	0.76
1:B:922:MET:HE3	1:B:1011:ASP:HB3	1.67	0.76

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	916/1026 (89%)	899 (98%)	17 (2%)	0	100	100
1	B	888/1026 (86%)	868 (98%)	20 (2%)	0	100	100
All	All	1804/2052 (88%)	1767 (98%)	37 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	721/894 (81%)	693 (96%)	28 (4%)	28	57
1	B	689/894 (77%)	658 (96%)	31 (4%)	24	52
All	All	1410/1788 (79%)	1351 (96%)	59 (4%)	26	55

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	450	VAL
1	B	1285	ILE
1	B	821	ILE
1	B	1269	THR
1	B	1219	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1097	ASN
1	A	1183	HIS
1	B	613	HIS
1	A	1184	ASN
1	A	827	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	10/14 (71%)	4 (40%)	2 (20%)
2	E	10/14 (71%)	3 (30%)	1 (10%)
3	D	6/7 (85%)	0	0
3	J	5/7 (71%)	2 (40%)	0
All	All	31/42 (73%)	9 (29%)	3 (9%)

5 of 9 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	0	A
2	C	6	G
2	C	7	G
2	C	8	A
2	E	0	A

All (3) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	C	-2	C
2	C	7	G
2	E	-2	C

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 11 ligands modelled in this entry, 7 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ZAN	A	1504	4	32,33,33	1.69	8 (25%)	46,52,52	2.01	10 (21%)
6	PEG	B	1501	-	6,6,6	0.13	0	5,5,5	0.09	0
7	GOL	B	1502	-	5,5,5	0.07	0	5,5,5	0.24	0
5	ZAN	B	1506	4	32,33,33	1.01	4 (12%)	46,52,52	1.13	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ZAN	A	1504	4	-	5/19/38/38	0/3/3/3
6	PEG	B	1501	-	-	2/4/4/4	-
7	GOL	B	1502	-	-	2/4/4/4	-
5	ZAN	B	1506	4	-	6/19/38/38	0/3/3/3

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1504	ZAN	C5-C4	4.73	1.47	1.39
5	A	1504	ZAN	PB-O2B	3.40	1.51	1.46
5	A	1504	ZAN	PA-O2A	3.37	1.51	1.46
5	B	1506	ZAN	PB-O2B	2.88	1.50	1.46
5	B	1506	ZAN	PA-O2A	2.69	1.50	1.46

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1504	ZAN	C5-C4-N3	-5.87	118.64	126.72
5	A	1504	ZAN	N3-C4-N9	4.72	135.20	127.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1504	ZAN	O1A-PA-O2A	4.58	119.68	109.87
5	A	1504	ZAN	O1B-PB-O2B	4.30	119.09	109.87
5	B	1506	ZAN	O1B-PB-O2B	4.21	118.90	109.87

There are no chirality outliers.

5 of 15 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	1504	ZAN	PB-N3A-PA-O2A
5	A	1504	ZAN	C5'-O5'-PA-O1A
5	A	1504	ZAN	C3'-C4'-C5'-O5'
5	B	1506	ZAN	PB-N3A-PA-O2A
5	B	1506	ZAN	C5'-O5'-PA-O1A

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1504	ZAN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	934/1026 (91%)	0.16	13 (1%) 73 72	37, 76, 114, 152	0
1	B	908/1026 (88%)	0.23	21 (2%) 61 58	39, 77, 122, 156	0
2	C	11/14 (78%)	-0.15	0 100 100	58, 71, 163, 164	0
2	E	11/14 (78%)	0.20	1 (9%) 15 12	64, 78, 169, 172	0
3	D	7/7 (100%)	-0.38	0 100 100	50, 72, 124, 138	0
3	J	6/7 (85%)	0.03	0 100 100	54, 85, 132, 142	0
All	All	1877/2094 (89%)	0.19	35 (1%) 66 63	37, 76, 119, 172	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	625	ASP	4.8
1	B	460	LEU	4.4
1	B	1137	LEU	4.0
1	B	849	TYR	3.7
1	B	557	GLN	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	GOL	B	1502	6/6	0.84	0.12	63,80,110,110	0
4	CA	B	1504	1/1	0.91	0.08	79,79,79,79	0
4	CA	C	101	1/1	0.92	0.08	122,122,122,122	0
6	PEG	B	1501	7/7	0.93	0.12	63,76,88,88	0
5	ZAN	B	1506	31/31	0.97	0.06	41,55,66,81	0
4	CA	A	1502	1/1	0.97	0.04	85,85,85,85	0
5	ZAN	A	1504	31/31	0.97	0.06	41,52,64,76	0
4	CA	A	1501	1/1	0.98	0.03	42,42,42,42	0
4	CA	B	1505	1/1	0.99	0.02	51,51,51,51	0
4	CA	A	1503	1/1	0.99	0.04	46,46,46,46	0
4	CA	B	1503	1/1	1.00	0.02	44,44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.