



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:19 PM UTC

PDB ID : 5Y82 / pdb_00005y82
Title : Crystal structure of the periplasmic domain of the Thermotoga maritima YidC
Authors : Huang, Y.; Xin, Y.
Deposited on : 2017-08-18
Resolution : 2.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7383 (2.54-2.50)
Clashscore	190562	8079 (2.54-2.50)
Ramachandran outliers	187476	7944 (2.54-2.50)
Sidechain outliers	187428	7946 (2.54-2.50)
RSRZ outliers	180081	7387 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	205	
1	B	205	
1	C	205	
1	D	205	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6581 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Membrane protein insertase YidC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	199	Total	C	N	O	S	0	0	0
			1611	1045	258	304	4			
1	A	199	Total	C	N	O	S	0	0	0
			1611	1045	258	304	4			
1	B	199	Total	C	N	O	S	0	0	0
			1611	1045	258	304	4			
1	C	199	Total	C	N	O	S	0	0	0
			1611	1045	258	304	4			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	57	MET	LEU	engineered mutation	UNP Q9X1H2
D	105	MET	ILE	engineered mutation	UNP Q9X1H2
D	153	MET	LEU	engineered mutation	UNP Q9X1H2
D	223	HIS	-	expression tag	UNP Q9X1H2
D	224	HIS	-	expression tag	UNP Q9X1H2
D	225	HIS	-	expression tag	UNP Q9X1H2
D	226	HIS	-	expression tag	UNP Q9X1H2
D	227	HIS	-	expression tag	UNP Q9X1H2
D	228	HIS	-	expression tag	UNP Q9X1H2
A	57	MET	LEU	engineered mutation	UNP Q9X1H2
A	105	MET	ILE	engineered mutation	UNP Q9X1H2
A	153	MET	LEU	engineered mutation	UNP Q9X1H2
A	223	HIS	-	expression tag	UNP Q9X1H2
A	224	HIS	-	expression tag	UNP Q9X1H2
A	225	HIS	-	expression tag	UNP Q9X1H2
A	226	HIS	-	expression tag	UNP Q9X1H2
A	227	HIS	-	expression tag	UNP Q9X1H2
A	228	HIS	-	expression tag	UNP Q9X1H2
B	57	MET	LEU	engineered mutation	UNP Q9X1H2
B	105	MET	ILE	engineered mutation	UNP Q9X1H2
B	153	MET	LEU	engineered mutation	UNP Q9X1H2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	223	HIS	-	expression tag	UNP Q9X1H2
B	224	HIS	-	expression tag	UNP Q9X1H2
B	225	HIS	-	expression tag	UNP Q9X1H2
B	226	HIS	-	expression tag	UNP Q9X1H2
B	227	HIS	-	expression tag	UNP Q9X1H2
B	228	HIS	-	expression tag	UNP Q9X1H2
C	57	MET	LEU	engineered mutation	UNP Q9X1H2
C	105	MET	ILE	engineered mutation	UNP Q9X1H2
C	153	MET	LEU	engineered mutation	UNP Q9X1H2
C	223	HIS	-	expression tag	UNP Q9X1H2
C	224	HIS	-	expression tag	UNP Q9X1H2
C	225	HIS	-	expression tag	UNP Q9X1H2
C	226	HIS	-	expression tag	UNP Q9X1H2
C	227	HIS	-	expression tag	UNP Q9X1H2
C	228	HIS	-	expression tag	UNP Q9X1H2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	36	Total O 36 36	0	0
2	A	28	Total O 28 28	0	0
2	B	31	Total O 31 31	0	0
2	C	42	Total O 42 42	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Membrane protein insertase YidC




• Molecule 1: Membrane protein insertase YidC



• Molecule 1: Membrane protein insertase YidC



• Molecule 1: Membrane protein insertase YidC



P221
G222
HIS
HIS
HIS
HIS
HIS
HIS

4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	79.95Å 79.95Å 135.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.97 – 2.52 29.97 – 2.52	Depositor EDS
% Data completeness (in resolution range)	88.2 (29.97-2.52) 88.0 (29.97-2.52)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 2.51Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.221 , 0.261 0.230 , 0.273	Depositor DCC
R_{free} test set	1349 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	33.7	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 24.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.449 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6581	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/1646	0.49	0/2225
1	B	0.34	0/1646	0.51	0/2225
1	C	0.35	0/1646	0.53	0/2225
1	D	0.38	0/1646	0.53	0/2225
All	All	0.35	0/6584	0.51	0/8900

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1611	0	1613	28	0
1	B	1611	0	1613	30	0
1	C	1611	0	1613	20	0
1	D	1611	0	1613	30	0
2	A	28	0	0	1	1
2	B	31	0	0	2	1
2	C	42	0	0	2	0
2	D	36	0	0	4	0
All	All	6581	0	6452	106	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:38:ARG:NH1	2:D:301:HOH:O	1.94	0.98
1:D:76:ASP:HB2	1:D:81:LEU:HD21	1.57	0.84
1:C:78:GLY:O	2:C:302:HOH:O	2.00	0.79
1:C:181:THR:OG1	2:C:301:HOH:O	1.94	0.78
1:D:94:LYS:NZ	2:D:303:HOH:O	2.13	0.74
1:D:135:PRO:O	2:D:302:HOH:O	2.05	0.73
1:C:76:ASP:HB2	1:C:81:LEU:HD21	1.76	0.66
1:A:211:ASP:N	1:A:211:ASP:OD1	2.28	0.65
1:C:131:GLU:HA	1:C:186:GLY:O	1.99	0.63
1:B:77:GLU:N	2:B:303:HOH:O	2.32	0.62
1:B:210:TYR:CE1	1:B:214:ILE:HD12	2.35	0.62
1:D:106:VAL:HG22	1:D:115:ILE:HG12	1.83	0.61
1:A:54:PHE:HB3	1:A:65:PHE:HB3	1.83	0.61
1:B:109:TYR:HB2	1:B:112:VAL:HB	1.81	0.61
1:A:76:ASP:OD2	1:A:111:TYR:OH	2.14	0.60
1:C:102:GLU:HG2	1:C:119:LYS:HG2	1.83	0.60
1:D:131:GLU:HA	1:D:186:GLY:O	2.02	0.60
1:B:41:GLU:HB3	1:B:55:TYR:HB2	1.85	0.59
1:B:33:ILE:HD12	1:B:105:MET:HE3	1.85	0.59
1:A:77:GLU:N	2:A:301:HOH:O	2.37	0.58
1:D:54:PHE:HB3	1:D:65:PHE:HB3	1.84	0.58
1:C:95:VAL:HG13	1:C:97:GLU:H	1.69	0.57
1:C:28:ARG:HG3	1:C:33:ILE:HG12	1.87	0.56
1:C:54:PHE:HB3	1:C:65:PHE:HB3	1.87	0.56
1:D:29:SER:C	1:D:31:LYS:H	2.14	0.55
1:B:54:PHE:HB3	1:B:65:PHE:HB3	1.87	0.55
1:C:124:TYR:HB3	1:C:194:MET:O	2.06	0.55
1:C:41:GLU:HB3	1:C:55:TYR:HB2	1.89	0.55
1:B:124:TYR:HB3	1:B:194:MET:O	2.06	0.55
1:A:109:TYR:HB2	1:A:112:VAL:HB	1.88	0.54
1:D:41:GLU:HB3	1:D:55:TYR:HB2	1.89	0.54
1:C:72:PHE:O	1:C:114:LYS:NZ	2.41	0.54
1:A:135:PRO:HB3	1:A:185:ASN:HB2	1.89	0.54
1:D:95:VAL:HG13	1:D:97:GLU:H	1.73	0.53
1:D:102:GLU:HG2	1:D:119:LYS:HG2	1.90	0.53
1:B:48:LYS:HD3	2:B:310:HOH:O	2.06	0.53
1:B:49:GLY:HA2	1:B:107:TYR:CE1	2.44	0.53
1:D:95:VAL:HG22	1:D:96:SER:H	1.73	0.53
1:A:130:ILE:HG23	1:A:184:VAL:HG11	1.90	0.53
1:C:95:VAL:HG22	1:C:96:SER:H	1.72	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:24:ILE:HD12	1:B:120:ASN:HD21	1.75	0.52
1:A:123:ASN:O	1:A:123:ASN:ND2	2.42	0.52
1:D:85:GLU:OE1	2:D:304:HOH:O	2.18	0.52
1:D:90:THR:HG23	1:D:104:SER:HB2	1.91	0.51
1:C:90:THR:HG23	1:C:104:SER:HB2	1.92	0.51
1:A:138:VAL:HB	1:A:182:LEU:HB2	1.93	0.51
1:A:135:PRO:HA	1:A:184:VAL:O	2.11	0.51
1:A:124:TYR:HB3	1:A:194:MET:O	2.11	0.50
1:B:123:ASN:O	1:B:123:ASN:ND2	2.38	0.50
1:B:28:ARG:HG3	1:B:33:ILE:HG12	1.94	0.49
1:A:41:GLU:HB3	1:A:55:TYR:HB2	1.95	0.49
1:D:181:THR:OG1	1:D:183:LYS:NZ	2.45	0.49
1:D:123:ASN:O	1:D:123:ASN:ND2	2.41	0.48
1:D:70:ASP:HA	1:D:139:THR:OG1	2.14	0.48
1:B:138:VAL:HB	1:B:182:LEU:HB2	1.96	0.48
1:B:44:PHE:HB2	1:B:105:MET:HE1	1.95	0.48
1:A:73:ASP:O	1:A:138:VAL:HA	2.14	0.47
1:A:102:GLU:HG2	1:A:119:LYS:HG2	1.97	0.47
1:A:199:ARG:HG2	1:A:210:TYR:OH	2.15	0.47
1:C:153:MET:HE3	1:C:153:MET:HB2	1.68	0.47
1:A:90:THR:HG23	1:A:104:SER:HB2	1.97	0.47
1:B:129:ASP:OD1	1:B:189:ARG:NE	2.28	0.47
1:D:73:ASP:O	1:D:138:VAL:HA	2.15	0.46
1:A:95:VAL:HG13	1:A:97:GLU:H	1.79	0.46
1:D:101:ASP:HB3	1:B:135:PRO:HG3	1.96	0.46
1:B:95:VAL:HG22	1:B:96:SER:H	1.80	0.46
1:C:31:LYS:HB3	1:C:31:LYS:HE2	1.70	0.46
1:A:28:ARG:HG3	1:A:33:ILE:HG12	1.98	0.46
1:A:200:THR:HG22	1:A:204:LYS:HE3	1.96	0.46
1:D:49:GLY:HA2	1:D:107:TYR:CE1	2.51	0.46
1:B:76:ASP:HA	1:B:136:VAL:HG12	1.97	0.46
1:D:90:THR:OG1	1:D:91:GLY:N	2.49	0.45
1:B:146:ASP:OD2	1:B:148:SER:HB3	2.17	0.45
1:C:44:PHE:HB2	1:C:105:MET:HE1	1.98	0.45
1:D:153:MET:HB2	1:D:153:MET:HE3	1.67	0.45
1:C:176:LEU:HD23	1:C:176:LEU:HA	1.84	0.44
1:A:99:PHE:CD1	1:A:103:VAL:HG22	2.51	0.44
1:B:70:ASP:HB2	1:B:140:VAL:O	2.17	0.44
1:A:33:ILE:HD12	1:A:105:MET:HE3	1.98	0.44
1:B:28:ARG:NH2	1:B:87:PRO:O	2.44	0.44
1:B:220:ILE:HG22	1:B:221:PRO:HD2	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:33:ILE:HG13	1:B:46:LEU:HD21	2.00	0.44
1:A:95:VAL:HG22	1:A:96:SER:H	1.83	0.43
1:A:183:LYS:HD2	1:A:183:LYS:H	1.82	0.43
1:C:33:ILE:HG13	1:C:46:LEU:HD21	2.01	0.43
1:B:153:MET:HE3	1:B:153:MET:HB2	1.83	0.43
1:D:57:MET:HE3	1:D:57:MET:HB2	1.86	0.42
1:D:156:TYR:CE2	1:D:213:LEU:HD11	2.54	0.42
1:C:156:TYR:CE2	1:C:213:LEU:HD11	2.54	0.42
1:A:43:HIS:O	1:A:52:LYS:N	2.50	0.42
1:D:28:ARG:HG3	1:D:33:ILE:HG12	2.00	0.42
1:B:105:MET:O	1:B:115:ILE:HA	2.19	0.42
1:D:199:ARG:HG3	1:D:214:ILE:HD13	2.02	0.41
1:B:75:LEU:HB2	1:B:137:ASP:HB2	2.02	0.41
1:B:146:ASP:N	1:B:146:ASP:OD1	2.53	0.41
1:B:95:VAL:N	1:B:98:GLY:O	2.51	0.41
1:D:72:PHE:O	1:D:114:LYS:NZ	2.53	0.41
1:B:198:LYS:HD2	1:B:198:LYS:HA	1.89	0.41
1:B:102:GLU:HG3	1:B:119:LYS:HG2	2.03	0.41
1:D:49:GLY:HA2	1:D:107:TYR:CZ	2.56	0.41
1:D:70:ASP:HB2	1:D:140:VAL:O	2.21	0.40
1:D:168:VAL:HG22	1:D:194:MET:HG3	2.03	0.40
1:A:76:ASP:HB2	1:A:81:LEU:HD21	2.03	0.40
1:A:135:PRO:HG3	1:C:101:ASP:CG	2.46	0.40
1:A:90:THR:OG1	1:A:91:GLY:N	2.54	0.40
1:A:44:PHE:HB2	1:A:105:MET:HE1	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:324:HOH:O	2:B:328:HOH:O[3_745]	1.85	0.35

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/205 (96%)	181 (92%)	14 (7%)	2 (1%)	12	23
1	B	197/205 (96%)	182 (92%)	13 (7%)	2 (1%)	12	23
1	C	197/205 (96%)	176 (89%)	21 (11%)	0	100	100
1	D	197/205 (96%)	180 (91%)	16 (8%)	1 (0%)	24	41
All	All	788/820 (96%)	719 (91%)	64 (8%)	5 (1%)	21	36

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	76	ASP
1	A	78	GLY
1	D	30	GLU
1	B	76	ASP
1	B	94	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/188 (97%)	175 (96%)	7 (4%)	29	53
1	B	182/188 (97%)	176 (97%)	6 (3%)	33	58
1	C	182/188 (97%)	179 (98%)	3 (2%)	55	78
1	D	182/188 (97%)	175 (96%)	7 (4%)	29	53
All	All	728/752 (97%)	705 (97%)	23 (3%)	34	60

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	24	ILE
1	D	30	GLU
1	D	38	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	90	THR
1	D	102	GLU
1	D	123	ASN
1	D	221	PRO
1	A	29	SER
1	A	40	GLU
1	A	79	THR
1	A	102	GLU
1	A	211	ASP
1	A	212	VAL
1	A	214	ILE
1	B	24	ILE
1	B	40	GLU
1	B	123	ASN
1	B	199	ARG
1	B	204	LYS
1	B	220	ILE
1	C	77	GLU
1	C	90	THR
1	C	221	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	69	ASN
1	A	219	ASN
1	B	185	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	199/205 (97%)	-1.30	0 100 100	18, 33, 58, 69	0
1	B	199/205 (97%)	-1.36	0 100 100	18, 33, 59, 73	0
1	C	199/205 (97%)	-1.38	0 100 100	16, 30, 56, 86	0
1	D	199/205 (97%)	-1.36	0 100 100	18, 30, 59, 81	0
All	All	796/820 (97%)	-1.35	0 100 100	16, 32, 59, 86	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.