



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:30 PM UTC

PDB ID : 1Y9Z / pdb_00001y9z
Title : Crystal Structure of Psychrophilic Subtilisin-like Serine Protease from Antarctic Psychrotroph Pseudoalteromonas sp. AS-11 at 0.14 nm resolution
Authors : Dong, D.; Watanabe, K.
Deposited on : 2004-12-16
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	441	11% 85% 13% .
1	B	441	4% 89% 9% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PMS	A	601	-	X	-	-
5	PMS	B	602	-	X	-	-
6	TRS	A	611	-	X	-	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6853 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alkaline serine protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	435	Total	C	N	O	S	0	0	0
			3129	1911	548	656	14			
1	B	436	Total	C	N	O	S	0	0	0
			3136	1915	549	658	14			


- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		
2	B	2	Total	Ca	0	0
			2	2		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Na	0	0
			2	2		
3	B	2	Total	Na	0	0
			2	2		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is phenylmethanesulfonic acid (CCD ID: PMS) (formula: $C_7H_8O_3S$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	S	0	0
			10	7	2	1		
5	B	1	Total	C	O	S	0	0
			10	7	2	1		

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	237	Total	O	0	0
			237	237		
7	B	305	Total	O	0	0
			305	305		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: alkaline serine protease

Chain A: 



- Molecule 1: alkaline serine protease

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	122.09Å 138.32Å 64.79Å 90.00° 97.49° 90.00°	Depositor
Resolution (Å)	34.58 – 1.40 34.58 – 1.40	Depositor EDS
% Data completeness (in resolution range)	91.2 (34.58-1.40) 91.3 (34.58-1.40)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.11 (at 1.40Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.197 , 0.207 0.190 , 0.200	Depositor DCC
R_{free} test set	19440 reflections (9.94%)	wwPDB-VP
Wilson B-factor (Å ²)	10.7	Xtriage
Anisotropy	0.240	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 31.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6853	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PMS, SO4, TRS, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.33	0/3185	0.89	12/4356 (0.3%)
1	B	0.34	0/3192	0.91	10/4366 (0.2%)
All	All	0.34	0/6377	0.90	22/8722 (0.3%)

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	76	ILE	N-CA-C	6.86	118.54	109.21
1	A	369	SER	CA-CB-OG	6.82	124.74	111.10
1	A	124	ASN	N-CA-C	-6.77	105.14	113.41
1	B	369	SER	CA-CB-OG	6.64	124.39	111.10
1	B	276	ILE	N-CA-C	-6.43	98.60	107.99
1	B	124	ASN	N-CA-C	-6.28	105.74	113.41
1	A	76	ILE	N-CA-C	6.11	117.47	109.58
1	B	177	VAL	N-CA-C	5.95	116.87	108.36
1	A	74	ALA	N-CA-C	5.88	119.74	112.03
1	A	232	VAL	N-CA-C	5.77	114.23	107.77
1	B	368	THR	N-CA-C	-5.74	105.54	112.54
1	A	88	ASN	N-CA-C	5.72	119.24	112.72
1	B	88	ASN	N-CA-C	5.70	119.01	112.57
1	A	100	ASN	N-CA-C	-5.53	102.01	110.14
1	A	368	THR	N-CA-C	-5.45	105.89	112.54
1	B	100	ASN	N-CA-C	-5.41	102.19	110.14
1	B	74	ALA	N-CA-C	5.37	119.06	112.03
1	B	232	VAL	N-CA-C	5.30	114.31	108.05
1	A	177	VAL	N-CA-C	5.23	115.84	108.36
1	A	276	ILE	N-CA-C	-5.10	100.54	107.99
1	A	42	ALA	N-CA-C	5.08	118.62	112.93
1	A	183	ASN	N-CA-C	-5.03	106.39	112.88

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3129	0	2958	39	0
1	B	3136	0	2965	28	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	A	10	0	7	0	0
5	B	10	0	7	0	0
6	A	8	0	12	0	0
7	A	237	0	0	2	0
7	B	305	0	0	0	0
All	All	6853	0	5949	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:37:HIS:HD2	1:B:39:ASP:H	1.14	0.92
1:A:37:HIS:HD2	1:A:39:ASP:H	1.16	0.90
1:A:160:ALA:H	1:A:191:GLN:HE21	1.25	0.84
1:B:159:ASN:HD21	1:B:368:THR:H	1.23	0.83
1:B:160:ALA:H	1:B:191:GLN:HE21	1.27	0.81
1:A:23:ASN:HD22	1:A:23:ASN:C	1.93	0.75
1:A:5:PRO:HG2	1:A:8:GLN:HG2	1.68	0.73
1:A:23:ASN:ND2	1:A:24:ARG:HG2	2.05	0.70
1:B:23:ASN:C	1:B:23:ASN:HD22	1.97	0.70
1:A:204:ALA:H	1:A:366:ASN:ND2	1.92	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:319:ASN:HB3	1:B:366:ASN:HD21	1.59	0.68
1:B:23:ASN:ND2	1:B:24:ARG:HG2	2.09	0.68
1:B:204:ALA:H	1:B:366:ASN:ND2	1.92	0.68
1:A:274:ASN:HA	1:A:304:LYS:HD2	1.76	0.67
1:B:390:SER:H	1:B:393:GLN:HE21	1.43	0.66
1:A:319:ASN:HB3	1:A:366:ASN:HD21	1.61	0.65
1:B:5:PRO:HG2	1:B:8:GLN:HG2	1.79	0.65
1:A:274:ASN:HA	1:A:304:LYS:CD	2.27	0.64
1:B:37:HIS:HE1	1:B:361:ASP:O	1.84	0.61
1:A:37:HIS:HE1	1:A:361:ASP:O	1.86	0.59
1:A:260:CYS:SG	1:A:267:PHE:HB3	2.43	0.59
1:B:204:ALA:H	1:B:366:ASN:HD22	1.51	0.57
1:A:204:ALA:H	1:A:366:ASN:HD22	1.53	0.57
1:B:390:SER:H	1:B:393:GLN:NE2	2.03	0.56
1:B:37:HIS:CD2	1:B:39:ASP:H	2.07	0.56
1:A:390:SER:H	1:A:393:GLN:NE2	2.05	0.55
1:B:234:HIS:HD2	1:B:333:SER:OG	1.90	0.55
1:A:390:SER:H	1:A:393:GLN:HE21	1.56	0.54
1:B:23:ASN:HD21	1:B:384:SER:HA	1.72	0.54
1:A:37:HIS:CD2	1:A:39:ASP:H	2.09	0.53
1:A:238:THR:HB	1:A:239:PRO:HD2	1.91	0.53
1:A:234:HIS:HD2	1:A:333:SER:OG	1.92	0.52
1:A:68:HIS:HA	1:A:208:THR:O	2.10	0.52
1:B:274:ASN:HA	1:B:304:LYS:HD2	1.93	0.51
1:B:68:HIS:HA	1:B:208:THR:O	2.11	0.51
1:A:55:ASN:HD22	1:A:58:GLN:NE2	2.08	0.51
1:A:23:ASN:HD21	1:A:384:SER:HA	1.77	0.50
1:A:234:HIS:HE1	1:A:291:GLU:OE1	1.94	0.49
1:A:256:ALA:HB3	1:A:275:LYS:HD3	1.94	0.49
1:B:234:HIS:HE1	1:B:291:GLU:OE1	1.96	0.48
1:B:23:ASN:C	1:B:23:ASN:ND2	2.67	0.47
1:B:160:ALA:H	1:B:191:GLN:NE2	2.05	0.46
1:B:430:GLU:OE2	1:B:435:PRO:HA	2.14	0.46
1:B:23:ASN:HD22	1:B:24:ARG:HG2	1.79	0.45
1:B:23:ASN:HD22	1:B:24:ARG:N	2.15	0.44
1:A:23:ASN:HD22	1:A:24:ARG:HG2	1.78	0.44
1:A:23:ASN:HD21	1:A:24:ARG:HG2	1.80	0.44
1:A:257:LEU:HG	1:A:348:LEU:HD23	1.99	0.43
1:A:12:GLY:O	1:A:421:ALA:HB3	2.18	0.43
1:B:33:TYR:CZ	1:B:40:LEU:HB3	2.53	0.43
1:A:385:TYR:CD2	1:A:425:LYS:HE2	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:ARG:NH1	7:A:837:HOH:O	2.47	0.42
1:B:130:LEU:C	1:B:130:LEU:HD12	2.44	0.42
1:A:224:GLN:NE2	1:A:224:GLN:HA	2.34	0.42
1:A:295:THR:HG22	1:A:329:ILE:HD12	2.02	0.42
1:B:159:ASN:ND2	1:B:368:THR:H	2.04	0.42
1:A:237:LEU:HD11	1:A:321:PHE:CD2	2.55	0.41
1:A:234:HIS:HD2	1:A:333:SER:CB	2.33	0.41
1:A:130:LEU:HD12	1:A:130:LEU:C	2.45	0.41
1:A:146:HIS:HD2	7:A:794:HOH:O	2.03	0.41
1:A:23:ASN:HD22	1:A:24:ARG:N	2.18	0.41
1:A:179:ALA:HB1	1:A:202:GLY:HA3	2.03	0.41
1:A:256:ALA:HB3	1:A:275:LYS:CD	2.51	0.41
1:A:69:VAL:HG13	1:A:373:PRO:HB3	2.02	0.40
1:B:238:THR:HB	1:B:239:PRO:HD2	2.03	0.40
1:A:23:ASN:C	1:A:23:ASN:ND2	2.66	0.40
1:B:69:VAL:HG13	1:B:373:PRO:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	433/441 (98%)	416 (96%)	17 (4%)	0	100	100
1	B	434/441 (98%)	420 (97%)	14 (3%)	0	100	100
All	All	867/882 (98%)	836 (96%)	31 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/339 (99%)	332 (99%)	3 (1%)	70	44
1	B	336/339 (99%)	332 (99%)	4 (1%)	63	34
All	All	671/678 (99%)	664 (99%)	7 (1%)	68	40

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	ASN
1	A	87	PRO
1	A	234	HIS
1	B	23	ASN
1	B	87	PRO
1	B	234	HIS
1	B	311	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (40) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	ASN
1	A	37	HIS
1	A	58	GLN
1	A	62	ASN
1	A	88	ASN
1	A	89	GLN
1	A	90	ASN
1	A	141	ASN
1	A	144	ASN
1	A	146	HIS
1	A	191	GLN
1	A	224	GLN
1	A	234	HIS
1	A	235	ASN
1	A	274	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	319	ASN
1	A	366	ASN
1	A	393	GLN
1	A	399	ASN
1	A	412	ASN
1	A	413	GLN
1	B	23	ASN
1	B	37	HIS
1	B	44	ASN
1	B	90	ASN
1	B	92	ASN
1	B	141	ASN
1	B	159	ASN
1	B	191	GLN
1	B	229	ASN
1	B	234	HIS
1	B	235	ASN
1	B	274	ASN
1	B	311	ASN
1	B	319	ASN
1	B	357	GLN
1	B	366	ASN
1	B	393	GLN
1	B	412	ASN
1	B	413	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 13 ligands modelled in this entry, 8 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	PMS	B	602	1	7,10,11	3.15	6 (85%)	11,12,15	1.97	6 (54%)
4	SO4	B	613	-	4,4,4	0.38	0	6,6,6	0.07	0
4	SO4	A	612	-	4,4,4	0.37	0	6,6,6	0.05	0
6	TRS	A	611	-	7,7,7	1.25	1 (14%)	9,9,9	6.34	6 (66%)
5	PMS	A	601	1	7,10,11	3.22	6 (85%)	11,12,15	1.96	6 (54%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PMS	B	602	1	-	1/4/4/5	0/1/1/1
6	TRS	A	611	-	-	6/9/9/9	-
5	PMS	A	601	1	-	1/4/4/5	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	601	PMS	C6-C1	5.15	1.49	1.38
5	B	602	PMS	C6-C1	5.07	1.48	1.38
5	A	601	PMS	C4-C3	3.24	1.45	1.38
5	B	602	PMS	C4-C3	3.19	1.45	1.38
5	A	601	PMS	C-C1	3.09	1.55	1.50
5	B	602	PMS	C2-C1	2.92	1.44	1.38
5	A	601	PMS	C2-C1	2.90	1.44	1.38
5	B	602	PMS	C-C1	2.86	1.55	1.50
5	A	601	PMS	C3-C2	2.72	1.43	1.38
5	B	602	PMS	C3-C2	2.71	1.43	1.38
5	A	601	PMS	C5-C6	2.66	1.43	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	602	PMS	C5-C6	2.48	1.43	1.38
6	A	611	TRS	O1-C1	2.01	1.48	1.42

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	611	TRS	C2-C-N	-11.12	79.81	108.17
6	A	611	TRS	C1-C-N	-9.42	84.15	108.17
6	A	611	TRS	C3-C-N	-8.18	87.32	108.17
6	A	611	TRS	C2-C-C1	7.01	129.31	110.66
6	A	611	TRS	O1-C1-C	4.25	122.71	110.88
5	A	601	PMS	C5-C6-C1	-3.35	115.89	120.61
5	B	602	PMS	C5-C6-C1	-3.33	115.92	120.61
6	A	611	TRS	O2-C2-C	3.01	119.28	110.88
5	B	602	PMS	C6-C1-C2	2.68	122.21	118.23
5	A	601	PMS	C6-C1-C2	2.65	122.17	118.23
5	B	602	PMS	O1S-S-C	2.58	110.01	105.55
5	A	601	PMS	O1S-S-C	2.44	109.76	105.55
5	B	602	PMS	C-C1-C2	-2.28	118.08	120.56
5	A	601	PMS	C-C1-C2	-2.24	118.11	120.56
5	B	602	PMS	C4-C3-C2	-2.10	117.64	120.24
5	B	602	PMS	C5-C4-C3	2.06	122.69	119.87
5	A	601	PMS	C4-C3-C2	-2.05	117.72	120.24
5	A	601	PMS	C5-C4-C3	2.02	122.64	119.87

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	601	PMS	C1-C-S-O1S
5	B	602	PMS	C1-C-S-O1S
6	A	611	TRS	C2-C-C1-O1
6	A	611	TRS	N-C-C1-O1
6	A	611	TRS	C1-C-C3-O3
6	A	611	TRS	C2-C-C3-O3
6	A	611	TRS	C1-C-C2-O2
6	A	611	TRS	N-C-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	435/441 (98%)	0.57	49 (11%)	10 10	8, 14, 28, 38	0
1	B	436/441 (98%)	0.06	19 (4%)	39 40	6, 10, 20, 36	0
All	All	871/882 (98%)	0.31	68 (7%)	19 20	6, 12, 26, 38	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	242	THR	7.0
1	B	436	THR	5.9
1	A	13	ALA	5.5
1	A	15	VAL	4.9
1	B	1	ALA	4.8
1	A	241	GLY	4.7
1	A	268	SER	4.3
1	B	241	GLY	4.2
1	B	242	THR	4.2
1	A	260	CYS	4.0
1	A	262	VAL	3.9
1	B	244	TYR	3.9
1	A	273	ALA	3.9
1	B	243	SER	3.8
1	A	239	PRO	3.8
1	B	239	PRO	3.8
1	A	261	THR	3.6
1	A	267	PHE	3.6
1	A	349	GLY	3.6
1	A	79	ASN	3.6
1	A	265	THR	3.5
1	A	269	CYS	3.5
1	A	12	GLY	3.4
1	A	266	SER	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	301	ALA	3.3
1	A	275	LYS	3.2
1	B	274	ASN	3.2
1	A	435	PRO	3.2
1	A	271	ASN	3.1
1	A	270	GLY	3.0
1	B	245	ALA	3.0
1	A	346	ALA	2.9
1	A	243	SER	2.9
1	A	1	ALA	2.8
1	A	427	TYR	2.8
1	B	246	PRO	2.7
1	B	293	ASN	2.6
1	A	264	GLY	2.6
1	A	325	ALA	2.5
1	A	274	ASN	2.5
1	A	240	SER	2.5
1	A	256	ALA	2.4
1	A	223	GLY	2.4
1	A	328	ASP	2.3
1	A	14	THR	2.3
1	A	342	LEU	2.3
1	A	302	GLY	2.3
1	B	304	LYS	2.3
1	A	263	ASN	2.3
1	A	293	ASN	2.3
1	A	276	ILE	2.3
1	A	300	THR	2.2
1	B	323	VAL	2.2
1	A	258	ALA	2.2
1	A	433	THR	2.2
1	B	325	ALA	2.1
1	A	312	SER	2.1
1	B	240	SER	2.1
1	A	287	SER	2.1
1	A	279	VAL	2.1
1	A	282	VAL	2.1
1	A	246	PRO	2.1
1	A	348	LEU	2.1
1	B	238	THR	2.0
1	A	244	TYR	2.0
1	B	80	GLU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	237	LEU	2.0
1	B	271	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	TRS	A	611	8/8	0.76	0.21	23,26,28,30	0
5	PMS	A	601	10/11	0.86	0.20	23,29,30,31	0
5	PMS	B	602	10/11	0.88	0.16	20,24,25,26	0
4	SO4	A	612	5/5	0.91	0.13	29,31,31,32	0
4	SO4	B	613	5/5	0.97	0.12	19,20,21,22	0
2	CA	A	604	1/1	0.98	0.04	12,12,12,12	0
2	CA	B	605	1/1	0.99	0.04	8,8,8,8	0
2	CA	B	606	1/1	0.99	0.02	11,11,11,11	0
3	NA	A	607	1/1	0.99	0.04	8,8,8,8	0
3	NA	A	608	1/1	0.99	0.05	17,17,17,17	0
3	NA	B	610	1/1	0.99	0.07	15,15,15,15	0
2	CA	A	603	1/1	1.00	0.03	8,8,8,8	0
3	NA	B	609	1/1	1.00	0.04	7,7,7,7	0

6.5 Other polymers [i](#)

There are no such residues in this entry.