



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 30, 2026 – 01:00 AM UTC

PDB ID : 2Y95 / pdb_00002y95
BMRB ID : 16714
Title : Solution structure of AUCG tetraloop hairpin found in human Xist RNA A-repeats essential for X-inactivation
Authors : Duszczuk, M.M.; Sattler, M.
Deposited on : 2011-02-11

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 74%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
RNA backbone	8273	777

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	14	

2 Ensemble composition and analysis ⓘ

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 452 atoms, of which 155 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called 5'-R(*GP*GP*CP*GP*CP*AP*UP*CP*GP*GP*CP*GP*CP*C)-3'.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	14	452	133	155	55	96	13	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4	G	C	engineered mutation	GB 340393
A	11	C	G	engineered mutation	GB 340393
A	14	C	U	engineered mutation	GB 340393


4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 5'-R(*GP*GP*CP*GP*CP*AP*UP*CP*GP*GP*CP*GP*CP*C)-3'

Chain A: 



4.2 Residue scores for the first model from the NMR ensemble

No representative models were identified. Colouring as in section 4.1 above.

- Molecule 1: 5'-R(*GP*GP*CP*GP*CP*AP*UP*CP*GP*GP*CP*GP*CP*C)-3'

Chain A: 



5 Refinement protocol and experimental data overview

The models were refined using the following method: *ARIA*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *LEAST RESTRAINT VIOLATION, LOW ENERGY, AGREEMENT WITH CHEMICAL SHIFTS CALCULATED BY NUCHEMICS*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	refinement	
NMRPipe	structure solution	
NMRView	structure solution	
ARIA	structure solution	1.2
Amber	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	251
Number of shifts mapped to atoms	251
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	74%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.67±0.01	0±0/331 (0.0± 0.0%)	1.00±0.02	1±0/515 (0.1± 0.1%)
All	All	0.67	0/3310 (0.0%)	1.00	6/5150 (0.1%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	10	G	C5'-C4'-C3'	-5.91	107.13	116.00	7	4
1	A	9	G	C5'-C4'-C3'	-5.12	107.52	115.20	6	1
1	A	10	G	O4'-C4'-C3'	5.04	109.04	104.00	9	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	2970	1550	1550	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	13/14 (93%)	3±0 (23±0%)	1±0 (9±3%)	0.69±0.00
All	All	130/140 (93%)	30 (23%)	12 (9%)	0.69

The overall RNA backbone suiteness is 0.58.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	8	C	10
1	A	9	G	10
1	A	10	G	10

All unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	9	G	10
1	A	10	G	2

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry

There are no ligands in this entry.

6.7 Other polymers

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 74% for the well-defined parts and 74% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	251
Number of shifts mapped to atoms	251
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 74%, i.e. 208 atoms were assigned a chemical shift out of a possible 280. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Sugar	151/154 (98%)	82/84 (98%)	69/70 (99%)	0/0 (—%)
Base	57/126 (45%)	21/74 (28%)	21/22 (95%)	15/30 (50%)
Overall	208/280 (74%)	103/158 (65%)	90/92 (98%)	15/30 (50%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots

No *random coil index*(RCI) plot could be generated from the current chemical shift list. RCI is only applicable to proteins

8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	297
Intra-residue ($ i-j =0$)	162
Sequential ($ i-j =1$)	92
Medium range ($ i-j >1$ and $ i-j <5$)	10
Long range ($ i-j \geq 5$)	8
Inter-chain	0
Hydrogen bond restraints	25
Disulfide bond restraints	0
Total dihedral-angle restraints	96
Number of unmapped restraints	0
Number of restraints per residue	28.1
Number of long range restraints per residue ¹	2.4

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	4.1	0.2
0.2-0.5 (Medium)	0.5	0.36
>0.5 (Large)	12.0	1.76

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	Max (°)
1.0-10.0 (Small)	5.2	5.97
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None

9 Distance violation analysis ⓘ


9.1 Summary of distance violations ⓘ

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue ($i-j =0$)	162	54.5	2	1.2	0.7	0	0.0	0.0
Backbone-Backbone	94	31.6	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	61	20.5	2	3.3	0.7	0	0.0	0.0
Sidechain-Sidechain	7	2.4	0	0.0	0.0	0	0.0	0.0
Sequential ($i-j =1$)	92	31.0	5	5.4	1.7	2	2.2	0.7
Backbone-Backbone	32	10.8	3	9.4	1.0	2	6.2	0.7
Backbone-Sidechain	53	17.8	2	3.8	0.7	0	0.0	0.0
Sidechain-Sidechain	7	2.4	0	0.0	0.0	0	0.0	0.0
Medium range ($i-j >1$ & $i-j <5$)	10	3.4	1	10.0	0.3	0	0.0	0.0
Backbone-Backbone	2	0.7	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	6	2.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	2	0.7	1	50.0	0.3	0	0.0	0.0
Long range ($i-j \geq 5$)	8	2.7	2	25.0	0.7	2	25.0	0.7
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	2	0.7	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	6	2.0	2	33.3	0.7	2	33.3	0.7
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	25	8.4	10	40.0	3.4	10	40.0	3.4
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	297	100.0	20	6.7	6.7	14	4.7	4.7
Backbone-Backbone	128	43.1	3	2.3	1.0	2	1.6	0.7
Backbone-Sidechain	122	41.1	4	3.3	1.3	0	0.0	0.0
Sidechain-Sidechain	47	15.8	13	27.7	4.4	12	25.5	4.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis


9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	1	4	0	12	0	17	1.11	1.7	0.61	1.52
2	1	3	0	12	0	16	1.17	1.62	0.59	1.52
3	1	4	1	12	0	18	1.06	1.76	0.66	1.49
4	0	3	1	12	0	16	1.16	1.65	0.62	1.52
5	0	3	1	12	0	16	1.14	1.66	0.62	1.5
6	1	4	0	12	0	17	1.11	1.65	0.63	1.5
7	1	4	1	12	0	18	1.05	1.76	0.65	1.47
8	1	3	0	12	0	16	1.18	1.67	0.59	1.51
9	0	4	0	12	0	16	1.19	1.73	0.62	1.54
10	0	4	1	12	0	17	1.08	1.61	0.63	1.47

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 262(IR:160, SQ:87, MR:9, LR:6, IC:0) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
0	0	0	0	0	0	1	10.0
1	1	0	0	0	2	2	20.0
0	0	0	0	0	0	3	30.0


Continued on next page...

Continued from previous page...

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
1	0	0	0	0	1	4	40.0
0	0	1	0	0	1	5	50.0
0	0	0	0	0	0	6	60.0
0	2	0	0	0	2	7	70.0
0	0	0	0	0	0	8	80.0
0	0	0	0	0	0	9	90.0
0	2	0	2	0	4	10	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶ Number of models with violations

9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)




9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models

in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(2,9)	1:2:A:G:O6	1:13:A:C:H42	10	1.62	0.07	1.63
(1,270)	1:4:A:G:H1	1:11:A:C:H42	10	1.62	0.03	1.62
(2,21)	1:5:A:C:H42	1:10:A:G:O6	10	1.61	0.09	1.59
(2,25)	1:5:A:C:O2	1:10:A:G:H22	10	1.58	0.05	1.59
(2,15)	1:3:A:C:O2	1:12:A:G:H22	10	1.58	0.05	1.58
(2,11)	1:3:A:C:H42	1:12:A:G:O6	10	1.57	0.05	1.57
(2,4)	1:1:A:G:O6	1:14:A:C:H42	10	1.56	0.03	1.56
(2,2)	1:1:A:G:H22	1:14:A:C:O2	10	1.52	0.03	1.52
(2,7)	1:2:A:G:H22	1:13:A:C:O2	10	1.52	0.04	1.52
(2,17)	1:4:A:G:H22	1:11:A:C:O2	10	1.51	0.05	1.52

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(2,21)	1:5:A:C:H42	1:10:A:G:O6	3	1.76
(2,21)	1:5:A:C:H42	1:10:A:G:O6	7	1.76
(2,9)	1:2:A:G:O6	1:13:A:C:H42	9	1.73
(2,9)	1:2:A:G:O6	1:13:A:C:H42	1	1.7
(2,9)	1:2:A:G:O6	1:13:A:C:H42	3	1.7
(1,270)	1:4:A:G:H1	1:11:A:C:H42	8	1.67
(2,15)	1:3:A:C:O2	1:12:A:G:H22	5	1.66
(2,25)	1:5:A:C:O2	1:10:A:G:H22	5	1.65
(2,21)	1:5:A:C:H42	1:10:A:G:O6	9	1.65
(2,9)	1:2:A:G:O6	1:13:A:C:H42	6	1.65

10 Dihedral-angle violation analysis [i](#)


10.1 Summary of dihedral-angle violations [i](#)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
ZETA	10	10.4	0	0.0	0.0	0	0.0	0.0
EPSILON	9	9.4	3	33.3	3.1	0	0.0	0.0
CHI	14	14.6	1	7.1	1.0	0	0.0	0.0
DELTA	11	11.5	0	0.0	0.0	0	0.0	0.0
UNNAMED	22	22.9	1	4.5	1.0	0	0.0	0.0
GAMMA	12	12.5	3	25.0	3.1	0	0.0	0.0
ALPHA	10	10.4	6	60.0	6.2	0	0.0	0.0
BETA	8	8.3	0	0.0	0.0	0	0.0	0.0
Total	96	100.0	14	14.6	14.6	0	0.0	0.0

¹ percentage calculated with respect to total number of dihedral-angle restraints, ² percentage calculated with respect to number of restraints in a particular dihedral-angle type, ³ violated in at least one model, ⁴ violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations [i](#)




Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model ⓘ

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Number of violations									Total	Mean (°)	Max (°)	SD (°)	Median (°)
	ZETA	EPSILON	CHI	DELTA	UNNAMED	GAMMA	ALPHA	BETA						
1	0	2	0	0	0	2	2	0	6	2.01	3.63	1.02	1.44	
2	0	1	1	0	1	0	2	0	5	1.86	2.28	0.3	1.74	
3	0	1	1	0	1	0	3	0	6	2.48	5.97	1.85	1.32	
4	0	0	1	0	1	1	2	0	5	2.54	3.41	0.74	2.44	
5	0	0	1	0	1	0	2	0	4	1.8	2.63	0.5	1.61	
6	0	1	0	0	1	0	3	0	5	1.82	2.43	0.5	1.79	
7	0	0	1	0	1	1	5	0	8	2.96	5.01	1.29	2.6	
8	0	1	0	0	1	1	2	0	5	2.04	3.66	0.95	1.82	
9	0	0	1	0	0	2	0	0	3	1.92	2.07	0.2	2.05	
10	0	1	1	0	1	0	2	0	5	2.65	3.69	0.93	3.22	

10.2.1 Bar graph : Dihedral violation statistics for each model ⓘ



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble ⓘ


Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated

restraints for a given fraction of ensemble.

Number of violated restraints									Fraction of the ensemble	
ZETA	EPSILON	CHI	DELTA	UNNAMED	GAMMA	ALPHA	BETA	Total	Count ¹	%
0	1	0	0	0	0	0	0	1	1	10.0
0	1	0	0	0	2	1	0	4	2	20.0
0	0	0	0	0	1	3	0	4	3	30.0
0	1	0	0	0	0	0	0	1	4	40.0
0	0	0	0	0	0	1	0	1	5	50.0
0	0	0	0	0	0	0	0	0	6	60.0
0	0	1	0	0	0	1	0	2	7	70.0
0	0	0	0	1	0	0	0	1	8	80.0
0	0	0	0	0	0	0	0	0	9	90.0
0	0	0	0	0	0	0	0	0	10	100.0

¹ Number of models with violations

10.3.1 Bar graph : Dihedral-angle Violation statistics for the ensemble [i](#)




10.4 Most violated dihedral-angle restraints in the ensemble [i](#)

10.4.1 Histogram : Distribution of mean dihedral-angle violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models

in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.


Key	Atom-1	Atom-2	Atom-3	Atom-4	Models ¹	Mean	SD ²	Median
(1,69)	1:7:A:U:H2'	1:7:A:U:C2'	1:7:A:U:C3'	1:7:A:U:H3'	8	3.39	1.19	3.14
(1,9)	1:2:A:G:O3'	1:3:A:C:P	1:3:A:C:O5'	1:3:A:C:C5'	7	2.18	0.86	1.74
(1,90)	1:8:A:C:O4'	1:8:A:C:Cl'	1:8:A:C:N1	1:8:A:C:C2	7	2.02	1.02	1.64
(1,42)	1:12:A:G:O3'	1:13:A:C:P	1:13:A:C:O5'	1:13:A:C:C5'	5	2.71	1.32	2.35
(1,17)	1:4:A:G:C4'	1:4:A:G:C3'	1:4:A:G:O3'	1:5:A:C:P	4	1.48	0.47	1.26
(1,14)	1:3:A:C:O3'	1:4:A:G:P	1:4:A:G:O5'	1:4:A:G:C5'	3	2.43	1.09	1.79
(1,21)	1:5:A:C:O5'	1:5:A:C:C5'	1:5:A:C:C4'	1:5:A:C:C3'	3	2.02	0.69	2.07
(1,37)	1:11:A:C:O3'	1:12:A:G:P	1:12:A:G:O5'	1:12:A:G:C5'	3	1.72	0.25	1.82
(1,47)	1:13:A:C:O3'	1:14:A:C:P	1:14:A:C:O5'	1:14:A:C:C5'	3	1.49	0.1	1.53
(1,4)	1:1:A:G:O3'	1:2:A:G:P	1:2:A:G:O5'	1:2:A:G:C5'	2	2.88	0.53	2.88

¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints [i](#)

10.5.1 Histogram : Distribution of violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints [i](#)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,69)	1:7:A:U:H2'	1:7:A:U:C2'	1:7:A:U:C3'	1:7:A:U:H3'	3	5.97
(1,42)	1:12:A:G:O3'	1:13:A:C:P	1:13:A:C:O5'	1:13:A:C:C5'	7	5.01
(1,90)	1:8:A:C:O4'	1:8:A:C:C1'	1:8:A:C:N1	1:8:A:C:C2	7	4.35
(1,69)	1:7:A:U:H2'	1:7:A:U:C2'	1:7:A:U:C3'	1:7:A:U:H3'	7	4.15
(1,14)	1:3:A:C:O3'	1:4:A:G:P	1:4:A:G:O5'	1:4:A:G:C5'	3	3.97
(1,69)	1:7:A:U:H2'	1:7:A:U:C2'	1:7:A:U:C3'	1:7:A:U:H3'	10	3.69
(1,69)	1:7:A:U:H2'	1:7:A:U:C2'	1:7:A:U:C3'	1:7:A:U:H3'	8	3.66
(1,9)	1:2:A:G:O3'	1:3:A:C:P	1:3:A:C:O5'	1:3:A:C:C5'	1	3.63
(1,4)	1:1:A:G:O3'	1:2:A:G:P	1:2:A:G:O5'	1:2:A:G:C5'	4	3.41
(1,12)	1:3:A:C:C4'	1:3:A:C:C3'	1:3:A:C:O3'	1:4:A:G:P	10	3.28