



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:46 AM UTC

PDB ID : 5Y90 / pdb_00005y90
Title : MAP2K7 mutant -C218S
Authors : Kinoshita, T.; Hashimoto, T.; Sogabe, Y.; Matsumoto, T.; Sawa, M.; Fukada, H.
Deposited on : 2017-08-22
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1553 (1.30-1.30)
Clashscore	190562	1595 (1.30-1.30)
Ramachandran outliers	187476	1551 (1.30-1.30)
Sidechain outliers	187428	1551 (1.30-1.30)
RSRZ outliers	180081	1549 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	503	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2925 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dual specificity mitogen-activated protein kinase kinase 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	304	2560	1644	441	453	22	0	23	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	218	SER	CYS	engineered mutation	UNP O14733
A	436	HIS	-	expression tag	UNP O14733
A	437	HIS	-	expression tag	UNP O14733
A	438	HIS	-	expression tag	UNP O14733
A	439	HIS	-	expression tag	UNP O14733
A	440	HIS	-	expression tag	UNP O14733
A	441	HIS	-	expression tag	UNP O14733

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	341	Total	O	0	0
			341	341		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dual specificity mitogen-activated protein kinase kinase 7



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.40Å 70.61Å 92.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.05 – 1.30 56.05 – 1.30	Depositor EDS
% Data completeness (in resolution range)	91.7 (56.05-1.30) 91.7 (56.05-1.30)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.61 (at 1.30Å)	Xtriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.127 , 0.167 (Not available) , 0.189	Depositor DCC
R_{free} test set	3853 reflections (4.56%)	wwPDB-VP
Wilson B-factor (Å ²)	13.1	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 71.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2925	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.34	16/2669 (0.6%)	1.15	5/3585 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	218	SER	CB-OG	10.35	1.62	1.42
1	A	276	CYS	CB-SG	-8.80	1.52	1.81
1	A	338	TYR	CA-CB	7.25	1.62	1.53
1	A	134	ASN	CA-C	6.63	1.61	1.52
1	A	425	VAL	C-O	-6.38	1.17	1.24
1	A	172	ASN	C-O	5.92	1.31	1.23
1	A	363	MET	C-O	-5.82	1.16	1.24
1	A	251	GLU	C-O	-5.75	1.17	1.24
1	A	295	GLY	N-CA	5.57	1.50	1.45
1	A	298	ALA	N-CA	-5.56	1.39	1.46
1	A	197	GLN	CD-NE2	5.50	1.44	1.33
1	A	316	ASP	C-O	5.48	1.30	1.24
1	A	425	VAL	CA-C	5.47	1.59	1.52
1	A	131	ALA	C-O	-5.42	1.17	1.23
1	A	363	MET	CG-SD	5.37	1.94	1.80
1	A	191	ASP	CG-OD2	5.20	1.35	1.25

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	307	PRO	O-C-N	8.90	125.33	121.15
1	A	135	ASP	N-CA-C	6.17	117.68	111.07
1	A	314	ASP	N-CA-C	5.69	122.92	110.80
1	A	363	MET	CA-CB-CG	5.33	124.76	114.10
1	A	425	VAL	N-CA-C	5.01	115.07	107.80

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	146	THR	Peptide
1	A	147	CYS	Peptide
1	A	315	TYR	Peptide
1	A	362	HIS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2560	0	2651	78	0
2	A	24	0	32	11	0
3	A	341	0	0	27	2
All	All	2925	0	2683	79	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:270[B]:ARG:NH2	1:A:272[B]:GLN:NE2	1.61	1.46
1:A:357[B]:PRO:O	1:A:358[B]:LEU:HD23	1.58	1.03
1:A:216[A]:GLY:HA2	3:A:604:HOH:O	1.64	0.97
1:A:312:LYS:HE2	3:A:612:HOH:O	1.68	0.94
1:A:132:GLU:O	1:A:135:ASP:HB3	1.74	0.86
1:A:135:ASP:CG	2:A:503:GOL:H2	1.99	0.86

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:270[B]:ARG:CZ	3:A:601:HOH:O	2.26	0.83
1:A:270[B]:ARG:NH2	3:A:601:HOH:O	2.11	0.82
1:A:133:ILE:HD12	1:A:136:LEU:HD12	1.67	0.76
1:A:190:HIS:HA	1:A:197:GLN:OE1	1.86	0.75
1:A:217[A]:THR:HG22	1:A:221:LYS:HB2	1.70	0.74
1:A:270[B]:ARG:CZ	1:A:272[B]:GLN:NE2	2.50	0.72
1:A:270[B]:ARG:NH1	3:A:601:HOH:O	2.24	0.68
1:A:135:ASP:OD1	3:A:602:HOH:O	2.14	0.65
1:A:217[A]:THR:CG2	1:A:221:LYS:HB2	2.27	0.65
1:A:181[A]:MET:HE3	3:A:777:HOH:O	1.97	0.64
1:A:135:ASP:HB2	2:A:503:GOL:C3	2.28	0.64
1:A:312:LYS:CE	3:A:612:HOH:O	2.34	0.62
1:A:314:ASP:O	1:A:316:ASP:N	2.33	0.61
1:A:337:PRO:HB2	1:A:357[A]:PRO:HB2	1.83	0.60
1:A:317:ILE:HA	3:A:633:HOH:O	2.01	0.60
1:A:146:THR:HB	3:A:810:HOH:O	2.00	0.60
1:A:120:GLY:HA3	1:A:202:PHE:CD2	2.36	0.59
1:A:181[A]:MET:CE	3:A:777:HOH:O	2.49	0.58
1:A:357[B]:PRO:O	1:A:358[B]:LEU:CD2	2.45	0.58
1:A:340:ASN:N	3:A:609:HOH:O	2.36	0.58
1:A:127[B]:GLN:NE2	3:A:610:HOH:O	2.37	0.57
1:A:135:ASP:HB2	2:A:503:GOL:H32	1.87	0.56
1:A:298:ALA:HB1	1:A:336:PHE:CZ	2.40	0.56
2:A:501:GOL:C3	3:A:660:HOH:O	2.54	0.55
1:A:165[B]:LYS:HE3	1:A:277:ASP:OD2	2.08	0.54
1:A:127[A]:GLN:NE2	3:A:610:HOH:O	2.41	0.54
1:A:312:LYS:HE2	3:A:887:HOH:O	2.07	0.53
1:A:135:ASP:CG	3:A:602:HOH:O	2.50	0.53
1:A:419:SER:HB2	1:A:420:PRO:HD3	1.90	0.52
1:A:217[A]:THR:HG22	1:A:218:SER:N	2.25	0.52
1:A:135:ASP:CB	2:A:503:GOL:H2	2.39	0.52
1:A:131:ALA:HA	2:A:503:GOL:O2	2.10	0.51
1:A:142[B]:MET:HE1	1:A:214:LEU:HD11	1.91	0.51
1:A:135:ASP:CG	2:A:503:GOL:C2	2.79	0.51
1:A:181[A]:MET:HE3	1:A:181[A]:MET:HA	1.92	0.50
1:A:183:LEU:HA	1:A:186[B]:VAL:HG22	1.93	0.50
1:A:313:PRO:O	1:A:315:TYR:N	2.45	0.49
1:A:297:ALA:HA	1:A:300[A]:MET:HE3	1.94	0.49
1:A:165[B]:LYS:HE2	1:A:212:MET:HG3	1.95	0.48
1:A:312:LYS:NZ	3:A:612:HOH:O	2.38	0.48
1:A:314:ASP:OD2	1:A:316:ASP:OD2	2.31	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:298:ALA:HA	1:A:347:VAL:HG11	1.95	0.47
1:A:338:TYR:O	3:A:603:HOH:O	2.20	0.47
1:A:165[B]:LYS:CE	1:A:277:ASP:OD2	2.62	0.47
1:A:411:LYS:NZ	3:A:624:HOH:O	2.48	0.47
1:A:169:ARG:NH1	1:A:205[B]:ASN:OD1	2.47	0.47
1:A:149:GLN:HB3	3:A:728:HOH:O	2.15	0.47
1:A:341:CYS:SG	1:A:350[A]:LYS:HD2	2.55	0.47
1:A:337:PRO:HB2	1:A:357[A]:PRO:CB	2.46	0.46
1:A:127[B]:GLN:HB2	1:A:129:TYR:CE2	2.51	0.46
1:A:196[B]:VAL:HG21	1:A:266:LEU:HD12	1.98	0.46
1:A:135:ASP:CB	2:A:503:GOL:H32	2.46	0.45
1:A:248:TYR:CE2	1:A:252:LYS:HG2	2.52	0.44
1:A:272[A]:GLN:NE2	3:A:626:HOH:O	2.50	0.44
1:A:120:GLY:HA3	1:A:202:PHE:CE2	2.52	0.44
1:A:258:ARG:HD2	3:A:736:HOH:O	2.18	0.44
1:A:132:GLU:HB2	2:A:503:GOL:H32	2.00	0.44
1:A:215[A]:MET:C	3:A:604:HOH:O	2.61	0.43
1:A:350[B]:LYS:HD2	1:A:354[B]:GLU:OE2	2.19	0.43
1:A:217[A]:THR:HG23	3:A:851:HOH:O	2.18	0.43
1:A:190:HIS:CD2	1:A:190:HIS:H	2.36	0.42
1:A:196[B]:VAL:HG13	1:A:213:GLU:HB3	2.01	0.42
1:A:276:CYS:SG	1:A:277:ASP:N	2.92	0.41
1:A:135:ASP:OD1	2:A:503:GOL:C1	2.67	0.41
1:A:215[A]:MET:O	3:A:604:HOH:O	2.22	0.41
1:A:360:PRO:HG2	1:A:363:MET:SD	2.60	0.41
1:A:337:PRO:O	2:A:504:GOL:C1	2.69	0.41
1:A:177:LYS:O	1:A:181[B]:MET:HG3	2.20	0.41
1:A:340:ASN:CA	3:A:609:HOH:O	2.69	0.41
1:A:362:HIS:CE1	1:A:363:MET:HG3	2.56	0.41
1:A:397:LYS:HA	1:A:397:LYS:HD3	1.92	0.40
1:A:410:PHE:CE1	1:A:414:MET:HG3	2.57	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:886:HOH:O	3:A:901:HOH:O[4_455]	1.97	0.23
3:A:619:HOH:O	3:A:658:HOH:O[1_565]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	321/323 (99%)	308 (96%)	8 (2%)	5 (2%)	7 1

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	146	THR
1	A	314	ASP
1	A	316	ASP
1	A	419	SER
1	A	315	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	288/284 (101%)	276 (96%)	12 (4%)	26 2

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130[A]	GLN
1	A	130[B]	GLN
1	A	135	ASP
1	A	217[A]	THR
1	A	217[B]	THR
1	A	252	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	272[A]	GLN
1	A	272[B]	GLN
1	A	314	ASP
1	A	363	MET
1	A	430	HIS
1	A	431	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	190	HIS
1	A	362	HIS
1	A	428	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	502	-	5,5,5	0.82	0	5,5,5	0.76	0
2	GOL	A	504	-	5,5,5	1.52	0	5,5,5	0.96	0
2	GOL	A	503	-	5,5,5	0.77	0	5,5,5	1.46	1 (20%)
2	GOL	A	501	-	5,5,5	0.59	0	5,5,5	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	502	-	-	2/4/4/4	-
2	GOL	A	504	-	-	2/4/4/4	-
2	GOL	A	503	-	-	3/4/4/4	-
2	GOL	A	501	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	503	GOL	O3-C3-C2	2.60	122.08	110.38

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	503	GOL	C1-C2-C3-O3
2	A	503	GOL	O2-C2-C3-O3
2	A	502	GOL	C1-C2-C3-O3
2	A	504	GOL	O1-C1-C2-C3
2	A	502	GOL	O2-C2-C3-O3
2	A	504	GOL	O1-C1-C2-O2
2	A	501	GOL	O2-C2-C3-O3
2	A	503	GOL	O1-C1-C2-O2
2	A	501	GOL	C1-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	504	GOL	1	0
2	A	503	GOL	9	0
2	A	501	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	304/323 (94%)	0.36	34 (11%)	10 11	9, 20, 75, 181	23 (7%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	315	TYR	6.0
1	A	420	PRO	5.8
1	A	294	ALA	5.0
1	A	135	ASP	4.7
1	A	147	CYS	4.6
1	A	433	PHE	4.3
1	A	146	THR	4.1
1	A	296	CYS	4.0
1	A	295	GLY	3.9
1	A	120	GLY	3.4
1	A	119	THR	3.4
1	A	311	THR	3.4
1	A	431	LEU	3.3
1	A	312	LYS	3.1
1	A	314	ASP	3.0
1	A	148	GLY	2.9
1	A	425	VAL	2.8
1	A	145	GLY	2.8
1	A	424	GLY	2.7
1	A	430	HIS	2.7
1	A	434	PHE	2.6
1	A	419	SER	2.5
1	A	415	ALA	2.5
1	A	313	PRO	2.5
1	A	133	ILE	2.5
1	A	216[A]	GLY	2.4
1	A	362	HIS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	144	SER	2.4
1	A	143	GLY	2.4
1	A	228	GLY	2.3
1	A	310	PRO	2.1
1	A	436	HIS	2.1
1	A	297	ALA	2.0
1	A	416	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	503	6/6	0.66	0.20	58,70,91,248	0
2	GOL	A	504	6/6	0.82	0.15	39,59,81,92	0
2	GOL	A	502	6/6	0.88	0.14	39,78,89,101	0
2	GOL	A	501	6/6	0.90	0.13	56,63,82,102	0

6.5 Other polymers [i](#)

There are no such residues in this entry.