



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 05:43 PM UTC

PDB ID : 1YAA / pdb_00001yaa
Title : ASPARTATE AMINOTRANSFERASE FROM SACCHAROMYCES CEREVISIAE CYTOPLASM
Authors : Jeffery, C.J.
Deposited on : 1998-01-27
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	412	 75% 22% .
1	B	412	 76% 21% .
1	C	412	 77% 21% .
1	D	412	 78% 20% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13596 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ASPARTATE AMINOTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	412	Total	C	N	O	S	0	0	0
			3197	2039	550	597	11			
1	B	412	Total	C	N	O	S	0	0	0
			3197	2039	550	597	11			
1	C	412	Total	C	N	O	S	0	0	0
			3197	2039	550	597	11			
1	D	412	Total	C	N	O	S	0	0	0
			3197	2039	550	597	11			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	95	LEU	PHE	conflict	UNP P23542
B	95	LEU	PHE	conflict	UNP P23542
C	95	LEU	PHE	conflict	UNP P23542
D	95	LEU	PHE	conflict	UNP P23542

- Molecule 2 is MALEIC ACID (CCD ID: MAE) (formula: C₄H₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			8	4	4		
2	B	1	Total	C	O	0	0
			8	4	4		
2	C	1	Total	C	O	0	0
			8	4	4		
2	D	1	Total	C	O	0	0
			8	4	4		

- Molecule 3 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
3	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
3	C	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
3	D	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	163	Total	O	0	0
			163	163		
4	B	183	Total	O	0	0
			183	183		
4	C	190	Total	O	0	0
			190	190		
4	D	180	Total	O	0	0
			180	180		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: ASPARTATE AMINOTRANSFERASE




• Molecule 1: ASPARTATE AMINOTRANSFERASE





• Molecule 1: ASPARTATE AMINOTRANSFERASE





● Molecule 1: ASPARTATE AMINOTRANSFERASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	130.31Å 134.63Å 98.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.05	Depositor
% Data completeness (in resolution range)	93.0 (10.00-2.05)	Depositor
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.231 , 0.299	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	13596	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAE, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.43	1/3271 (0.0%)	0.98	19/4439 (0.4%)
1	B	0.40	0/3271	1.25	20/4439 (0.5%)
1	C	0.41	0/3271	0.97	10/4439 (0.2%)
1	D	0.41	0/3271	0.99	17/4439 (0.4%)
All	All	0.42	1/13084 (0.0%)	1.05	66/17756 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	258	LYS	CD-CE	8.12	1.76	1.52

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	137	LYS	CA-C-N	32.43	160.38	119.84
1	B	137	LYS	C-N-CA	32.43	160.38	119.84
1	B	137	LYS	N-CA-C	-15.57	86.61	109.50
1	B	194	ASN	N-CA-C	-15.34	90.55	109.72
1	B	137	LYS	C-N-CD	-12.20	74.98	125.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3197	0	3182	64	0
1	B	3197	0	3182	59	0
1	C	3197	0	3182	54	0
1	D	3197	0	3181	49	0
2	A	8	0	2	0	0
2	B	8	0	2	0	0
2	C	8	0	2	0	0
2	D	8	0	2	0	0
3	A	15	0	6	2	0
3	B	15	0	6	1	0
3	C	15	0	6	1	0
3	D	15	0	6	1	0
4	A	163	0	0	2	0
4	B	183	0	0	3	0
4	C	190	0	0	2	0
4	D	180	0	0	4	0
All	All	13596	0	12759	217	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 217 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:LYS:CE	1:A:258:LYS:CD	1.76	1.61
1:A:280:LYS:HB3	1:A:281:PRO:HD3	1.69	0.74
1:B:20:ILE:HG22	1:B:23:ARG:HH22	1.51	0.73
1:B:393:ASN:O	1:B:397:VAL:HG23	1.89	0.72
1:A:244:LEU:HA	1:A:247:VAL:HG13	1.71	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	410/412 (100%)	393 (96%)	16 (4%)	1 (0%)	43	37
1	B	410/412 (100%)	393 (96%)	15 (4%)	2 (0%)	24	16
1	C	410/412 (100%)	393 (96%)	16 (4%)	1 (0%)	43	37
1	D	410/412 (100%)	403 (98%)	6 (2%)	1 (0%)	43	37
All	All	1640/1648 (100%)	1582 (96%)	53 (3%)	5 (0%)	36	30

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	263	TYR
1	B	138	PRO
1	C	263	TYR
1	D	263	TYR
1	B	263	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/342 (100%)	330 (96%)	12 (4%)	32	27
1	B	342/342 (100%)	326 (95%)	16 (5%)	23	17
1	C	342/342 (100%)	327 (96%)	15 (4%)	25	19
1	D	342/342 (100%)	327 (96%)	15 (4%)	25	19
All	All	1368/1368 (100%)	1310 (96%)	58 (4%)	26	20

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	17	LEU
1	D	341	LEU
1	C	153	LYS
1	D	329	ARG
1	D	137	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 46 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	335	HIS
1	D	82	ASN
1	C	340	HIS
1	C	377	HIS
1	D	113	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MAE	A	901	-	7,7,7	0.97	0	8,8,8	0.87	0
3	PLP	C	907	1	15,15,16	2.12	3 (20%)	21,22,23	1.69	4 (19%)
2	MAE	D	904	-	7,7,7	1.00	0	8,8,8	0.84	0
3	PLP	B	906	1	15,15,16	2.20	3 (20%)	21,22,23	2.13	6 (28%)
2	MAE	C	903	-	7,7,7	1.00	1 (14%)	8,8,8	0.81	0
3	PLP	D	908	1	15,15,16	1.86	4 (26%)	21,22,23	1.89	4 (19%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MAE	B	902	-	7,7,7	1.00	0	8,8,8	0.88	0
3	PLP	A	905	1	15,15,16	2.57	3 (20%)	21,22,23	1.94	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAE	A	901	-	-	2/5/5/5	-
3	PLP	C	907	1	-	2/6/6/8	0/1/1/1
2	MAE	D	904	-	-	2/5/5/5	-
3	PLP	B	906	1	-	2/6/6/8	0/1/1/1
2	MAE	C	903	-	-	2/5/5/5	-
3	PLP	D	908	1	-	2/6/6/8	0/1/1/1
2	MAE	B	902	-	-	2/5/5/5	-
3	PLP	A	905	1	-	2/6/6/8	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	905	PLP	C4A-C4	-8.57	1.34	1.51
3	B	906	PLP	C3-C2	-6.84	1.33	1.41
3	C	907	PLP	C3-C2	-6.16	1.34	1.41
3	D	908	PLP	C3-C2	-3.75	1.37	1.41
3	D	908	PLP	C4A-C4	2.96	1.57	1.51

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	906	PLP	O4P-C5A-C5	6.20	120.97	109.36
3	D	908	PLP	O4P-C5A-C5	6.04	120.68	109.36
3	A	905	PLP	O4P-C5A-C5	4.96	118.66	109.36
3	A	905	PLP	C6-C5-C4	4.77	122.01	118.10
3	C	907	PLP	O4P-C5A-C5	4.65	118.07	109.36

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	905	PLP	C4-C5-C5A-O4P
3	A	905	PLP	C6-C5-C5A-O4P
3	B	906	PLP	C4-C5-C5A-O4P
3	B	906	PLP	C6-C5-C5A-O4P
3	C	907	PLP	C4-C5-C5A-O4P

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	907	PLP	1	0
3	B	906	PLP	1	0
3	D	908	PLP	1	0
3	A	905	PLP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.