



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:26 AM UTC

PDB ID : 2YA9 / pdb_00002ya9
Title : Crystal structure of the autoinhibited form of mouse DAPK2
Authors : Patel, A.K.; Kursula, P.
Deposited on : 2011-02-18
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	
1	B	361	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	D1D	B	1302	-	X	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5357 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DEATH-ASSOCIATED PROTEIN KINASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	299	Total	C	N	O	S	0	6	0
			2481	1587	418	471	5			
1	B	299	Total	C	N	O	S	0	2	0
			2446	1568	411	462	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q8VDF3
B	0	GLY	-	expression tag	UNP Q8VDF3

- Molecule 2 is (4S,5S)-1,2-DITHIANE-4,5-DIOL (CCD ID: D1D) (formula: C₄H₈O₂S₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	S	0	0
			8	4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	O	S	0	0
			8	4	2	2		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	207	Total	O	0	0
			207	207		
4	B	206	Total	O	0	0
			206	206		

- Molecule 1: DEATH-ASSOCIATED PROTEIN KINASE 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.04Å 86.44Å 124.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.82 – 2.30 40.82 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.2 (40.82-2.30) 97.2 (40.82-2.30)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.60 (at 2.29Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.206 , 0.261 (Not available) , 0.222	Depositor DCC
R_{free} test set	1525 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	24.3	Xtriage
Anisotropy	0.502	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5357	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.11 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0405e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, D1D

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/2528	0.73	3/3411 (0.1%)
1	B	0.40	0/2492	0.73	0/3362
All	All	0.41	0/5020	0.73	3/6773 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	71	VAL	N-CA-C	5.32	115.52	107.75
1	A	154	ILE	CA-C-N	5.12	125.06	119.78
1	A	154	ILE	C-N-CA	5.12	125.06	119.78

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2481	0	2481	22	0
1	B	2446	0	2462	26	0
2	A	8	0	6	2	0
2	B	8	0	6	4	0
3	A	1	0	0	0	0
4	A	207	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	206	0	0	2	0
All	All	5357	0	4955	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1302:D1D:C3	2:A:1302:D1D:C4	1.80	1.57
2:B:1302:D1D:C3	2:B:1302:D1D:C4	1.80	1.52
1:A:47[A]:ARG:HG3	1:A:47[A]:ARG:HH11	1.10	1.11
1:A:47[B]:ARG:HG2	1:A:47[B]:ARG:HH11	1.14	1.08
1:B:26:ILE:HD11	1:B:28:LYS:HE3	1.62	0.82
1:A:47[A]:ARG:HG3	1:A:47[A]:ARG:NH1	1.89	0.81
1:A:47[B]:ARG:HG2	1:A:47[B]:ARG:NH1	1.94	0.73
1:A:47[B]:ARG:HH11	1:A:47[B]:ARG:CG	1.98	0.73
1:B:109:GLU:H	1:B:109:GLU:CD	1.95	0.72
1:A:208:TYR:CD1	1:A:216:PRO:HD3	2.29	0.68
1:A:47[A]:ARG:HD3	1:A:49:SER:O	1.94	0.67
1:A:261:ARG:HH11	1:A:261:ARG:HG2	1.63	0.63
2:B:1302:D1D:C3	2:B:1302:D1D:S4	2.87	0.59
1:B:169:GLU:HB2	1:B:172:VAL:HB	1.87	0.56
1:B:147:LEU:HD13	1:B:155:PRO:HB3	1.88	0.55
1:B:10:GLU:H	1:B:10:GLU:CD	2.15	0.54
1:A:259:GLU:CD	1:A:259:GLU:H	2.16	0.54
1:B:45:LYS:HE2	1:B:88:ASP:OD1	2.07	0.53
1:A:254:LYS:HB3	1:A:264:LEU:HG	1.91	0.52
1:B:73:HIS:CG	1:B:74:PRO:HD2	2.44	0.52
1:B:74:PRO:O	1:B:158:LYS:HE2	2.08	0.52
1:B:208:TYR:CD1	1:B:216:PRO:HD3	2.45	0.51
1:B:293:LEU:O	1:B:297:LYS:HG3	2.11	0.51
1:A:264:LEU:HD22	1:A:268:GLU:HB3	1.92	0.51
1:A:10:GLU:H	1:A:10:GLU:CD	2.19	0.50
1:A:47[B]:ARG:NH1	1:A:47[B]:ARG:CG	2.66	0.50
1:B:209:ILE:HG21	1:B:300:TYR:OH	2.12	0.49
1:B:253:ARG:HD2	4:B:2167:HOH:O	2.13	0.49
1:A:261:ARG:HG2	1:A:261:ARG:NH1	2.27	0.49
1:B:104:PHE:CE2	1:B:155:PRO:HG2	2.48	0.48
1:A:18:GLU:HG2	1:A:20:GLY:H	1.80	0.47
2:A:1302:D1D:C3	2:A:1302:D1D:S4	2.87	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:209:ILE:HD13	1:B:300:TYR:OH	2.15	0.47
1:B:53:ARG:HG3	4:B:2038:HOH:O	2.14	0.46
1:A:267[B]:GLN:HG3	4:A:2180:HOH:O	2.15	0.46
2:B:1302:D1D:C4	2:B:1302:D1D:H3	2.20	0.45
1:B:71:VAL:O	1:B:71:VAL:HG23	2.18	0.44
1:B:192:GLU:HB3	1:B:193:PRO:HD2	2.00	0.44
1:B:45:LYS:HB2	1:B:45:LYS:HE3	1.63	0.43
1:B:222:LYS:HD2	2:B:1302:D1D:H4C1	2.00	0.43
1:B:146:MET:HG3	1:B:160:ILE:HD13	2.01	0.42
1:A:146:MET:HG3	1:A:160:ILE:HD13	2.01	0.42
1:A:104:PHE:CE2	1:A:155:PRO:HG2	2.54	0.42
1:B:134:LYS:HE2	1:B:134:LYS:HB3	1.92	0.42
1:A:73:HIS:CG	1:A:74:PRO:HD2	2.54	0.42
1:B:120:ILE:O	1:B:124:LEU:HG	2.20	0.42
1:B:19:LEU:HD12	1:B:19:LEU:HA	1.78	0.41
1:B:210:LEU:O	1:B:210:LEU:HD23	2.20	0.41
1:A:47[A]:ARG:NH1	1:A:47[A]:ARG:CG	2.68	0.41
1:A:253:ARG:HD2	4:A:2165:HOH:O	2.21	0.41
1:B:210:LEU:HD23	1:B:210:LEU:C	2.46	0.41
1:B:9:VAL:HG13	1:B:10:GLU:N	2.36	0.40
1:A:178:PHE:CD1	1:A:178:PHE:C	3.00	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/361 (84%)	294 (97%)	9 (3%)	0	100	100
1	B	299/361 (83%)	293 (98%)	6 (2%)	0	100	100
All	All	602/722 (83%)	587 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/327 (69%)	219 (96%)	8 (4%)	32	48
1	B	270/327 (83%)	258 (96%)	12 (4%)	25	38
All	All	497/654 (76%)	477 (96%)	20 (4%)	28	42

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47[A]	ARG
1	A	47[B]	ARG
1	A	50	ARG
1	A	58	ARG
1	A	154	ILE
1	A	169	GLU
1	A	261	ARG
1	A	291	VAL
1	B	7	GLN
1	B	19	LEU
1	B	26	ILE
1	B	45	LYS
1	B	58	ARG
1	B	109	GLU
1	B	118	SER
1	B	170	ASP
1	B	280	THR
1	B	293	LEU
1	B	294	GLU
1	B	301	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	223	GLN
1	A	299	GLN
1	B	128	ASN
1	B	223	GLN
1	B	267	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	D1D	B	1302	-	8,8,8	9.80	8 (100%)	6,10,10	2.05	4 (66%)
2	D1D	A	1302	-	8,8,8	9.85	8 (100%)	6,10,10	1.84	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	D1D	B	1302	-	-	-	0/1/1/1
2	D1D	A	1302	-	-	-	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1302	D1D	C4-S4	-17.24	1.61	1.82
2	B	1302	D1D	C4-S4	-17.10	1.61	1.82
2	B	1302	D1D	C4-C3	15.07	1.80	1.52
2	A	1302	D1D	C4-C3	15.00	1.80	1.52
2	A	1302	D1D	C3-C2	11.36	1.69	1.52
2	B	1302	D1D	C3-C2	11.12	1.69	1.52
2	A	1302	D1D	C1-C2	6.74	1.65	1.52
2	B	1302	D1D	C1-C2	6.73	1.65	1.52
2	A	1302	D1D	C1-S1	-6.63	1.74	1.82
2	B	1302	D1D	C1-S1	-6.58	1.74	1.82
2	A	1302	D1D	O3-C3	-4.16	1.34	1.43
2	B	1302	D1D	O3-C3	-4.10	1.34	1.43
2	B	1302	D1D	O2-C2	-3.75	1.35	1.43
2	A	1302	D1D	O2-C2	-3.63	1.35	1.43
2	A	1302	D1D	S4-S1	-2.32	1.81	2.02
2	B	1302	D1D	S4-S1	-2.24	1.81	2.02

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1302	D1D	O3-C3-C4	2.79	114.49	109.81
2	A	1302	D1D	O2-C2-C1	2.49	113.98	109.81
2	B	1302	D1D	C1-C2-C3	2.46	117.32	112.45
2	B	1302	D1D	O2-C2-C1	2.35	113.75	109.81
2	A	1302	D1D	O3-C3-C4	2.32	113.70	109.81
2	B	1302	D1D	C4-C3-C2	2.25	116.89	112.45

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1302	D1D	4	0
2	A	1302	D1D	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	299/361 (82%)	-0.15	9 (3%)	52	54	7, 20, 50, 78	6 (2%)
1	B	299/361 (82%)	-0.19	5 (1%)	69	70	11, 22, 46, 71	2 (0%)
All	All	598/722 (82%)	-0.17	14 (2%)	61	63	7, 22, 47, 78	8 (1%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	300	TYR	4.9
1	A	22	GLY	4.7
1	B	301	VAL	4.1
1	B	300	TYR	3.7
1	A	20	GLY	3.4
1	A	3	THR	2.9
1	A	21	SER	2.4
1	A	301	VAL	2.3
1	A	23	GLN	2.3
1	B	86[A]	ARG	2.3
1	B	3	THR	2.2
1	A	107	GLN	2.1
1	B	23	GLN	2.1
1	A	24	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	A	1303	1/1	0.92	0.10	63,63,63,63	0
2	D1D	B	1302	8/8	0.93	0.11	21,38,43,43	0
2	D1D	A	1302	8/8	0.96	0.08	12,35,40,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.