



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2026 – 02:20 AM UTC

PDB ID : 4YAQ / pdb_00004yaq
Title : Crystal structure of a computationally optimized PG9 mutant
Authors : Murrell, S.; Julien, J.P.; Wilson, I.A.
Deposited on : 2015-02-17
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	
1	H	248	
2	B	216	
2	L	216	
3	C	3	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	3	 33% 67%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7215 atoms, of which 52 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PG9_N100FY Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	239	Total	C	N	O	S	0	0	0
			1843	1160	308	365	10			
1	A	239	Total	C	N	O	S	0	0	0
			1843	1160	308	365	10			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
H	233	GLY	-	expression tag	UNP A0A087X1C7
H	234	LEU	-	expression tag	UNP A0A087X1C7
H	235	GLU	-	expression tag	UNP A0A087X1C7
H	236	VAL	-	expression tag	UNP A0A087X1C7
H	237	LEU	-	expression tag	UNP A0A087X1C7
H	238	PHE	-	expression tag	UNP A0A087X1C7
H	239	GLN	-	expression tag	UNP A0A087X1C7
A	233	GLY	-	expression tag	UNP A0A087X1C7
A	234	LEU	-	expression tag	UNP A0A087X1C7
A	235	GLU	-	expression tag	UNP A0A087X1C7
A	236	VAL	-	expression tag	UNP A0A087X1C7
A	237	LEU	-	expression tag	UNP A0A087X1C7
A	238	PHE	-	expression tag	UNP A0A087X1C7
A	239	GLN	-	expression tag	UNP A0A087X1C7

- Molecule 2 is a protein called PG9_N100FY Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	214	Total	C	N	O	S	0	0	0
			1587	985	270	327	5			
2	B	214	Total	C	N	O	S	0	0	0
			1587	985	270	327	5			


- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[al

pha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	3	Total	C	N	O	0	0	0
			38	22	2	14			
3	D	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	H	1	Total	C	H	N	O S	0	0
			25	6	13	1	4 1		
4	L	1	Total	C	H	N	O S	0	0
			25	6	13	1	4 1		
4	L	1	Total	C	H	N	O S	0	0
			25	6	13	1	4 1		
4	B	1	Total	C	H	N	O S	0	0
			25	6	13	1	4 1		


- Molecule 5 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	H	1	Total	O	P	0	0
			5	4	1		
5	L	1	Total	O	P	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	54	Total	O	0	0
			54	54		
6	L	38	Total	O	0	0
			38	38		
6	A	41	Total	O	0	0
			41	41		
6	B	36	Total	O	0	0
			36	36		

Chain C:  33% 67%

MAG1
MAG2
FUC3

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  33% 67%

MAG1
MAG2
FUC3

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.74Å 122.98Å 69.92Å 90.00° 96.27° 90.00°	Depositor
Resolution (Å)	37.26 – 2.30 37.26 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.3 (37.26-2.30) 98.3 (37.26-2.30)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.204 , 0.238 0.209 , 0.243	Depositor DCC
R_{free} test set	2000 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	45.2	Xtriage
Anisotropy	0.282	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7215	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TYS, NAG, PCA, MES, PO4, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/1852	0.66	0/2519
1	H	0.23	0/1852	0.66	0/2519
2	B	0.24	0/1622	0.68	0/2209
2	L	0.25	0/1622	0.68	0/2209
All	All	0.24	0/6948	0.67	0/9456

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1843	0	1741	15	0
1	H	1843	0	1741	19	0
2	B	1587	0	1547	14	0
2	L	1587	0	1547	24	0
3	C	38	0	34	1	0
3	D	38	0	34	1	0
4	B	12	13	13	0	0
4	H	12	13	13	0	0
4	L	24	26	26	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	H	5	0	0	0	0
5	L	5	0	0	0	0
6	A	41	0	0	0	0
6	B	36	0	0	0	0
6	H	54	0	0	1	0
6	L	38	0	0	1	0
All	All	7163	52	6696	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:SER:HB3	1:A:228:LYS:HE3	1.62	0.80
2:L:120:PRO:HD3	2:L:132:LEU:CD2	2.26	0.65
2:L:115:VAL:O	2:L:204:LYS:NZ	2.30	0.65
1:A:190:VAL:HG21	2:B:135:LEU:HD13	1.80	0.63
2:L:110:LYS:HD3	2:L:198:GLU:CB	2.29	0.62
1:H:208:CYS:SG	1:H:221:LYS:HB3	2.41	0.61
2:B:12:SER:HB2	2:B:106(A):LEU:HD23	1.85	0.57
2:L:42:LYS:NZ	6:L:403:HOH:O	2.38	0.56
2:L:110:LYS:HD3	2:L:198:GLU:HB3	1.87	0.56
2:B:120:PRO:HD3	2:B:132:LEU:CD2	2.37	0.55
2:L:95(A):ARG:NH2	2:B:170:ASN:OD1	2.42	0.53
1:A:208:CYS:SG	1:A:221:LYS:HB3	2.48	0.53
1:A:127:SER:HB3	1:A:228:LYS:CE	2.37	0.52
1:A:152:VAL:CG2	1:A:187:LEU:HD21	2.40	0.52
1:A:190:VAL:HG21	2:B:135:LEU:CD1	2.41	0.51
1:H:190:VAL:HG21	2:L:135:LEU:CD1	2.40	0.51
2:L:120:PRO:HD3	2:L:132:LEU:HD23	1.93	0.51
3:C:2:NAG:H82	3:C:3:FUC:C1	2.41	0.51
1:H:127:SER:HB2	1:H:230:CYS:HB2	1.93	0.50
2:L:149:LYS:HG3	2:L:194:GLN:NE2	2.25	0.50
2:L:136:ILE:N	2:L:136:ILE:HD12	2.26	0.50
1:H:152:VAL:CG2	1:H:187:LEU:HD21	2.41	0.50
3:D:2:NAG:H82	3:D:3:FUC:C1	2.42	0.50
1:H:177:VAL:CG1	2:L:162:THR:HG23	2.42	0.50
1:A:166:LEU:HD21	1:A:191:VAL:HG21	1.94	0.50
2:B:12:SER:HB2	2:B:106(A):LEU:CD2	2.42	0.50
1:A:187:LEU:C	1:A:187:LEU:HD12	2.38	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:187:LEU:C	1:H:187:LEU:HD12	2.38	0.49
2:B:15:PRO:HD3	2:B:106(A):LEU:O	2.14	0.48
2:L:39:HIS:HB3	2:L:40:PRO:HD2	1.95	0.48
2:L:106:VAL:O	2:L:108:GLN:NE2	2.46	0.47
2:B:14:SER:O	2:B:17:GLN:HG2	2.14	0.47
1:A:22:CYS:HB3	1:A:78:LEU:HB3	1.95	0.47
2:B:111:ALA:O	2:B:139:PHE:HA	2.15	0.47
1:A:127:SER:O	1:A:133:THR:HG23	2.15	0.47
1:H:100(B):ARG:HB2	1:H:100(G):TYS:HE1	1.97	0.46
1:H:127:SER:O	1:H:133:THR:HG23	2.15	0.46
1:H:22:CYS:HB3	1:H:78:LEU:HB3	1.97	0.46
2:L:83:GLU:HG3	2:L:106:VAL:HG23	1.98	0.46
2:L:149:LYS:HD3	2:L:154:PRO:HA	1.97	0.46
1:A:147:TYR:CE1	1:A:152:VAL:HG13	2.51	0.45
2:B:140:TYR:HA	2:B:141:PRO:C	2.40	0.45
1:H:32:GLN:NE2	6:H:406:HOH:O	2.50	0.45
2:L:35:TRP:CD2	2:L:73:LEU:HB2	2.52	0.45
2:L:140:TYR:HA	2:L:141:PRO:C	2.42	0.44
2:L:153:SER:HA	2:L:154:PRO:HD3	1.88	0.44
1:H:100(B):ARG:HB2	1:H:100(G):TYS:CE1	2.48	0.43
2:L:132:LEU:HD12	2:L:178:LEU:HD23	1.99	0.43
2:B:194:GLN:NE2	2:B:203:GLU:OE1	2.43	0.43
2:B:183:GLU:OE1	2:B:183:GLU:N	2.51	0.43
1:H:193:VAL:HG11	1:H:206:TYR:CE1	2.53	0.43
1:H:190:VAL:HG21	2:L:135:LEU:HD13	2.00	0.42
2:B:35:TRP:CD2	2:B:73:LEU:HB2	2.55	0.42
1:H:213:LYS:N	1:H:214:PRO:CD	2.82	0.42
1:A:100(A):TYR:HD2	1:A:100(F):TYR:CE1	2.38	0.42
1:H:93:VAL:CG1	1:H:100(T):MET:HB3	2.50	0.42
1:H:58:TYR:OH	2:L:95:ARG:HD3	2.20	0.41
1:H:152:VAL:HG23	1:H:187:LEU:HD21	2.03	0.41
1:H:166:LEU:HD21	1:H:191:VAL:HG21	2.02	0.41
1:A:213:LYS:N	1:A:214:PRO:CD	2.83	0.41
2:B:4:LEU:HB2	2:B:99:GLY:HA2	2.03	0.41
1:H:52:LYS:O	1:H:71:ARG:NH1	2.53	0.41
2:L:14:SER:O	2:L:17:GLN:HG2	2.21	0.41
2:L:150:ALA:HB2	2:L:155:VAL:HG21	2.01	0.41
2:L:91:LEU:HD21	1:A:167:THR:HB	2.03	0.41
1:A:100(B):ARG:O	1:A:100(G):TYS:HE1	2.19	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/248 (95%)	230 (98%)	5 (2%)	0	100	100
1	H	235/248 (95%)	231 (98%)	4 (2%)	0	100	100
2	B	212/216 (98%)	203 (96%)	9 (4%)	0	100	100
2	L	212/216 (98%)	206 (97%)	6 (3%)	0	100	100
All	All	894/928 (96%)	870 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/207 (96%)	199 (100%)	0	100	100
1	H	199/207 (96%)	199 (100%)	0	100	100
2	B	181/183 (99%)	181 (100%)	0	100	100
2	L	181/183 (99%)	181 (100%)	0	100	100
All	All	760/780 (97%)	760 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	81	GLN
1	H	82(A)	ASN
1	H	172	HIS
2	L	17	GLN
2	L	167	GLN
2	L	197	HIS
1	A	100(R)	HIS
2	B	17	GLN
2	B	39	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	TYS	H	100(H)	1	15,16,17	1.27	3 (20%)	15,22,24	0.59	0
1	PCA	H	2	1	7,8,9	1.92	1 (14%)	9,10,12	2.13	5 (55%)
1	PCA	A	2	1	7,8,9	1.92	1 (14%)	9,10,12	2.16	5 (55%)
1	TYS	H	100(G)	1	15,16,17	1.27	3 (20%)	15,22,24	0.63	0
1	TYS	A	100(H)	1	15,16,17	1.28	3 (20%)	15,22,24	0.63	0
1	TYS	A	100(G)	1	15,16,17	1.28	3 (20%)	15,22,24	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TYS	H	100(H)	1	-	0/10/11/13	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	H	2	1	-	0/0/11/13	0/1/1/1
1	PCA	A	2	1	-	0/0/11/13	0/1/1/1
1	TYS	H	100(G)	1	-	3/10/11/13	0/1/1/1
1	TYS	A	100(H)	1	-	3/10/11/13	0/1/1/1
1	TYS	A	100(G)	1	-	0/10/11/13	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2	PCA	CD-N	4.98	1.46	1.34
1	H	2	PCA	CD-N	4.95	1.46	1.34
1	A	100(G)	TYS	OH-CZ	-3.29	1.37	1.42
1	A	100(H)	TYS	OH-CZ	-3.20	1.37	1.42
1	H	100(G)	TYS	OH-CZ	-3.20	1.37	1.42
1	H	100(H)	TYS	OH-CZ	-3.17	1.37	1.42
1	H	100(G)	TYS	OH-S	-2.31	1.54	1.58
1	A	100(G)	TYS	OH-S	-2.30	1.54	1.58
1	A	100(H)	TYS	OH-S	-2.28	1.54	1.58
1	H	100(H)	TYS	OH-S	-2.27	1.54	1.58
1	A	100(H)	TYS	O3-S	2.17	1.64	1.50
1	H	100(G)	TYS	O3-S	2.17	1.64	1.50
1	H	100(H)	TYS	O3-S	2.17	1.64	1.50
1	A	100(G)	TYS	O3-S	2.16	1.64	1.50

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	2	PCA	OE-CD-CG	-3.02	121.33	126.72
1	A	2	PCA	OE-CD-CG	-2.97	121.42	126.72
1	H	2	PCA	CA-N-CD	-2.84	103.85	113.58
1	H	2	PCA	CB-CA-N	2.83	111.04	103.24
1	A	2	PCA	CA-N-CD	-2.83	103.88	113.58
1	A	2	PCA	CB-CA-N	2.78	110.89	103.24
1	H	2	PCA	CG-CD-N	2.59	114.74	108.39
1	A	2	PCA	CG-CD-N	2.55	114.63	108.39
1	A	2	PCA	CB-CA-C	-2.52	109.20	112.66
1	H	2	PCA	CB-CA-C	-2.23	109.59	112.66

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	H	100(G)	TYS	CZ-OH-S-O1
1	H	100(G)	TYS	CZ-OH-S-O3
1	A	100(H)	TYS	CZ-OH-S-O3
1	H	100(G)	TYS	CZ-OH-S-O2
1	A	100(H)	TYS	CZ-OH-S-O1
1	A	100(H)	TYS	CZ-OH-S-O2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	H	100(G)	TYS	2	0
1	A	100(G)	TYS	1	0

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	3,2	14,14,15	0.48	0	17,19,21	0.77	0
3	NAG	C	2	3	14,14,15	0.75	0	17,19,21	0.50	0
3	FUC	C	3	3	10,10,11	0.80	0	14,14,16	0.58	0
3	NAG	D	1	3,2	14,14,15	0.24	0	17,19,21	0.72	0
3	NAG	D	2	3	14,14,15	0.83	1 (7%)	17,19,21	0.52	0
3	FUC	D	3	3	10,10,11	0.84	1 (10%)	14,14,16	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,2	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
3	FUC	C	3	3	-	-	0/1/1/1
3	NAG	D	1	3,2	-	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
3	FUC	D	3	3	-	-	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2	NAG	O5-C1	-2.73	1.39	1.43
3	D	3	FUC	O5-C1	-2.07	1.40	1.43

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	2	NAG	O5-C5-C6-O6
3	D	2	NAG	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2
3	D	1	NAG	C8-C7-N2-C2
3	D	1	NAG	O7-C7-N2-C2


There are no ring outliers.

4 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	2	NAG	1	0
3	C	2	NAG	1	0
3	C	3	FUC	1	0
3	D	3	FUC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MES	H	301	-	12,12,12	2.32	1 (8%)	15,16,16	1.35	1 (6%)
5	PO4	L	306	-	4,4,4	0.97	0	6,6,6	0.46	0
4	MES	L	305	-	12,12,12	2.29	1 (8%)	15,16,16	1.36	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MES	L	304	-	12,12,12	2.32	1 (8%)	15,16,16	1.34	1 (6%)
4	MES	B	304	-	12,12,12	2.36	1 (8%)	15,16,16	1.34	2 (13%)
5	PO4	H	302	-	4,4,4	0.96	0	6,6,6	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MES	B	304	-	-	0/6/14/14	0/1/1/1
4	MES	H	301	-	-	1/6/14/14	0/1/1/1
4	MES	L	305	-	-	1/6/14/14	0/1/1/1
4	MES	L	304	-	-	1/6/14/14	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	304	MES	C8-S	-7.92	1.66	1.77
4	L	304	MES	C8-S	-7.78	1.66	1.77
4	H	301	MES	C8-S	-7.76	1.66	1.77
4	L	305	MES	C8-S	-7.68	1.66	1.77

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	L	305	MES	C5-N4-C3	2.82	114.91	108.84
4	L	304	MES	C5-N4-C3	2.42	114.06	108.84
4	H	301	MES	C5-N4-C3	2.29	113.78	108.84
4	B	304	MES	C6-C5-N4	-2.27	106.67	110.12
4	B	304	MES	C5-N4-C3	2.24	113.67	108.84

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	305	MES	C8-C7-N4-C3
4	H	301	MES	C8-C7-N4-C3
4	L	304	MES	C8-C7-N4-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	236/248 (95%)	0.24	5 (2%) 63 65	30, 50, 77, 134	0
1	H	236/248 (95%)	0.03	5 (2%) 63 65	32, 44, 71, 104	0
2	B	214/216 (99%)	0.33	9 (4%) 40 42	32, 50, 78, 89	0
2	L	214/216 (99%)	0.21	4 (1%) 66 68	31, 48, 77, 101	0
All	All	900/928 (96%)	0.20	23 (2%) 57 59	30, 48, 77, 134	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	211	CYS	4.2
2	B	199	GLY	4.1
2	B	106(A)	LEU	3.2
2	B	28	GLY	3.0
2	B	2	SER	3.0
2	L	199	GLY	2.5
1	A	100(J)	PHE	2.4
1	A	129	LYS	2.4
1	A	100(K)	TYR	2.4
1	H	115	SER	2.3
1	H	100(A)	TYR	2.3
1	H	229	SER	2.3
2	B	200	SER	2.3
1	H	100(F)	TYR	2.3
2	L	2	SER	2.3
2	L	211	CYS	2.2
2	B	168	SER	2.2
2	L	109	PRO	2.2
1	A	100(E)	TYR	2.1
2	B	9	SER	2.1
1	A	100(A)	TYR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	100(D)	GLY	2.1
2	B	110	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TYS	A	100(G)	16/17	0.72	0.14	73,108,125,126	0
1	TYS	H	100(G)	16/17	0.85	0.12	53,85,108,109	0
1	PCA	H	2	8/9	0.86	0.12	49,55,60,67	0
1	PCA	A	2	8/9	0.87	0.11	65,69,70,71	0
1	TYS	A	100(H)	16/17	0.90	0.11	62,68,78,89	0
1	TYS	H	100(H)	16/17	0.92	0.11	43,50,63,77	0

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	C	1	14/15	-	-	79,82,87,88	0
3	NAG	C	2	14/15	-	-	76,87,92,92	0
3	FUC	C	3	10/11	-	-	84,97,100,105	0
3	NAG	D	1	14/15	-	-	80,86,91,94	0
3	NAG	D	2	14/15	-	-	90,94,99,101	0
3	FUC	D	3	10/11	-	-	95,102,108,111	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	PO4	L	306	5/5	0.66	0.14	112,113,114,119	0
4	MES	B	304	12/12	0.77	0.18	90,110,122,124	0
4	MES	L	304	12/12	0.84	0.14	81,100,104,106	0
4	MES	L	305	12/12	0.84	0.11	52,68,83,85	0
4	MES	H	301	12/12	0.92	0.07	49,60,66,69	0
5	PO4	H	302	5/5	0.94	0.10	55,58,62,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.