



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 11:45 PM UTC

PDB ID : 6YAJ / pdb_00006yaj
Title : Split gene transketolase, inactive beta4 tetramer
Authors : Isupov, M.N.; Littlechild, J.A.; James, P.
Deposited on : 2020-03-12
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	341	4% 77% 11% • 11%
1	BBB	341	5% 77% 10% • 12%
1	CCC	341	5% 79% 8% • 12%
1	DDD	341	4% 80% 11% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	AAA	403	-	-	X	-
3	GOL	DDD	401	-	-	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9891 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C-terminal chain of split transketolase from Carboxydotherrmus hydrogenoformans.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	303	Total	C	N	O	S	0	23	0
			2390	1533	410	438	9			
1	BBB	299	Total	C	N	O	S	0	17	0
			2305	1479	391	426	9			
1	CCC	300	Total	C	N	O	S	0	22	0
			2344	1507	393	436	8			
1	DDD	311	Total	C	N	O	S	0	20	0
			2415	1548	412	447	8			

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	AAA	1	Total	C	O	0	0
			4	2	2		
2	AAA	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	AAA	1	Total	C	O	0	0
			4	2	2		
2	AAA	1	Total	C	O	0	0
			4	2	2		
2	BBB	1	Total	C	O	0	0
			4	2	2		
2	BBB	1	Total	C	O	0	0
			4	2	2		
2	BBB	1	Total	C	O	0	0
			4	2	2		
2	CCC	1	Total	C	O	0	0
			4	2	2		
2	CCC	1	Total	C	O	0	0
			4	2	2		
2	CCC	1	Total	C	O	0	0
			4	2	2		
2	DDD	1	Total	C	O	0	0
			4	2	2		
2	DDD	1	Total	C	O	0	0
			4	2	2		
2	DDD	1	Total	C	O	0	0
			4	2	2		
2	DDD	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	C	O	0	0
			6	3	3		
3	BBB	1	Total	C	O	0	0
			6	3	3		
3	BBB	1	Total	C	O	0	0
			6	3	3		
3	CCC	1	Total	C	O	0	0
			6	3	3		
3	DDD	1	Total	C	O	0	0
			6	3	3		
3	DDD	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	BBB	1	Total	C	O	0	0
			7	4	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	AAA	91	Total	O	0	0
			91	91		
5	BBB	81	Total	O	0	0
			81	81		
5	CCC	66	Total	O	0	0
			66	66		
5	DDD	96	Total	O	0	0
			96	96		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: C-terminal chain of split transketolase from *Carboxydothermus hydrogenofor*mans




- Molecule 1: C-terminal chain of split transketolase from *Carboxydothermus hydrogenofor*mans



- Molecule 1: C-terminal chain of split transketolase from *Carboxydothermus hydrogenofor*mans



- Molecule 1: C-terminal chain of split transketolase from Carboxydotherrmus hydrogenofor-mans



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	101.88Å 101.88Å 164.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.01 – 1.90 64.01 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (64.01-1.90) 99.8 (64.01-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.178 , 0.208 0.177 , 0.208	Depositor DCC
R_{free} test set	6613 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	44.9	Xtriage
Anisotropy	0.424	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.042 for h,-k,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	9891	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, EDO, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	0.65	1/2485 (0.0%)	1.00	0/3352
1	BBB	0.63	0/2387	1.02	4/3225 (0.1%)
1	CCC	0.64	0/2441	0.99	3/3295 (0.1%)
1	DDD	0.64	0/2509	1.01	1/3391 (0.0%)
All	All	0.64	1/9822 (0.0%)	1.00	8/13263 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	213	GLU	CD-OE1	5.43	1.35	1.25

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	7	ARG	CB-CG-CD	7.20	127.86	111.30
1	DDD	84	ARG	CG-CD-NE	5.29	123.64	112.00
1	BBB	7	ARG	CG-CD-NE	5.29	123.64	112.00
1	CCC	145	ARG	CB-CG-CD	5.28	123.44	111.30
1	BBB	231[A]	GLU	CB-CG-CD	5.25	121.52	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2390	0	2549	24	0
1	BBB	2305	0	2449	25	0
1	CCC	2344	0	2492	22	0
1	DDD	2415	0	2569	25	0
2	AAA	16	0	24	1	0
2	BBB	12	0	18	1	0
2	CCC	12	0	18	2	0
2	DDD	20	0	30	1	0
3	AAA	6	0	8	4	0
3	BBB	12	0	16	2	0
3	CCC	6	0	8	3	0
3	DDD	12	0	16	4	0
4	BBB	7	0	10	0	0
5	AAA	91	0	0	3	0
5	BBB	81	0	0	1	0
5	CCC	66	0	0	0	0
5	DDD	96	0	0	5	0
All	All	9891	0	10207	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 92 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:296:THR:HB	3:AAA:403:GOL:H12	1.03	1.02
1:DDD:46:GLU:HB2	3:DDD:401:GOL:H12	1.39	1.01
1:AAA:296:THR:CB	3:AAA:403:GOL:H12	1.99	0.86
1:CCC:280:PHE:O	1:CCC:282:GLU:HG3	1.76	0.86
2:DDD:407:EDO:H22	5:DDD:592:HOH:O	1.75	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	320/341 (94%)	309 (97%)	10 (3%)	1 (0%)	36	29
1	BBB	310/341 (91%)	301 (97%)	9 (3%)	0	100	100
1	CCC	316/341 (93%)	308 (98%)	8 (2%)	0	100	100
1	DDD	329/341 (96%)	313 (95%)	14 (4%)	2 (1%)	21	13
All	All	1275/1364 (94%)	1231 (96%)	41 (3%)	3 (0%)	36	36

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	279	VAL
1	DDD	283	SER
1	DDD	112	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	256/260 (98%)	246 (96%)	10 (4%)	28	21
1	BBB	246/260 (95%)	238 (97%)	8 (3%)	33	26
1	CCC	251/260 (96%)	242 (96%)	9 (4%)	31	23
1	DDD	258/260 (99%)	245 (95%)	13 (5%)	22	14
All	All	1011/1040 (97%)	971 (96%)	40 (4%)	30	20

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	DDD	111	VAL
1	DDD	164	ARG
1	DDD	113	GLU
1	DDD	119[A]	GLN
1	DDD	196	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	DDD	404	-	3,3,3	0.24	0	2,2,2	0.67	0
2	EDO	BBB	403	-	3,3,3	0.16	0	2,2,2	0.60	0
2	EDO	AAA	405	-	3,3,3	0.18	0	2,2,2	0.54	0
2	EDO	DDD	405	-	3,3,3	0.10	0	2,2,2	0.27	0
2	EDO	BBB	402	-	3,3,3	0.10	0	2,2,2	0.16	0
2	EDO	CCC	501	-	3,3,3	0.19	0	2,2,2	0.53	0
3	GOL	DDD	401	-	5,5,5	0.13	0	5,5,5	0.39	0
3	GOL	DDD	406	-	5,5,5	0.12	0	5,5,5	0.50	0
3	GOL	BBB	401	-	5,5,5	0.12	0	5,5,5	0.34	0
3	GOL	CCC	504	-	5,5,5	0.13	0	5,5,5	0.65	0
2	EDO	AAA	401	-	3,3,3	0.08	0	2,2,2	0.44	0
2	EDO	AAA	402	-	3,3,3	0.17	0	2,2,2	0.71	0
2	EDO	CCC	503	-	3,3,3	0.19	0	2,2,2	0.46	0
2	EDO	DDD	403	-	3,3,3	0.17	0	2,2,2	0.33	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	DDD	402	-	3,3,3	0.05	0	2,2,2	0.16	0
4	PEG	BBB	406	-	6,6,6	0.39	0	5,5,5	0.21	0
3	GOL	BBB	405	-	5,5,5	0.10	0	5,5,5	0.32	0
3	GOL	AAA	403	-	5,5,5	0.07	0	5,5,5	0.31	0
2	EDO	AAA	404	-	3,3,3	0.25	0	2,2,2	0.23	0
2	EDO	BBB	404	-	3,3,3	0.22	0	2,2,2	0.53	0
2	EDO	DDD	407	-	3,3,3	0.12	0	2,2,2	0.51	0
2	EDO	CCC	502	-	3,3,3	0.15	0	2,2,2	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	DDD	404	-	-	1/1/1/1	-
2	EDO	BBB	403	-	-	1/1/1/1	-
2	EDO	AAA	405	-	-	1/1/1/1	-
2	EDO	DDD	405	-	-	1/1/1/1	-
2	EDO	BBB	402	-	-	1/1/1/1	-
2	EDO	CCC	501	-	-	0/1/1/1	-
3	GOL	DDD	401	-	-	2/4/4/4	-
3	GOL	DDD	406	-	-	4/4/4/4	-
3	GOL	BBB	401	-	-	2/4/4/4	-
3	GOL	CCC	504	-	-	2/4/4/4	-
2	EDO	AAA	401	-	-	1/1/1/1	-
2	EDO	AAA	402	-	-	0/1/1/1	-
2	EDO	CCC	503	-	-	1/1/1/1	-
2	EDO	DDD	403	-	-	1/1/1/1	-
2	EDO	DDD	402	-	-	1/1/1/1	-
4	PEG	BBB	406	-	-	3/4/4/4	-
3	GOL	BBB	405	-	-	2/4/4/4	-
3	GOL	AAA	403	-	-	2/4/4/4	-
2	EDO	AAA	404	-	-	1/1/1/1	-
2	EDO	BBB	404	-	-	0/1/1/1	-
2	EDO	DDD	407	-	-	0/1/1/1	-
2	EDO	CCC	502	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 27 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	BBB	401	GOL	C1-C2-C3-O3
3	BBB	405	GOL	O1-C1-C2-O2
3	BBB	405	GOL	O1-C1-C2-C3
3	CCC	504	GOL	O1-C1-C2-C3
3	DDD	406	GOL	O1-C1-C2-O2

There are no ring outliers.

9 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	BBB	403	EDO	1	0
3	DDD	401	GOL	4	0
3	CCC	504	GOL	3	0
2	CCC	503	EDO	1	0
3	BBB	405	GOL	2	0
3	AAA	403	GOL	4	0
2	AAA	404	EDO	1	0
2	DDD	407	EDO	1	0
2	CCC	502	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	303/341 (88%)	0.15	14 (4%) 37 40	22, 49, 110, 178	23 (7%)
1	BBB	299/341 (87%)	0.15	16 (5%) 31 33	28, 51, 101, 175	17 (5%)
1	CCC	300/341 (87%)	0.28	18 (6%) 27 29	26, 53, 99, 179	22 (7%)
1	DDD	311/341 (91%)	0.02	13 (4%) 40 43	26, 49, 107, 165	20 (6%)
All	All	1213/1364 (88%)	0.15	61 (5%) 34 36	22, 51, 107, 179	82 (6%)

The worst 5 of 61 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	289	LEU	5.6
1	CCC	279	VAL	5.5
1	BBB	118[A]	HIS	4.9
1	BBB	279	VAL	4.9
1	DDD	111	VAL	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	AAA	405	4/4	0.57	0.12	89,96,105,114	0
2	EDO	CCC	503	4/4	0.63	0.15	85,98,102,104	0
2	EDO	BBB	403	4/4	0.74	0.17	93,97,98,103	0
2	EDO	AAA	404	4/4	0.74	0.15	90,93,101,102	0
2	EDO	DDD	404	4/4	0.79	0.16	75,79,81,92	0
3	GOL	BBB	401	6/6	0.79	0.12	79,106,109,109	0
2	EDO	CCC	502	4/4	0.80	0.17	76,78,85,89	0
2	EDO	BBB	404	4/4	0.81	0.16	79,84,87,94	0
3	GOL	AAA	403	6/6	0.82	0.15	98,100,103,105	0
3	GOL	CCC	504	6/6	0.83	0.14	80,90,97,121	0
2	EDO	AAA	402	4/4	0.84	0.15	77,81,81,85	0
3	GOL	DDD	406	6/6	0.84	0.14	58,76,94,98	0
3	GOL	DDD	401	6/6	0.85	0.12	60,84,89,93	0
2	EDO	DDD	402	4/4	0.85	0.17	100,112,122,122	0
2	EDO	DDD	405	4/4	0.86	0.13	94,95,100,103	0
3	GOL	BBB	405	6/6	0.89	0.13	64,75,82,83	6
4	PEG	BBB	406	7/7	0.89	0.18	77,85,99,108	0
2	EDO	DDD	403	4/4	0.90	0.14	92,95,95,99	0
2	EDO	AAA	401	4/4	0.90	0.20	65,82,92,93	0
2	EDO	CCC	501	4/4	0.92	0.14	74,88,89,97	0
2	EDO	BBB	402	4/4	0.92	0.20	80,97,104,108	0
2	EDO	DDD	407	4/4	0.95	0.16	70,86,93,96	0

6.5 Other polymers ⓘ

There are no such residues in this entry.