



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 06:03 PM UTC

PDB ID : 2YBR / pdb_00002ybr
Title : Crystal structure of the human derived single chain antibody fragment (scFv) 9004G in complex with Cn2 toxin from the scorpion *Centruroides noxius* Hoffmann
Authors : Canul-Tec, J.C.; Riano-Umbarila, L.; Rudino-Pinera, E.; Becerril, B.; Possani, L.D.; Torres-Larios, A.
Deposited on : 2011-03-09
Resolution : 2.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	117	2% 80% 17% ..
1	D	117	% 74% 22% ..
1	G	117	2% 77% 19% ..
2	B	146	68% 8% 23%
2	E	146	% 58% 13% 27%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	146	70%8%22%
3	C	66	91%8%
3	F	66	2% 83%15%
3	I	66	2% 83%14%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6941 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SINGLE CHAIN ANTIBODY FRAGMENT 9004G.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	116	Total	C	N	O	S	0	0	0
			869	538	152	174	5			
1	D	115	Total	C	N	O	S	0	0	0
			866	537	151	173	5			
1	G	115	Total	C	N	O	S	0	0	0
			866	537	151	173	5			

- Molecule 2 is a protein called SINGLE CHAIN ANTIBODY FRAGMENT 9004G.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	112	Total	C	N	O	S	0	0	0
			851	532	149	168	2			
2	E	107	Total	C	N	O	S	0	0	0
			826	518	143	163	2			
2	H	114	Total	C	N	O	S	0	0	0
			868	541	154	171	2			

- Molecule 3 is a protein called BETA-MAMMAL TOXIN CN2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	65	Total	C	N	O	S	0	0	0
			523	333	88	94	8			
3	F	65	Total	C	N	O	S	0	0	0
			523	333	88	94	8			
3	I	66	Total	C	N	O	S	0	0	0
			529	336	89	96	8			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	9	Total	O	0	0
			9	9		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	42	Total 42	O 42	0	0
4	C	30	Total 30	O 30	0	0
4	D	33	Total 33	O 33	0	0
4	E	27	Total 27	O 27	0	0
4	F	32	Total 32	O 32	0	0
4	G	9	Total 9	O 9	0	0
4	H	23	Total 23	O 23	0	0
4	I	15	Total 15	O 15	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SINGLE CHAIN ANTIBODY FRAGMENT 9004G




- Molecule 1: SINGLE CHAIN ANTIBODY FRAGMENT 9004G



- Molecule 1: SINGLE CHAIN ANTIBODY FRAGMENT 9004G



- Molecule 2: SINGLE CHAIN ANTIBODY FRAGMENT 9004G



- Molecule 2: SINGLE CHAIN ANTIBODY FRAGMENT 9004G



GLN
LYS
LEU
ILE
SER
GLU
GLU
ASP
LEU
ASN
GLY
ALA
ALA
HIS
HIS
HIS
HIS
HIS

• Molecule 2: SINGLE CHAIN ANTIBODY FRAGMENT 9004G

Chain H:  70% 8% 22%

GLY
GLY
GLY
SER
SER
GLY
GLY
GLY
S127
S139
P140
V145
L153
Q170
D182
A183
I190
P191
I207
L210
S226
E240
ALA
ALA
ALA
GLU
GLN
LYS
LEU
ILE
SER
SER
GLU
GLU
ASP
LEU
ASN
GLY
ALA
ALA
HIS
HIS
HIS
HIS
HIS

• Molecule 3: BETA-MAMMAL TOXIN CN2

Chain C:  91% 8%

K1
E2
N9
K18
T49
K63
R64
C65
SER

• Molecule 3: BETA-MAMMAL TOXIN CN2

Chain F:  2% 83% 15%

K1
Y4
K8
K13
C16
L17
K18
N22
D23
Y24
A45
L60
P61
R62
C85
SER

• Molecule 3: BETA-MAMMAL TOXIN CN2

Chain I:  2% 83% 14%

K1
Y4
K8
K13
L17
K18
N22
D23
Y24
A45
L60
P61
R62
K63
S66

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.84Å 104.49Å 152.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.72 – 2.55 45.72 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.72-2.55) 99.9 (45.72-2.55)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.37 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.192 , 0.247 0.187 , 0.240	Depositor DCC
R_{free} test set	1858 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6941	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/885	0.81	0/1195
1	D	0.54	0/882	0.83	0/1191
1	G	0.47	0/882	0.79	0/1191
2	B	0.54	0/870	0.86	2/1179 (0.2%)
2	E	0.55	0/845	0.89	2/1148 (0.2%)
2	H	0.52	0/887	0.87	0/1201
3	C	0.50	0/537	0.79	0/723
3	F	0.54	0/537	0.71	0/723
3	I	0.54	0/543	0.79	0/731
All	All	0.52	0/6868	0.83	4/9282 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	190	ILE	CA-C-N	6.59	126.57	119.78
2	B	190	ILE	C-N-CA	6.59	126.57	119.78
2	E	190	ILE	CA-C-N	5.81	125.74	119.76
2	E	190	ILE	C-N-CA	5.81	125.74	119.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	869	0	829	16	0
1	D	866	0	828	28	0
1	G	866	0	828	24	0
2	B	851	0	825	8	0
2	E	826	0	800	17	0
2	H	868	0	843	9	0
3	C	523	0	492	3	0
3	F	523	0	492	8	0
3	I	529	0	497	7	0
4	A	9	0	0	0	0
4	B	42	0	0	1	0
4	C	30	0	0	0	0
4	D	33	0	0	1	0
4	E	27	0	0	0	0
4	F	32	0	0	0	0
4	G	9	0	0	0	0
4	H	23	0	0	0	0
4	I	15	0	0	0	0
All	All	6941	0	6434	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 105 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:PRO:HD3	1:A:116:SER:O	1.64	0.98
1:D:39:GLN:HE22	2:E:170:GLN:HE22	1.00	0.94
1:G:39:GLN:HE22	2:H:170:GLN:HE22	1.06	0.91
1:G:91:THR:OG1	1:G:115:VAL:HG12	1.83	0.78
1:D:50:GLY:C	3:F:17:LEU:HD21	2.09	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	114/117 (97%)	106 (93%)	8 (7%)	0	100	100
1	D	113/117 (97%)	109 (96%)	4 (4%)	0	100	100
1	G	113/117 (97%)	107 (95%)	6 (5%)	0	100	100
2	B	110/146 (75%)	108 (98%)	2 (2%)	0	100	100
2	E	105/146 (72%)	102 (97%)	3 (3%)	0	100	100
2	H	112/146 (77%)	111 (99%)	1 (1%)	0	100	100
3	C	63/66 (96%)	62 (98%)	1 (2%)	0	100	100
3	F	63/66 (96%)	62 (98%)	1 (2%)	0	100	100
3	I	64/66 (97%)	63 (98%)	1 (2%)	0	100	100
All	All	857/987 (87%)	830 (97%)	27 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	93/94 (99%)	90 (97%)	3 (3%)	34	51
1	D	92/94 (98%)	88 (96%)	4 (4%)	26	39
1	G	92/94 (98%)	88 (96%)	4 (4%)	26	39
2	B	91/111 (82%)	88 (97%)	3 (3%)	33	50
2	E	90/111 (81%)	84 (93%)	6 (7%)	15	21
2	H	93/111 (84%)	91 (98%)	2 (2%)	45	64
3	C	54/55 (98%)	54 (100%)	0	100	100
3	F	54/55 (98%)	53 (98%)	1 (2%)	50	69
3	I	55/55 (100%)	53 (96%)	2 (4%)	31	46
All	All	714/780 (92%)	689 (96%)	25 (4%)	32	48

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	229	THR
1	G	54	SER
3	I	18	LYS
3	F	18	LYS
1	G	64	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	39	GLN
1	G	109	GLN
2	H	232	GLN
2	B	169	GLN
3	C	50	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	116/117 (99%)	-0.07	2 (1%) 69 69	21, 37, 59, 78	0
1	D	115/117 (98%)	-0.32	1 (0%) 81 83	20, 30, 47, 63	0
1	G	115/117 (98%)	0.10	2 (1%) 69 69	22, 42, 65, 79	0
2	B	112/146 (76%)	-0.39	0 100 100	19, 29, 44, 62	0
2	E	107/146 (73%)	-0.32	1 (0%) 81 83	18, 29, 58, 64	0
2	H	114/146 (78%)	-0.38	0 100 100	18, 29, 49, 69	0
3	C	65/66 (98%)	-0.51	0 100 100	17, 25, 50, 70	0
3	F	65/66 (98%)	-0.54	1 (1%) 72 73	17, 22, 42, 72	0
3	I	66/66 (100%)	-0.34	1 (1%) 72 73	18, 24, 57, 73	0
All	All	875/987 (88%)	-0.28	8 (0%) 81 83	17, 30, 58, 79	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	I	62	ASN	4.8
3	F	62	ASN	4.2
2	E	238	ILE	3.3
1	A	117	SER	3.2
1	G	115	VAL	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.