



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 12:27 PM UTC

PDB ID : 4YBJ / pdb_00004ybj
Title : Type II Dasatinib Analog Crystallized with c-Src Kinase
Authors : Kwarcinski, F.E.; Brandvold, K.B.; Johnson, T.K.; Phadke, S.; Meagher, J.L.;
Seeliger, M.A.; Stuckey, J.A.; Soellner, M.B.
Deposited on : 2015-02-18
Resolution : 2.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4951 (2.64-2.60)
Clashscore	190562	5303 (2.64-2.60)
Ramachandran outliers	187476	5217 (2.64-2.60)
Sidechain outliers	187428	5217 (2.64-2.60)
RSRZ outliers	180081	4950 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	10% 86% • 11%
1	B	286	20% 84% 5% 12%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4149 atoms, of which 60 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	255	Total	C	N	O	S	0	0	0
			1987	1280	329	362	16			
1	B	253	Total	C	N	O	S	0	0	0
			1950	1253	330	352	15			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	GLY	-	expression tag	UNP P00523
A	249	HIS	-	expression tag	UNP P00523
A	250	MET	-	expression tag	UNP P00523
B	248	GLY	-	expression tag	UNP P00523
B	249	HIS	-	expression tag	UNP P00523
B	250	MET	-	expression tag	UNP P00523

- Molecule 2 is 2-({6-[4-(2-hydroxyethyl)piperazin-1-yl]-2-methylpyrimidin-4-yl}amino)-N-(3-{[3-(trifluoromethyl)benzoyl]amino}phenyl)-1,3-thiazole-5-carboxamide (CCD ID: 4A9) (formula: C₂₉H₂₉F₃N₈O₃S).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
2	A	1	Total	C	F	H	N	O	S	30	0
			74	29	3	30	8	3	1		
2	B	1	Total	C	F	H	N	O	S	30	0
			74	29	3	30	8	3	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	30	Total	O	0	0
			30	30		
3	B	34	Total	O	0	0
			34	34		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.12Å 120.06Å 63.32Å 90.00° 90.26° 90.00°	Depositor
Resolution (Å)	43.56 – 2.61 43.56 – 2.61	Depositor EDS
% Data completeness (in resolution range)	99.9 (43.56-2.61) 99.9 (43.56-2.61)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.21 (at 2.61Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.10.0	Depositor
R, R_{free}	0.218 , 0.250 0.218 , 0.254	Depositor DCC
R_{free} test set	1034 reflections (5.40%)	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtriage
Anisotropy	0.030	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.158 for h,-k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4149	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 4A9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	0/2036	1.23	2/2767 (0.1%)
1	B	0.77	0/1998	1.21	4/2719 (0.1%)
All	All	0.78	0/4034	1.22	6/5486 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	352	GLY	CA-C-N	6.02	128.63	120.38
1	A	352	GLY	C-N-CA	6.02	128.63	120.38
1	B	377	VAL	CA-C-N	5.13	127.11	120.44
1	B	377	VAL	C-N-CA	5.13	127.11	120.44
1	B	425	PRO	CA-C-N	5.09	127.44	120.46
1	B	425	PRO	C-N-CA	5.09	127.44	120.46

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1987	0	1936	4	0
1	B	1950	0	1859	5	0
2	A	44	30	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	44	30	0	1	0
3	A	30	0	0	0	0
3	B	34	0	0	0	0
All	All	4089	60	3795	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:262:ILE:HG13	1:B:263:PRO:HD2	1.78	0.65
1:A:262:ILE:HG12	1:A:327:ALA:HB1	1.89	0.55
1:B:262:ILE:HG12	1:B:286:TRP:HE1	1.73	0.53
2:B:601:4A9:S	2:B:601:4A9:N5	2.82	0.52
2:A:601:4A9:S	2:A:601:4A9:N5	2.84	0.50
1:A:283:MET:HG3	1:A:340:TYR:CE1	2.49	0.47
1:A:366:MET:HE3	1:A:398:LEU:HG	1.99	0.43
1:B:474:GLN:O	1:B:477:ARG:HG3	2.20	0.42
1:B:269:LEU:HG	1:B:294:ILE:HD13	2.02	0.41
1:A:383:VAL:HG12	1:A:385:ARG:HG3	2.03	0.41
1:B:320:GLU:HG3	1:B:321:LYS:HG2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/286 (87%)	245 (98%)	4 (2%)	0	100	100
1	B	247/286 (86%)	241 (98%)	5 (2%)	1 (0%)	30	49
All	All	496/572 (87%)	486 (98%)	9 (2%)	1 (0%)	43	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	277	CYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/245 (84%)	205 (99%)	2 (1%)	68	85
1	B	195/245 (80%)	194 (100%)	1 (0%)	81	91
All	All	402/490 (82%)	399 (99%)	3 (1%)	76	89

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	296	THR
1	A	398	LEU
1	B	322	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	4A9	A	601	-	48,48,48	1.45	8 (16%)	68,68,68	1.14	7 (10%)
2	4A9	B	601	-	48,48,48	1.46	7 (14%)	68,68,68	1.07	4 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4A9	A	601	-	-	6/33/43/43	0/5/5/5
2	4A9	B	601	-	-	8/33/43/43	0/5/5/5

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	4A9	C22-N6	3.92	1.40	1.34
2	A	601	4A9	C22-N6	3.73	1.39	1.34
2	A	601	4A9	C18-N4	3.39	1.41	1.36
2	B	601	4A9	C18-N4	3.29	1.41	1.36
2	B	601	4A9	C18-N3	3.15	1.36	1.31
2	A	601	4A9	C19-N5	2.92	1.40	1.34
2	B	601	4A9	C19-N5	2.90	1.40	1.34
2	A	601	4A9	C18-N3	2.85	1.36	1.31
2	B	601	4A9	C16-S	-2.80	1.67	1.73
2	A	601	4A9	C16-S	-2.75	1.67	1.73
2	B	601	4A9	C22-N7	2.58	1.43	1.37
2	B	601	4A9	C1-N1	2.32	1.40	1.35
2	A	601	4A9	C22-N7	2.26	1.42	1.37
2	A	601	4A9	C1-N1	2.18	1.40	1.35
2	A	601	4A9	C15-C6	2.11	1.42	1.39

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	4A9	C17-C16-C1	-2.97	126.01	130.64
2	B	601	4A9	C17-C16-C1	-2.91	126.11	130.64
2	A	601	4A9	N4-C19-N5	2.55	119.50	113.33
2	A	601	4A9	C27-C28-N7	-2.53	105.47	110.78
2	B	601	4A9	N4-C19-N5	2.52	119.44	113.33
2	A	601	4A9	C17-C16-S	2.24	112.84	109.51
2	A	601	4A9	C2-N1-C1	-2.23	123.50	127.45
2	B	601	4A9	C17-C16-S	2.17	112.74	109.51
2	B	601	4A9	N6-C22-N7	-2.12	113.89	116.56
2	A	601	4A9	N6-C22-N7	-2.06	113.96	116.56
2	A	601	4A9	C8-C9-C10	-2.04	118.39	121.04

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	4A9	O2-C7-C8-C9
2	A	601	4A9	N2-C7-C8-C9
2	A	601	4A9	O2-C7-C8-C14
2	A	601	4A9	C3-C2-N1-C1
2	A	601	4A9	N2-C7-C8-C14
2	A	601	4A9	C15-C2-N1-C1
2	B	601	4A9	C3-C2-N1-C1
2	B	601	4A9	C15-C2-N1-C1
2	B	601	4A9	O2-C7-C8-C9
2	B	601	4A9	N2-C7-C8-C9
2	B	601	4A9	O2-C7-C8-C14
2	B	601	4A9	N2-C7-C8-C14
2	B	601	4A9	O1-C1-C16-S
2	B	601	4A9	N1-C1-C16-S


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	4A9	1	0
2	B	601	4A9	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	255/286 (89%)	0.57	28 (10%) 10 8	13, 33, 92, 131	0
1	B	253/286 (88%)	0.92	57 (22%) 2 2	13, 33, 110, 184	0
All	All	508/572 (88%)	0.75	85 (16%) 4 3	13, 33, 107, 184	0

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	267	LEU	7.7
1	B	259	ALA	7.2
1	B	260	TRP	6.9
1	A	259	ALA	6.8
1	B	262	ILE	6.7
1	B	269	LEU	6.5
1	B	263	PRO	6.3
1	B	264	ARG	6.2
1	A	262	ILE	5.8
1	B	261	GLU	5.8
1	A	261	GLU	5.6
1	B	287	ASN	5.5
1	B	265	GLU	5.4
1	B	282	TRP	5.3
1	A	260	TRP	4.9
1	A	263	PRO	4.8
1	B	268	ARG	4.8
1	A	264	ARG	4.7
1	A	269	LEU	4.6
1	B	277	CYS	4.6
1	B	330	SER	4.4
1	B	280	GLU	4.3
1	B	288	GLY	4.2
1	B	335	TYR	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	267	LEU	4.1
1	A	266	SER	4.0
1	A	406	GLY	3.9
1	B	266	SER	3.9
1	B	270	GLU	3.9
1	B	296	THR	3.8
1	B	305	GLU	3.8
1	B	276	GLY	3.8
1	A	268	ARG	3.6
1	A	276	GLY	3.6
1	B	329	VAL	3.6
1	B	285	THR	3.6
1	B	271	VAL	3.6
1	B	289	THR	3.5
1	B	332	GLU	3.4
1	A	265	GLU	3.3
1	B	273	LEU	3.3
1	B	283	MET	3.3
1	A	287	ASN	3.1
1	B	290	THR	3.1
1	B	284	GLY	3.1
1	B	291	ARG	3.1
1	A	289	THR	3.0
1	B	286	TRP	3.0
1	B	272	LYS	2.9
1	B	302	MET	2.9
1	B	307	PHE	2.9
1	B	275	GLN	2.8
1	B	303	SER	2.8
1	A	299	PRO	2.8
1	B	292	VAL	2.8
1	A	333	PRO	2.7
1	B	281	VAL	2.7
1	B	333	PRO	2.6
1	B	326	TYR	2.6
1	B	406	GLY	2.6
1	B	328	VAL	2.6
1	B	278	PHE	2.6
1	A	472	LEU	2.5
1	B	304	PRO	2.5
1	B	337	VAL	2.5
1	B	424	PHE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	308	LEU	2.4
1	B	331	GLU	2.4
1	A	312	GLN	2.3
1	B	316	LYS	2.3
1	A	330	SER	2.3
1	A	300	GLY	2.3
1	A	305	GLU	2.2
1	B	473	ASP	2.2
1	B	477	ARG	2.2
1	B	334	ILE	2.2
1	A	270	GLU	2.2
1	A	279	GLY	2.2
1	B	336	ILE	2.1
1	A	286	TRP	2.1
1	B	438	ARG	2.1
1	A	308	LEU	2.1
1	A	478	GLY	2.1
1	B	309	GLN	2.1
1	A	284	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	4A9	A	601	44/44	0.83	0.17	41,49,57,58	30
2	4A9	B	601	44/44	0.86	0.15	33,43,50,51	30

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 4A9 A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 4A9 B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.