



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:22 PM UTC

PDB ID : 5YC6 / pdb_00005yc6
Title : The crystal structure of uPA in complex with 4-Bromobenzylamine at pH4.6
Authors : Jiang, L.G.; Zhang, X.; Huang, M.D.
Deposited on : 2017-09-06
Resolution : 1.18 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1789 (1.20-1.16)
Clashscore	190562	1845 (1.20-1.16)
Ramachandran outliers	187476	1789 (1.20-1.16)
Sidechain outliers	187428	1789 (1.20-1.16)
RSRZ outliers	180081	1787 (1.20-1.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	U	246	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PZH	U	301	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Urokinase-type plasminogen activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	U	246	1942	1226	340	361	15	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
U	122	ALA	CYS	conflict	UNP P00749
U	145	GLN	ASN	conflict	UNP P00749

- Molecule 2 is 1-(4-BROMOPHENYL)METHANAMINE (CCD ID: PZH) (formula: C_7H_8BrN).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	Br	C	N		
2	U	1	9	1	7	1	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	U	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	U	1	Total	C	O	0	0
			10	6	4		
4	U	1	Total	C	O	0	0
			10	6	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	U	214	Total 214	O 214	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Urokinase-type plasminogen activator



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	120.85Å 120.85Å 42.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 1.18 50.00 – 1.18	Depositor EDS
% Data completeness (in resolution range)	95.9 (50.00-1.18) 96.3 (50.00-1.18)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 1.18Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.197 , 0.228 0.203 , 0.232	Depositor DCC
R_{free} test set	3695 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	12.3	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 38.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.016 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2190	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PZH, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	U	0.82	8/1991 (0.4%)	0.82	2/2697 (0.1%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	U	97(B)	LEU	C-N	-7.52	1.23	1.33
1	U	170(A)	PRO	N-CD	7.10	1.57	1.47
1	U	71	SER	C-O	-5.88	1.16	1.24
1	U	99	HIS	C-O	-5.78	1.17	1.24
1	U	34	TYR	C-O	-5.73	1.16	1.23
1	U	72	ARG	C-O	-5.39	1.17	1.24
1	U	231	VAL	C-O	-5.35	1.18	1.24
1	U	39	THR	C-O	-5.14	1.17	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	U	97(B)	LEU	CA-C-N	6.54	130.15	120.87
1	U	97(B)	LEU	C-N-CA	6.54	130.15	120.87

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	U	1942	0	1895	19	2
2	U	9	0	8	6	0
3	U	5	0	0	1	0
4	U	20	0	28	0	0
5	U	214	0	0	6	0
All	All	2190	0	1931	21	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:U:301:PZH:HE2	5:U:493:HOH:O	1.75	0.86
2:U:301:PZH:BR	5:U:516:HOH:O	2.59	0.76
1:U:84:GLU:OE1	5:U:401:HOH:O	2.10	0.68
1:U:37:HIS:HE1	1:U:39:THR:OG1	1.79	0.65
1:U:57:HIS:HD2	1:U:102:ASP:OD2	1.80	0.64
1:U:57:HIS:HE1	3:U:302:SO4:O2	1.84	0.61
1:U:126:MET:HG2	1:U:127:TYR:HD1	1.66	0.59
1:U:72:ARG:NH2	1:U:153:GLU:HG2	2.20	0.57
1:U:22:THR:HG21	1:U:157:MET:HG3	1.88	0.56
1:U:72:ARG:NH2	1:U:153:GLU:CG	2.71	0.54
1:U:126:MET:HG2	1:U:127:TYR:CD1	2.48	0.48
1:U:101:ASN:HD21	1:U:179:LYS:NZ	2.12	0.48
1:U:89:ILE:HD11	5:U:403:HOH:O	2.13	0.48
1:U:189:ASP:CG	2:U:301:PZH:BR	3.12	0.47
1:U:72:ARG:HH21	1:U:153:GLU:HG2	1.80	0.46
1:U:87:ASN:ND2	5:U:403:HOH:O	2.49	0.46
1:U:189:ASP:OD1	2:U:301:PZH:BR	2.91	0.44
1:U:224:LYS:NZ	5:U:404:HOH:O	2.52	0.43
1:U:191:CYS:HA	2:U:301:PZH:CE1	2.48	0.42
1:U:189:ASP:OD2	2:U:301:PZH:BR	2.94	0.41
1:U:72:ARG:CZ	1:U:153:GLU:HG3	2.51	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:U:86:GLU:OE1	1:U:166:ARG:NH2[5_454]	1.94	0.26
1:U:86:GLU:CD	1:U:166:ARG:NH2[5_454]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	U	244/246 (99%)	238 (98%)	5 (2%)	1 (0%)	30	10

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	U	97(A)	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	U	214/214 (100%)	208 (97%)	6 (3%)	38	6

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	U	36	ARG
1	U	97	ASP
1	U	97(A)	THR
1	U	97(B)	LEU
1	U	109	ARG
1	U	187	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	U	37	HIS
1	U	57	HIS
1	U	87	ASN
1	U	99	HIS
1	U	101	ASN
1	U	169	GLN
1	U	170(B)	HIS
1	U	204	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PGE	U	303	-	9,9,9	0.41	0	8,8,8	0.39	0
4	PGE	U	304	-	9,9,9	0.51	0	8,8,8	0.99	0
3	SO4	U	302	-	4,4,4	0.37	0	6,6,6	0.33	0
2	PZH	U	301	-	9,9,9	0.38	0	11,11,11	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PGE	U	303	-	-	0/7/7/7	-
4	PGE	U	304	-	-	2/7/7/7	-
2	PZH	U	301	-	-	0/2/2/2	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	U	304	PGE	O3-C5-C6-O4
4	U	304	PGE	O2-C3-C4-O3

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	U	302	SO4	1	0
2	U	301	PZH	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	U	246/246 (100%)	1.35	62 (25%) 1 3	8, 15, 32, 40	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	U	97(A)	THR	7.5
1	U	97(B)	LEU	5.9
1	U	127	TYR	5.8
1	U	97	ASP	5.4
1	U	96	ALA	5.2
1	U	204	GLN	5.0
1	U	37(B)	GLY	4.5
1	U	181	LEU	4.5
1	U	132	PHE	4.5
1	U	242	THR	4.3
1	U	37(C)	GLY	4.2
1	U	110(D)	ARG	4.1
1	U	37(A)	ARG	4.1
1	U	170(A)	PRO	4.1
1	U	175	GLU	4.0
1	U	131	GLN	3.9
1	U	243	LYS	3.9
1	U	126	MET	3.8
1	U	206	ARG	3.5
1	U	37(D)	SER	3.5
1	U	203	LEU	3.4
1	U	240	SER	3.3
1	U	98	ALA	3.3
1	U	113	GLN	3.3
1	U	114	PRO	3.2
1	U	60	ILE	3.1
1	U	173	GLY	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	U	111	CYS	3.0
1	U	110(B)	GLU	3.0
1	U	166	ARG	3.0
1	U	170(B)	HIS	2.9
1	U	60(B)	TYR	2.9
1	U	174	SER	2.9
1	U	112	ALA	2.9
1	U	205	GLY	2.8
1	U	110(A)	LYS	2.8
1	U	178	THR	2.7
1	U	109	ARG	2.7
1	U	170	GLN	2.7
1	U	37	HIS	2.7
1	U	61	LYS	2.7
1	U	124	PRO	2.6
1	U	38	VAL	2.6
1	U	86	GLU	2.5
1	U	110(C)	GLY	2.5
1	U	165	HIS	2.4
1	U	169	GLN	2.4
1	U	125	SER	2.4
1	U	63	ASP	2.3
1	U	76	ASN	2.2
1	U	119	GLN	2.2
1	U	51	TRP	2.2
1	U	123	LEU	2.2
1	U	75	SER	2.2
1	U	151	TYR	2.2
1	U	83	PHE	2.1
1	U	186	TRP	2.1
1	U	128	ASN	2.1
1	U	217	ARG	2.1
1	U	168	CYS	2.0
1	U	72	ARG	2.0
1	U	222	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PZH	U	301	9/9	0.60	0.23	20,22,25,43	0
4	PGE	U	304	10/10	0.82	0.16	32,41,47,54	0
4	PGE	U	303	10/10	0.85	0.16	20,34,42,44	0
3	SO4	U	302	5/5	0.96	0.09	17,18,21,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.