



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:40 PM UTC

PDB ID : 5YD4 / pdb_00005yd4
Title : Crystal structure of the scFv antibody 4B08 with epitope peptide (mutation T6A)
Authors : Caaveiro, J.M.M.; Miyanabe, K.; Tsumoto, K.
Deposited on : 2017-09-11
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1216 (1.36-1.36)
Clashscore	190562	1232 (1.36-1.36)
Ramachandran outliers	187476	1220 (1.36-1.36)
Sidechain outliers	187428	1220 (1.36-1.36)
RSRZ outliers	180081	1214 (1.36-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	2% 88% • 8%
1	C	251	2% 87% • 9%
1	E	251	90% • 8%
1	G	251	28% 80% 8% • 10%
2	B	9	11% 89% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	9	11%89%11%
2	F	9	11%100%
2	H	9	44%67%11%22%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8594 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called scFv 4B08.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	232	Total	C	N	O	S	0	14	0
			1874	1188	314	362	10			
1	C	229	Total	C	N	O	S	0	12	0
			1856	1180	309	357	10			
1	E	230	Total	C	N	O	S	0	7	0
			1824	1158	305	352	9			
1	G	226	Total	C	N	O	S	0	7	0
			1776	1125	298	343	10			

- Molecule 2 is a protein called Epitope peptide (mutation T6A).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	8	Total	C	N	O	0	0	0
			68	44	9	15			
2	D	8	Total	C	N	O	0	0	0
			68	44	9	15			
2	F	9	Total	C	N	O	0	0	0
			76	48	10	18			
2	H	7	Total	C	N	O	0	1	0
			60	38	8	14			

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	1
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	233	Total	O	0	12
			242	242		
4	B	12	Total	O	0	0
			12	12		
4	C	247	Total	O	0	12
			256	256		
4	D	13	Total	O	0	3
			16	16		


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	257	Total 268	O 268	0	13
4	F	19	Total 20	O 20	0	1
4	G	137	Total 138	O 138	0	3





- Molecule 2: Epitope peptide (mutation T6A)



- Molecule 2: Epitope peptide (mutation T6A)



- Molecule 2: Epitope peptide (mutation T6A)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.47Å 72.90Å 86.46Å 90.00° 114.15° 90.00°	Depositor
Resolution (Å)	33.10 – 1.35 33.10 – 1.35	Depositor EDS
% Data completeness (in resolution range)	99.1 (33.10-1.35) 99.1 (33.10-1.35)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.44 (at 1.35Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.135 , 0.173 (Not available) , 0.199	Depositor DCC
R_{free} test set	3797 reflections (1.90%)	wwPDB-VP
Wilson B-factor (Å ²)	10.8	Xtriage
Anisotropy	0.168	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8594	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.7710e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/1958	0.89	1/2649 (0.0%)
1	C	0.86	0/1936	0.92	0/2622
1	E	0.80	0/1887	0.86	1/2557 (0.0%)
1	G	1.00	4/1833 (0.2%)	1.03	5/2486 (0.2%)
2	B	0.79	0/70	0.66	0/94
2	D	0.72	0/70	0.70	0/94
2	F	0.90	0/78	0.84	0/105
2	H	0.94	0/62	1.07	0/83
All	All	0.87	4/7894 (0.1%)	0.92	7/10690 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	233	CYS	CA-CB	7.77	1.66	1.53
1	G	242	THR	C-O	-6.41	1.15	1.23
1	G	242	THR	CA-C	5.75	1.59	1.52
1	G	217	THR	N-CA	-5.60	1.39	1.46

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	252	LYS	N-CA-C	9.47	123.96	110.23
1	G	144	MET	CG-SD-CE	6.37	114.91	100.90
1	G	105	ASN	CB-CA-C	-5.93	100.07	109.80
1	G	10	GLY	N-CA-C	5.72	122.06	112.85
1	G	242	THR	CA-C-O	5.33	127.35	121.11
1	E	153	VAL	N-CA-CB	-5.27	102.80	111.44
1	G	233	CYS	CB-CA-C	5.24	118.99	109.83

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1874	0	1835	6	0
1	C	1856	0	1807	8	0
1	E	1824	0	1776	3	0
1	G	1776	0	1713	21	0
2	B	68	0	57	0	0
2	D	68	0	57	0	0
2	F	76	0	64	0	0
2	H	60	0	46	3	0
3	A	10	0	0	0	0
3	C	15	0	0	0	0
3	E	15	0	0	0	0
4	A	242	0	0	0	0
4	B	12	0	0	0	0
4	C	256	0	0	2	0
4	D	16	0	0	0	0
4	E	268	0	0	2	0
4	F	20	0	0	0	0
4	G	138	0	0	2	0
All	All	8594	0	7355	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:144:MET:HE1	1:G:169:LEU:HD11	1.29	1.14
1:G:104:THR:HG21	2:H:9[A]:PRO:HG3	1.55	0.87
1:G:144:MET:CE	1:G:169:LEU:HD11	2.09	0.77
1:G:104:THR:CG2	2:H:9[A]:PRO:HG3	2.17	0.74
1:G:104:THR:HG21	2:H:9[A]:PRO:CG	2.24	0.66
1:G:175:ASN:ND2	4:G:301:HOH:O	2.28	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:235:GLN:HE22	1:G:238:GLU:H	1.48	0.62
1:G:175:ASN:OD1	1:G:195[B]:ARG:NE	2.34	0.60
1:G:196[B]:MET:SD	1:G:197:SER:HB3	2.49	0.53
1:A:93[A]:SER:HB3	1:G:222:ARG:HH22	1.75	0.52
1:C:251:LEU:HB3	4:C:435[A]:HOH:O	2.09	0.50
1:G:235:GLN:HE21	1:G:237:LEU:H	1.60	0.50
1:C:158:SER:HB3	1:C:221:SER:O	2.12	0.50
1:E:219[B]:ARG:HG3	4:E:457[B]:HOH:O	2.10	0.49
1:G:235:GLN:NE2	1:G:237:LEU:H	2.11	0.49
1:G:144:MET:HE1	1:G:169:LEU:CD1	2.20	0.48
1:A:219[B]:ARG:NH1	1:A:221[B]:SER:OG	2.47	0.48
1:G:5:GLN:NE2	1:G:7:GLN:OE1	2.47	0.47
1:G:144:MET:HE2	1:G:235:GLN:HB3	1.96	0.47
1:C:42[B]:ARG:HD2	4:C:453:HOH:O	2.15	0.46
1:C:5[B]:GLN:HE21	1:C:7:GLN:HE21	1.65	0.45
1:A:251:LEU:HD12	1:G:198:ASN:HD21	1.81	0.45
1:C:5[A]:GLN:HB2	1:C:27:ALA:HB3	1.99	0.44
1:G:176:THR:O	1:G:195[B]:ARG:HA	2.15	0.44
1:A:14:VAL:HG21	1:A:88:LEU:HD13	2.00	0.44
1:E:223:VAL:HG12	1:E:251:LEU:HD21	2.00	0.43
1:C:234[B]:MET:HE1	1:C:236:HIS:HB2	2.00	0.43
1:A:11:ALA:HB3	1:G:157:GLU:HG2	2.00	0.43
1:C:234[B]:MET:HB2	1:C:243:PHE:CD2	2.54	0.42
1:A:15:ARG:HH11	1:A:118:SER:CB	2.32	0.41
1:E:42:ARG:HD2	4:E:470:HOH:O	2.21	0.41
1:C:42[B]:ARG:HD2	1:C:42[B]:ARG:HH11	1.71	0.40
1:G:164:ARG:NH2	4:G:306:HOH:O	2.50	0.40
1:G:95:ILE:HD13	1:G:114:SER:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	242/251 (96%)	240 (99%)	2 (1%)	0	100	100
1	C	237/251 (94%)	235 (99%)	2 (1%)	0	100	100
1	E	233/251 (93%)	230 (99%)	3 (1%)	0	100	100
1	G	226/251 (90%)	219 (97%)	7 (3%)	0	100	100
2	B	6/9 (67%)	6 (100%)	0	0	100	100
2	D	6/9 (67%)	6 (100%)	0	0	100	100
2	F	7/9 (78%)	7 (100%)	0	0	100	100
2	H	5/9 (56%)	5 (100%)	0	0	100	100
All	All	962/1040 (92%)	948 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/199 (105%)	207 (99%)	2 (1%)	68	40
1	C	206/199 (104%)	206 (100%)	0	100	100
1	E	202/199 (102%)	202 (100%)	0	100	100
1	G	195/199 (98%)	193 (99%)	2 (1%)	68	40
2	B	7/8 (88%)	7 (100%)	0	100	100
2	D	7/8 (88%)	7 (100%)	0	100	100
2	F	8/8 (100%)	8 (100%)	0	100	100
2	H	6/8 (75%)	6 (100%)	0	100	100
All	All	840/828 (101%)	836 (100%)	4 (0%)	81	64

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	226	GLU
1	A	252	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	175	ASN
1	G	235	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	61	ASN
1	C	41	GLN
1	C	45	HIS
1	C	183	GLN
1	C	198	ASN
1	G	5	GLN
1	G	7	GLN
1	G	198	ASN
1	G	235	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	301	-	4,4,4	0.47	0	6,6,6	0.33	0
3	SO4	A	302	-	4,4,4	0.36	0	6,6,6	0.74	0
3	SO4	E	302	-	4,4,4	0.51	0	6,6,6	0.52	0
3	SO4	C	303	-	4,4,4	0.43	0	6,6,6	0.38	0
3	SO4	E	301	-	4,4,4	0.65	0	6,6,6	0.53	0
3	SO4	C	302[A]	-	4,4,4	0.42	0	6,6,6	0.40	0
3	SO4	C	301	-	4,4,4	0.40	0	6,6,6	1.11	0
3	SO4	E	303	-	4,4,4	0.57	0	6,6,6	0.46	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	232/251 (92%)	-0.30	5 (2%)	62	72	5, 10, 26, 45	14 (6%)
1	C	229/251 (91%)	-0.24	5 (2%)	62	72	5, 11, 26, 49	12 (5%)
1	E	230/251 (91%)	-0.10	1 (0%)	88	93	7, 14, 27, 46	7 (3%)
1	G	226/251 (90%)	1.48	71 (31%)	1	2	7, 24, 41, 53	7 (3%)
2	B	8/9 (88%)	0.56	1 (12%)	8	12	12, 14, 28, 37	0
2	D	8/9 (88%)	0.16	1 (12%)	8	12	9, 11, 19, 27	0
2	F	9/9 (100%)	0.34	1 (11%)	10	15	13, 16, 28, 53	0
2	H	7/9 (77%)	4.44	4 (57%)	0	0	22, 35, 63, 68	1 (14%)
All	All	949/1040 (91%)	0.24	89 (9%)	14	19	5, 14, 34, 68	41 (4%)

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	9[A]	PRO	10.3
2	H	4	TYR	6.5
1	G	177	TYR	6.0
2	H	5	TYR	5.7
1	A	253	VAL	5.5
1	G	102	TYR	5.1
1	G	169	LEU	5.0
1	G	103	TYR	4.8
1	G	4	VAL	4.4
1	C	251	LEU	4.3
1	G	117	VAL	4.2
1	G	142	ILE	4.2
1	G	170	LEU	4.1
1	G	239	TYR	4.1
1	G	194[B]	TYR	4.0
1	G	193[B]	ILE	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	2	ILE	3.9
1	G	180	TRP	3.9
1	G	178	LEU	3.8
1	G	171	HIS	3.8
2	H	6	ALA	3.8
1	G	104	THR	3.8
1	G	213	GLY	3.7
1	G	176	THR	3.7
1	G	216	PHE	3.5
1	C	252	LYS	3.5
1	G	14	VAL	3.5
1	G	214	THR	3.4
1	G	13	LEU	3.4
1	G	68	GLY	3.4
1	G	200	ALA	3.4
1	G	204	PRO	3.4
1	G	237	LEU	3.4
1	G	179	PHE	3.4
1	G	241	TYR	3.4
1	G	143	VAL	3.4
1	G	225	ALA	3.4
1	G	202	GLY	3.3
1	G	163	CYS	3.3
1	G	11	ALA	3.3
1	G	166	SER	3.2
1	G	203	VAL	3.2
1	G	207	PHE	3.2
1	G	89	THR	3.2
1	A	118	SER	3.1
1	G	28	GLY	3.1
1	G	251	LEU	3.0
1	G	115	VAL	3.0
1	G	88	LEU	3.0
1	G	211	GLY	3.0
1	G	118	SER	2.9
1	G	236	HIS	2.9
1	G	164	ARG	2.8
1	G	220	ILE	2.8
1	G	141	ASP	2.8
1	C	118	SER	2.8
1	C	213	GLY	2.7
1	G	199	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	172	ARG	2.6
1	G	198	ASN	2.5
1	G	90	SER	2.5
1	G	116	THR	2.4
1	G	205	ASP	2.4
1	A	226	GLU	2.3
1	C	221	SER	2.3
1	G	233	CYS	2.3
1	G	162	SER	2.3
1	G	210	SER	2.3
1	G	18	THR	2.3
2	D	2	ILE	2.3
1	G	173	ASN	2.3
1	G	27	ALA	2.3
1	G	165	SER	2.3
1	G	145	THR	2.3
1	G	155	PRO	2.3
1	G	238	GLU	2.2
1	A	252	LYS	2.2
1	E	118	SER	2.2
1	G	156	GLY	2.1
1	G	196[A]	MET	2.1
1	G	235	GLN	2.1
1	A	138	GLY	2.1
1	G	174	GLY	2.1
1	G	186	GLY	2.1
1	G	218	LEU	2.1
2	F	2	ILE	2.1
1	G	59	TYR	2.1
1	G	215	ALA	2.0
1	G	167	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	C	302[A]	5/5	0.87	0.11	30,31,44,47	5
3	SO4	C	303	5/5	0.93	0.12	29,30,40,47	5
3	SO4	A	301	5/5	0.94	0.08	43,44,53,57	0
3	SO4	E	301	5/5	0.96	0.14	16,17,20,21	5
3	SO4	C	301	5/5	0.97	0.07	20,23,32,34	0
3	SO4	E	302	5/5	0.97	0.06	25,25,27,28	5
3	SO4	E	303	5/5	0.97	0.06	24,28,33,40	0
3	SO4	A	302	5/5	0.98	0.06	20,25,32,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.