



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:37 PM UTC

PDB ID : 1YFQ / pdb_00001yfq
Title : High resolution S. cerevisiae Bub3 mitotic checkpoint protein
Authors : Wilson, D.K.; Cerna, D.; Chew, E.
Deposited on : 2005-01-03
Resolution : 1.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1656 (1.12-1.08)
Ramachandran outliers	187476	1616 (1.12-1.08)
Sidechain outliers	187428	1612 (1.12-1.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	342	 83% 13% . .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3278 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell cycle arrest protein BUB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	0	0
			2709	1714	454	528	13			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	342	PRO	-	cloning artifact	UNP P26449

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Ca	0	0
			4	4		

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	561	Total	O	0	0
			561	561		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Cell cycle arrest protein BUB3



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.27Å 74.09Å 94.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.10	Depositor
% Data completeness (in resolution range)	(Not available) (30.00-1.10)	Depositor
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.152 , 0.186	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3278	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.95	0/2757	1.54	49/3734 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	ASP	CA-CB-CG	15.40	128.00	112.60
1	A	143	ASN	CA-CB-CG	10.43	123.03	112.60
1	A	148	LYS	CA-C-N	9.83	139.67	121.97
1	A	148	LYS	C-N-CA	9.83	139.67	121.97
1	A	231	ASN	CA-C-N	-9.66	103.08	121.54
1	A	231	ASN	C-N-CA	-9.66	103.08	121.54
1	A	143	ASN	CA-C-N	9.37	138.57	121.70
1	A	143	ASN	C-N-CA	9.37	138.57	121.70
1	A	106	ARG	CG-CD-NE	9.09	132.01	112.00
1	A	145	ASN	CA-CB-CG	-8.53	104.08	112.60
1	A	314	ASP	CA-CB-CG	8.22	120.82	112.60
1	A	235	ARG	CD-NE-CZ	8.02	135.62	124.40
1	A	230	TYR	CA-C-N	7.91	134.95	122.61
1	A	230	TYR	C-N-CA	7.91	134.95	122.61
1	A	231	ASN	CA-CB-CG	7.68	120.28	112.60
1	A	288	ASN	CA-CB-CG	7.67	120.27	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	197	ARG	CG-CD-NE	6.93	127.25	112.00
1	A	304	ASP	CA-CB-CG	6.92	119.52	112.60
1	A	341	ASN	CA-CB-CG	6.82	119.42	112.60
1	A	143	ASN	CA-C-O	-6.45	111.29	120.51
1	A	106	ARG	CD-NE-CZ	6.38	133.34	124.40
1	A	326	ILE	CB-CG1-CD1	-6.35	100.46	113.80
1	A	65	ILE	CA-C-O	-6.29	113.84	120.39
1	A	299	LYS	CG-CD-CE	-6.12	97.23	111.30
1	A	7	GLU	N-CA-C	5.97	119.04	111.69
1	A	87	ILE	O-C-N	-5.90	117.44	121.98
1	A	41	ASP	CA-CB-CG	5.89	118.49	112.60
1	A	239	ARG	CA-CB-CG	5.76	125.62	114.10
1	A	232	SER	CA-C-O	-5.74	112.31	120.51
1	A	143	ASN	OD1-CG-ND2	5.62	128.22	122.60
1	A	162	ARG	NE-CZ-NH1	5.60	127.10	121.50
1	A	141	ASN	OD1-CG-ND2	-5.54	117.06	122.60
1	A	326	ILE	CA-CB-CG1	5.50	119.75	110.40
1	A	242	ARG	NE-CZ-NH2	5.41	124.07	119.20
1	A	68	THR	N-CA-CB	5.40	118.25	110.20
1	A	215	ASP	CA-CB-CG	5.38	117.98	112.60
1	A	222	PHE	CA-CB-CG	-5.36	108.44	113.80
1	A	197	ARG	CD-NE-CZ	5.36	131.90	124.40
1	A	168	ASN	OD1-CG-ND2	-5.32	117.28	122.60
1	A	130	ARG	CD-NE-CZ	5.30	131.81	124.40
1	A	325	THR	CA-C-N	-5.29	113.77	121.71
1	A	325	THR	C-N-CA	-5.29	113.77	121.71
1	A	85	ASP	CA-C-O	-5.24	114.56	120.32
1	A	195	GLN	OE1-CD-NE2	5.19	127.79	122.60
1	A	340	GLU	CB-CG-CD	5.09	121.26	112.60
1	A	143	ASN	O-C-N	-5.09	115.82	122.59
1	A	197	ARG	NE-CZ-NH1	5.07	126.57	121.50
1	A	149	VAL	N-CA-CB	5.07	119.59	111.23
1	A	162	ARG	NE-CZ-NH2	-5.06	114.65	119.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	144	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2709	0	2685	19	0
2	A	4	0	0	0	0
3	A	4	0	3	0	0
4	A	561	0	0	4	0
All	All	3278	0	2688	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229:ASP:O	1:A:231:ASN:N	2.04	0.91
1:A:341:ASN:HB3	4:A:2497:HOH:O	1.84	0.76
1:A:224:ASP:HB2	4:A:2197:HOH:O	1.99	0.62
1:A:232:SER:O	1:A:235:ARG:HG2	2.06	0.56
1:A:150:LYS:CE	1:A:152:LYS:HB2	2.37	0.54
1:A:45:LYS:HE3	1:A:338:ASP:OD2	2.11	0.50
1:A:144:SER:O	1:A:146:ASN:OD1	2.29	0.50
1:A:134:ASP:HB2	4:A:2257:HOH:O	2.11	0.49
1:A:134:ASP:HB2	4:A:2189:HOH:O	2.12	0.48
1:A:150:LYS:HG2	1:A:151:ASN:N	2.28	0.48
1:A:229:ASP:O	1:A:230:TYR:C	2.57	0.47
1:A:149:VAL:O	1:A:149:VAL:HG12	2.15	0.46
1:A:143:ASN:O	1:A:146:ASN:ND2	2.49	0.45
1:A:69:ASP:OD1	1:A:69:ASP:N	2.49	0.43
1:A:202:LEU:HG	1:A:208:GLY:C	2.43	0.43
1:A:229:ASP:C	1:A:231:ASN:N	2.75	0.43
1:A:116:ILE:HD13	1:A:163:LEU:HD12	2.00	0.42
1:A:229:ASP:OD1	1:A:233:SER:HB2	2.19	0.42
1:A:169:ASN:O	1:A:171:GLN:HG3	2.19	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	340/342 (99%)	322 (95%)	14 (4%)	4 (1%)	10 1

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	148	LYS
1	A	149	VAL
1	A	230	TYR
1	A	145	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	306/306 (100%)	293 (96%)	13 (4%)	26 2

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	GLU
1	A	87	ILE
1	A	89	SER
1	A	149	VAL
1	A	150	LYS
1	A	179	LEU
1	A	204	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	205	GLU
1	A	235	ARG
1	A	309	LEU
1	A	317	LYS
1	A	326	ILE
1	A	341	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	67	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACT	A	900	-	3,3,3	0.67	0	3,3,3	0.81	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.