



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 09:42 PM UTC

PDB ID : 2YFC / pdb_00002yfc
Title : STRUCTURAL AND FUNCTIONAL INSIGHTS OF DR2231 PROTEIN,
THE MAZG-LIKE NUCLEOSIDE TRIPHOSPHATE PYROPHOSPHO-
HYDROLASE FROM DEINOCOCCUS RADIODURANS, COMPLEXED
WITH Mn and dUMP
Authors : Goncalves, A.M.D.; De Sanctis, D.; Mcsweeney, S.M.
Deposited on : 2011-04-05
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	154	3% 78% 10% • 11%
1	B	154	5% 69% 13% • 15%
1	C	154	% 80% 9% 11%
1	D	154	4% 76% 7% • 16%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	1147	-	-	X	-
4	GOL	B	1148	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8621 atoms, of which 4120 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MAZG-LIKE NUCLEOSIDE TRIPHOSPHATE PYROPHOSPHOHYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	137	Total	C	H	N	O	0	0	0
			2094	655	1041	199	199			
1	B	131	Total	C	H	N	O	0	2	0
			2006	632	995	188	191			
1	C	137	Total	C	H	N	O	0	0	0
			2086	652	1038	197	199			
1	D	130	Total	C	H	N	O	0	2	0
			1997	627	992	188	190			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	expression tag	UNP Q9RS96
A	-4	ILE	-	expression tag	UNP Q9RS96
A	-3	ASP	-	expression tag	UNP Q9RS96
A	-2	PRO	-	expression tag	UNP Q9RS96
A	-1	PHE	-	expression tag	UNP Q9RS96
A	0	THR	-	expression tag	UNP Q9RS96
B	-5	GLY	-	expression tag	UNP Q9RS96
B	-4	ILE	-	expression tag	UNP Q9RS96
B	-3	ASP	-	expression tag	UNP Q9RS96
B	-2	PRO	-	expression tag	UNP Q9RS96
B	-1	PHE	-	expression tag	UNP Q9RS96
B	0	THR	-	expression tag	UNP Q9RS96
C	-5	GLY	-	expression tag	UNP Q9RS96
C	-4	ILE	-	expression tag	UNP Q9RS96
C	-3	ASP	-	expression tag	UNP Q9RS96
C	-2	PRO	-	expression tag	UNP Q9RS96
C	-1	PHE	-	expression tag	UNP Q9RS96
C	0	THR	-	expression tag	UNP Q9RS96
D	-5	GLY	-	expression tag	UNP Q9RS96
D	-4	ILE	-	expression tag	UNP Q9RS96

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-3	ASP	-	expression tag	UNP Q9RS96
D	-2	PRO	-	expression tag	UNP Q9RS96
D	-1	PHE	-	expression tag	UNP Q9RS96
D	0	THR	-	expression tag	UNP Q9RS96

- Molecule 2 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (CCD ID: UMP) (formula: $C_9H_{13}N_2O_8P$).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	0	0
			31	9	11	2	8	1		
2	C	1	Total	C	H	N	O	P	0	0
			31	9	11	2	8	1		

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		
3	B	2	Total	Mn	0	0
			2	2		
3	C	1	Total	Mn	0	0
			1	1		
3	D	2	Total	Mn	0	0
			2	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	B	1	Total	C	H	O	0	0
			14	3	8	3		
4	C	1	Total	C	H	O	0	0
			14	3	8	3		
4	C	1	Total	C	H	O	0	0
			14	3	8	3		



- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	85	Total	O	0	0
			85	85		
5	B	91	Total	O	0	0
			91	91		
5	C	64	Total	O	0	0
			64	64		
5	D	74	Total	O	0	0
			74	74		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MAZG-LIKE NUCLEOSIDE TRIPHOSPHATE PYROPHOSPHOHYDROLASE





- Molecule 1: MAZG-LIKE NUCLEOSIDE TRIPHOSPHATE PYROPHOSPHOHYDROLASE



- Molecule 1: MAZG-LIKE NUCLEOSIDE TRIPHOSPHATE PYROPHOSPHOHYDROLASE



- Molecule 1: MAZG-LIKE NUCLEOSIDE TRIPHOSPHATE PYROPHOSPHOHYDROLASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	78.73Å 150.19Å 52.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.01 20.00 – 2.01	Depositor EDS
% Data completeness (in resolution range)	98.6 (20.00-2.01) 98.5 (20.00-2.01)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.33 (at 2.01Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.177 , 0.219 0.171 , 0.211	Depositor DCC
R_{free} test set	2082 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8621	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 31.94 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0187e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MN, GOL, UMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	6/1074 (0.6%)	0.86	5/1460 (0.3%)
1	B	1.08	5/1041 (0.5%)	0.95	8/1418 (0.6%)
1	C	0.68	3/1069 (0.3%)	0.86	6/1455 (0.4%)
1	D	1.03	3/1033 (0.3%)	0.79	2/1406 (0.1%)
All	All	0.93	17/4217 (0.4%)	0.87	21/5739 (0.4%)

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	126	PRO	C-N	-26.35	0.97	1.33
1	D	27	GLU	C-N	-22.14	1.08	1.33
1	D	28	ARG	C-N	14.35	1.50	1.33
1	B	127	GLU	C-N	8.89	1.46	1.33
1	D	124	LEU	C-N	-8.26	1.19	1.33
1	C	122	LYS	C-N	-7.94	1.22	1.33
1	B	19	ARG	C-N	-7.04	1.24	1.34
1	A	57	HIS	CD2-NE2	-5.73	1.31	1.37
1	A	109	ASN	C-O	-5.63	1.17	1.24
1	C	142	GLN	CD-OE1	5.34	1.33	1.23
1	B	143	HIS	ND1-CE1	5.31	1.37	1.32
1	C	15	HIS	ND1-CE1	5.20	1.37	1.32
1	A	18	HIS	ND1-CE1	5.19	1.37	1.32
1	B	27	GLU	C-N	-5.09	1.27	1.33
1	A	93	GLN	CD-OE1	5.09	1.33	1.23
1	A	58	LEU	C-O	-5.06	1.18	1.24
1	A	123	GLN	CD-OE1	5.05	1.33	1.23

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	28	ARG	O-C-N	8.37	127.76	121.65
1	B	145	PRO	N-CA-CB	8.13	111.94	103.00
1	C	122	LYS	CA-C-N	7.65	131.66	120.95
1	C	122	LYS	C-N-CA	7.65	131.66	120.95
1	B	125	LYS	N-CA-C	7.03	119.83	109.50
1	A	7	PRO	N-CA-CB	6.73	110.41	103.00
1	C	7	PRO	N-CA-CB	6.72	110.39	103.00
1	B	143	HIS	CB-CG-CD2	-6.55	122.68	131.20
1	A	18	HIS	CB-CG-CD2	-6.55	122.69	131.20
1	C	15	HIS	CB-CG-CD2	-6.51	122.73	131.20
1	D	7	PRO	N-CA-CB	6.47	110.11	103.00
1	A	57	HIS	CB-CG-CD2	-6.14	123.22	131.20
1	B	28	ARG	O-C-N	-5.85	117.38	121.65
1	B	127	GLU	O-C-N	-5.82	115.73	122.95
1	A	18	HIS	CB-CG-ND1	5.66	131.19	122.70
1	B	143	HIS	CB-CG-ND1	5.65	131.18	122.70
1	C	15	HIS	CB-CG-ND1	5.51	130.97	122.70
1	C	122	LYS	O-C-N	-5.43	116.21	122.95
1	A	57	HIS	CB-CG-ND1	5.33	130.70	122.70
1	B	127	GLU	CA-C-N	5.25	131.88	121.53
1	B	127	GLU	C-N-CA	5.25	131.88	121.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1053	1041	1038	10	3
1	B	1011	995	979	17	0
1	C	1048	1038	1034	8	0
1	D	1005	992	978	14	0
2	A	20	11	11	0	0
2	C	20	11	10	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
3	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	2	0	0	0	0
4	A	6	8	8	1	3
4	B	6	8	8	5	0
4	C	12	16	16	3	0
5	A	85	0	0	1	0
5	B	91	0	0	1	0
5	C	64	0	0	0	0
5	D	74	0	0	0	0
All	All	4501	4120	4082	45	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:28:ARG:NH2	1:D:144:ALA:HB2	1.18	1.45
1:D:28:ARG:NH2	1:D:144:ALA:CB	1.89	1.33
1:D:28:ARG:CZ	1:D:144:ALA:HB2	1.95	0.96
1:D:28:ARG:HH21	1:D:144:ALA:HB2	1.11	0.87
1:D:28:ARG:HH22	1:D:144:ALA:CB	1.92	0.82
1:B:35:GLU:HA	4:B:1148:GOL:H32	1.64	0.79
1:D:28:ARG:NH2	1:D:144:ALA:HB3	2.05	0.69
1:D:28:ARG:HH21	1:D:144:ALA:CB	1.79	0.69
1:C:12:GLU:O	4:C:1147:GOL:H2	1.95	0.65
1:B:38:ARG:HB2	4:B:1148:GOL:H31	1.81	0.62
1:C:104:GLU:HG3	1:C:132:ALA:HB1	1.83	0.61
1:C:16:GLU:HB2	4:C:1147:GOL:O2	2.00	0.60
1:B:38:ARG:HD2	4:B:1148:GOL:H11	1.85	0.59
1:A:74:ALA:HA	1:B:138:ILE:HD12	1.85	0.59
1:B:7:PRO:N	1:B:8:PRO:HD3	2.18	0.58
1:A:104:GLU:HG3	1:A:132:ALA:HB1	1.88	0.55
1:B:35:GLU:CA	4:B:1148:GOL:H32	2.36	0.54
1:B:7:PRO:N	1:B:8:PRO:CD	2.72	0.53
1:B:144:ALA:O	1:B:145:PRO:CB	2.56	0.53
1:B:38:ARG:CD	4:B:1148:GOL:H11	2.38	0.53
1:C:61:ARG:O	1:C:66:GLU:HB2	2.08	0.53
1:B:115:GLY:HA3	1:B:124:LEU:O	2.10	0.52
1:B:138:ILE:O	1:B:142:GLN:HG3	2.10	0.51
1:D:124:LEU:O	1:D:124:LEU:HD23	2.10	0.51
1:A:130:ARG:HH12	1:C:135:ARG:NH2	2.11	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:36:LEU:HD13	5:B:2073:HOH:O	2.13	0.49
1:A:130:ARG:NH1	1:C:135:ARG:NH2	2.61	0.48
1:D:125:LYS:HD3	1:D:129:TRP:CE3	2.49	0.48
1:A:74:ALA:HA	1:B:138:ILE:CD1	2.44	0.47
1:D:28:ARG:HH22	1:D:144:ALA:HB3	1.68	0.47
1:A:60:ALA:CB	4:A:1147:GOL:H2	2.44	0.47
1:A:32:PRO:HG3	1:A:94:LEU:CD2	2.46	0.45
1:B:126:PRO:HG2	1:B:129:TRP:HB2	1.99	0.45
1:C:135:ARG:H	4:C:1146:GOL:H32	1.81	0.45
1:D:143:HIS:O	1:D:144:ALA:C	2.61	0.44
1:A:130:ARG:NH1	5:A:2073:HOH:O	2.51	0.43
1:C:130:ARG:HA	1:C:131:PRO:HD3	1.89	0.43
1:A:130:ARG:HH11	1:A:130:ARG:HG3	1.82	0.43
1:B:79:GLU:HA	1:B:79:GLU:OE1	2.19	0.42
1:D:36:LEU:HD23	1:D:36:LEU:C	2.45	0.41
1:A:7:PRO:HA	1:A:8:PRO:HD3	1.86	0.41
1:D:25:THR:HG23	1:D:25:THR:O	2.21	0.41
1:B:15:HIS:NE2	1:B:19:ARG:HD2	2.36	0.40
1:B:130:ARG:HA	1:B:131:PRO:HD3	1.83	0.40
1:D:138:ILE:O	1:D:142:GLN:HG3	2.21	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:ARG:HH22	4:A:1147:GOL:O1[4_445]	1.13	0.47
1:A:130:ARG:HH22	4:A:1147:GOL:HO1[4_445]	1.34	0.26
1:A:130:ARG:NH2	4:A:1147:GOL:O1[4_445]	2.01	0.19

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/154 (86%)	131 (98%)	2 (2%)	0	100	100
1	B	129/154 (84%)	126 (98%)	3 (2%)	0	100	100
1	C	135/154 (88%)	132 (98%)	3 (2%)	0	100	100
1	D	128/154 (83%)	125 (98%)	3 (2%)	0	100	100
All	All	525/616 (85%)	514 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	105/120 (88%)	103 (98%)	2 (2%)	50	56
1	B	102/120 (85%)	100 (98%)	2 (2%)	48	54
1	C	104/120 (87%)	103 (99%)	1 (1%)	68	75
1	D	101/120 (84%)	100 (99%)	1 (1%)	68	75
All	All	412/480 (86%)	406 (98%)	6 (2%)	57	64

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LEU
1	A	140	ARG
1	B	14	LEU
1	B	124	LEU
1	C	93	GLN
1	D	14	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	10	ASN
1	C	62	GLN
1	D	62	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	1147	-	5,5,5	0.23	0	5,5,5	0.54	0
4	GOL	B	1148	-	5,5,5	0.24	0	5,5,5	0.67	0
2	UMP	A	1145	3	21,21,21	2.18	5 (23%)	30,31,31	2.24	11 (36%)
2	UMP	C	1144	3	21,21,21	2.16	5 (23%)	30,31,31	2.24	8 (26%)
4	GOL	C	1146	-	5,5,5	0.39	0	5,5,5	0.28	0
4	GOL	C	1147	-	5,5,5	0.43	0	5,5,5	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1147	-	-	4/4/4/4	-
4	GOL	B	1148	-	-	2/4/4/4	-
2	UMP	A	1145	3	-	3/10/22/22	0/2/2/2
2	UMP	C	1144	3	-	3/10/22/22	0/2/2/2
4	GOL	C	1146	-	-	2/4/4/4	-
4	GOL	C	1147	-	-	2/4/4/4	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1145	UMP	O4-C4	6.54	1.37	1.24
2	C	1144	UMP	O4-C4	6.38	1.36	1.24
2	A	1145	UMP	O4'-C1'	3.75	1.50	1.42
2	C	1144	UMP	O4'-C1'	3.56	1.50	1.42
2	C	1144	UMP	C6-N1	3.14	1.45	1.38
2	A	1145	UMP	C6-N1	3.01	1.45	1.38
2	C	1144	UMP	C2-N1	-2.83	1.34	1.38
2	A	1145	UMP	C2-N1	-2.55	1.34	1.38
2	A	1145	UMP	O3'-C3'	-2.55	1.38	1.43
2	C	1144	UMP	O3'-C3'	-2.16	1.38	1.43

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1144	UMP	C4-N3-C2	-5.64	119.62	126.61
2	A	1145	UMP	C4-N3-C2	-5.23	120.11	126.61
2	C	1144	UMP	C5-C4-N3	5.07	121.90	114.80
2	C	1144	UMP	C2'-C1'-N1	4.56	125.21	113.81
2	A	1145	UMP	C5-C4-N3	4.54	121.16	114.80
2	A	1145	UMP	C2'-C1'-N1	4.46	124.96	113.81
2	C	1144	UMP	O4'-C1'-N1	3.75	114.51	107.86
2	C	1144	UMP	N3-C2-N1	3.62	119.61	114.89
2	A	1145	UMP	N3-C2-N1	3.60	119.58	114.89
2	A	1145	UMP	O4'-C1'-N1	3.50	114.08	107.86
2	A	1145	UMP	O4-C4-C5	-2.61	120.65	125.16
2	A	1145	UMP	O5'-C5'-C4'	-2.59	100.17	108.99
2	C	1144	UMP	O4-C4-C5	-2.50	120.85	125.16
2	A	1145	UMP	O5'-P-OP1	2.50	113.19	106.44
2	A	1145	UMP	O3'-C3'-C4'	2.33	118.89	110.07
2	A	1145	UMP	C4'-O4'-C1'	-2.16	104.37	109.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1144	UMP	O4'-C4'-C5'	2.13	116.16	109.33
2	A	1145	UMP	O4'-C4'-C5'	2.04	115.86	109.33
2	C	1144	UMP	O5'-P-OP1	2.02	111.90	106.44

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1145	UMP	C5'-O5'-P-OP2
2	A	1145	UMP	C5'-O5'-P-OP3
2	C	1144	UMP	C5'-O5'-P-OP2
2	C	1144	UMP	C5'-O5'-P-OP3
4	A	1147	GOL	C1-C2-C3-O3
4	B	1148	GOL	C1-C2-C3-O3
4	C	1146	GOL	O1-C1-C2-O2
4	C	1146	GOL	O1-C1-C2-C3
4	C	1147	GOL	O1-C1-C2-C3
4	A	1147	GOL	O2-C2-C3-O3
4	B	1148	GOL	O2-C2-C3-O3
4	A	1147	GOL	O1-C1-C2-C3
4	C	1147	GOL	O1-C1-C2-O2
4	A	1147	GOL	O1-C1-C2-O2
2	A	1145	UMP	C5'-O5'-P-OP1
2	C	1144	UMP	C5'-O5'-P-OP1


There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1147	GOL	1	3
4	B	1148	GOL	5	0
4	C	1146	GOL	1	0
4	C	1147	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	D	2
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	124:LEU	C	125:LYS	N	1.19
1	D	27:GLU	C	28:ARG	N	1.08
1	B	126:PRO	C	127:GLU	N	0.97

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	137/154 (88%)	-0.53	5 (3%)	46	45	12, 20, 44, 66	0
1	B	131/154 (85%)	-0.43	7 (5%)	32	31	9, 19, 53, 84	1 (0%)
1	C	137/154 (88%)	-0.49	2 (1%)	72	71	10, 21, 51, 58	0
1	D	130/154 (84%)	-0.50	6 (4%)	37	36	9, 20, 53, 73	1 (0%)
All	All	535/616 (86%)	-0.49	20 (3%)	45	44	9, 20, 52, 84	2 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	145	PRO	5.1
1	B	115	GLY	4.3
1	A	144	ALA	4.1
1	C	143	HIS	3.8
1	B	124	LEU	3.6
1	D	144	ALA	3.5
1	B	7	PRO	3.5
1	D	114	SER	3.5
1	A	7	PRO	3.1
1	B	127	GLU	3.0
1	D	124	LEU	2.9
1	D	115	GLY	2.8
1	B	144	ALA	2.6
1	B	143	HIS	2.6
1	A	143	HIS	2.6
1	C	24	ALA	2.6
1	A	23	ALA	2.3
1	D	127	GLU	2.1
1	A	25	THR	2.1
1	D	7	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	C	1147	6/6	0.74	0.20	51,53,56,56	0
4	GOL	C	1146	6/6	0.75	0.18	54,55,57,57	0
4	GOL	A	1147	6/6	0.80	0.24	24,46,52,52	0
4	GOL	B	1148	6/6	0.83	0.23	22,30,35,35	0
2	UMP	A	1145	20/20	0.98	0.04	8,12,18,20	0
2	UMP	C	1144	20/20	0.98	0.05	8,12,15,17	0
3	MN	D	1146	1/1	0.99	0.02	15,15,15,15	0
3	MN	A	1146	1/1	0.99	0.02	30,30,30,30	0
3	MN	B	1147	1/1	0.99	0.02	23,23,23,23	0
3	MN	C	1145	1/1	0.99	0.02	25,25,25,25	0
3	MN	D	1145	1/1	0.99	0.02	20,20,20,20	0
3	MN	B	1146	1/1	1.00	0.01	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around UMP A 1145:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UMP C 1144:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.