



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:32 PM UTC

PDB ID : 2YGQ / pdb_00002ygq
Title : WIF domain-epidermal growth factor (EGF)-like domains 1-3 of human Wnt inhibitory factor 1 in complex with 1,2- dipalmitoylphosphatidylcholine
Authors : Malinauskas, T.; Aricescu, A.R.; Lu, W.; Siebold, C.; Jones, E.Y.
Deposited on : 2011-04-19
Resolution : 3.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1046 (4.16-3.76)
Clashscore	190562	1019 (4.14-3.78)
Ramachandran outliers	187476	1031 (4.16-3.76)
Sidechain outliers	187428	1024 (4.16-3.76)
RSRZ outliers	180081	1046 (4.16-3.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	29% 61% 12% • 26%
2	B	2	100%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 1969 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called WNT INHIBITORY FACTOR 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	241	1826	1153	312	335	26	0	0	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	GLU	-	expression tag	UNP Q9Y5W5
A	33	THR	-	expression tag	UNP Q9Y5W5
A	34	GLY	-	expression tag	UNP Q9Y5W5
A	166	LYS	GLN	variant	UNP Q9Y5W5
A	347	GLY	-	expression tag	UNP Q9Y5W5
A	348	THR	-	expression tag	UNP Q9Y5W5
A	349	LYS	-	expression tag	UNP Q9Y5W5
A	350	HIS	-	expression tag	UNP Q9Y5W5
A	351	HIS	-	expression tag	UNP Q9Y5W5
A	352	HIS	-	expression tag	UNP Q9Y5W5
A	353	HIS	-	expression tag	UNP Q9Y5W5
A	354	HIS	-	expression tag	UNP Q9Y5W5
A	355	HIS	-	expression tag	UNP Q9Y5W5


- Molecule 2 is an oligosaccharide called 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	O	S			
2	B	2	55	12	35	8	0	0	0


- Molecule 3 is 1,2-DIACYL-SN-GLYCERO-3-PHOSHOCHOLINE (CCD ID: PCF) (formula:

C₄₀H₈₀NO₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			50	40	1	8	1		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O		0	0
			14	8	1	5			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is alpha-L-fucopyranose (CCD ID: FUC) (formula: $C_6H_{12}O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			10	6	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: WNT INHIBITORY FACTOR 1



• Molecule 2: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	178.01Å 178.01Å 178.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.80 – 3.95 39.80 – 3.95	Depositor EDS
% Data completeness (in resolution range)	96.0 (39.80-3.95) 99.8 (39.80-3.95)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.51 (at 4.00Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.7_650)	Depositor
R, R_{free}	0.297 , 0.331 0.299 , 0.340	Depositor DCC
R_{free} test set	399 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	147.7	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 175.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.032 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	1969	wwPDB-VP
Average B, all atoms (Å ²)	230.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, PCF, NAG, GU4, YYJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	1/1874 (0.1%)	1.14	3/2541 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	78	PRO	N-CD	12.47	1.65	1.47

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	200	CYS	CA-C-N	6.16	126.61	120.52
1	A	200	CYS	C-N-CA	6.16	126.61	120.52
1	A	92	GLN	N-CA-C	5.16	116.25	108.46

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1826	0	1735	55	0
2	B	55	0	6	0	0
3	A	50	0	80	18	0
4	A	28	0	26	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	10	0	10	0	0
All	All	1969	0	1857	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:VAL:CG1	3:A:1275:PCF:H411	2.07	0.84
1:A:90:THR:HG22	1:A:133:VAL:HG22	1.59	0.84
1:A:152:VAL:HG11	3:A:1275:PCF:H411	1.61	0.83
1:A:152:VAL:CB	3:A:1275:PCF:H411	2.13	0.79
1:A:152:VAL:HG11	3:A:1275:PCF:H422	1.65	0.78
1:A:215:THR:HB	1:A:216:PRO:HD3	1.70	0.72
1:A:77:MET:HE1	3:A:1275:PCF:H282	1.72	0.70
1:A:225:VAL:HG22	1:A:228:GLY:O	1.91	0.70
1:A:152:VAL:HG11	3:A:1275:PCF:C41	2.22	0.69
1:A:152:VAL:HB	3:A:1275:PCF:H411	1.73	0.69
3:A:1275:PCF:O13	3:A:1275:PCF:H143	1.91	0.69
1:A:152:VAL:HG11	3:A:1275:PCF:C42	2.22	0.68
1:A:216:PRO:HB2	1:A:240:ASN:OD1	1.94	0.67
1:A:247:SER:H	1:A:259:PRO:HB3	1.59	0.67
1:A:49:ILE:HG12	1:A:51:PHE:HD1	1.60	0.66
1:A:80:ILE:HD11	3:A:1275:PCF:H351	1.77	0.66
1:A:214:CYS:SG	1:A:229:PHE:O	2.56	0.64
1:A:63:MET:HB3	1:A:67:THR:OG1	1.97	0.64
1:A:215:THR:CB	1:A:216:PRO:HD3	2.29	0.63
1:A:90:THR:CG2	1:A:133:VAL:HG22	2.27	0.62
1:A:216:PRO:CB	1:A:240:ASN:OD1	2.48	0.62
1:A:49:ILE:O	3:A:1275:PCF:H142	2.00	0.61
1:A:90:THR:HG22	1:A:133:VAL:CG2	2.31	0.60
1:A:48:LEU:HD22	1:A:87:MET:HE3	1.84	0.58
1:A:49:ILE:HG12	1:A:51:PHE:CD1	2.38	0.58
1:A:214:CYS:SG	1:A:229:PHE:N	2.77	0.57
1:A:56:LEU:N	1:A:56:LEU:HD12	2.21	0.56
1:A:92:GLN:OE1	1:A:131:ALA:N	2.39	0.56
1:A:152:VAL:HG21	3:A:1275:PCF:H392	1.89	0.55
1:A:215:THR:HB	1:A:216:PRO:CD	2.38	0.54
1:A:45:ALA:O	1:A:49:ILE:HG22	2.07	0.54
1:A:216:PRO:HB3	1:A:240:ASN:HD21	1.74	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:216:PRO:CB	1:A:240:ASN:HD21	2.24	0.51
1:A:225:VAL:CG2	1:A:228:GLY:O	2.58	0.51
1:A:248:THR:HG23	1:A:248:THR:O	2.12	0.49
1:A:92:GLN:OE1	1:A:131:ALA:HB2	2.11	0.49
1:A:152:VAL:CG1	3:A:1275:PCF:C41	2.82	0.49
1:A:56:LEU:N	1:A:56:LEU:CD1	2.76	0.49
1:A:226:THR:HB	1:A:227:PRO:HD3	1.95	0.48
1:A:77:MET:CE	3:A:1275:PCF:H282	2.40	0.47
1:A:80:ILE:HD11	3:A:1275:PCF:C35	2.44	0.47
1:A:100:PHE:HD1	1:A:126:THR:HG22	1.80	0.47
1:A:92:GLN:OE1	1:A:131:ALA:CA	2.64	0.45
1:A:41:ASP:OD1	1:A:44:GLN:HB3	2.18	0.44
1:A:152:VAL:CG1	3:A:1275:PCF:H422	2.42	0.44
1:A:215:THR:CB	1:A:216:PRO:CD	2.96	0.44
1:A:49:ILE:HG23	1:A:51:PHE:H	1.83	0.44
1:A:169:GLN:HG2	1:A:170:ASN:OD1	2.19	0.43
1:A:216:PRO:HB2	1:A:240:ASN:CG	2.45	0.42
1:A:78:PRO:HD2	3:A:1275:PCF:H32	2.01	0.42
1:A:87:MET:SD	3:A:1275:PCF:H361	2.61	0.41
1:A:123:LEU:HD22	1:A:123:LEU:H	1.86	0.41
1:A:100:PHE:CD1	1:A:126:THR:HG22	2.56	0.40
1:A:122:PRO:HG3	1:A:220:ASN:HB3	2.03	0.40
1:A:216:PRO:CB	1:A:240:ASN:ND2	2.84	0.40
1:A:39:TRP:CE2	1:A:90:THR:OG1	2.71	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/324 (74%)	218 (91%)	21 (9%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	199/269 (74%)	194 (98%)	5 (2%)	42 62

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	LEU
1	A	48	LEU
1	A	167	THR
1	A	218	CYS
1	A	263	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	HIS
1	A	144	GLN
1	A	205	HIS
1	A	220	ASN
1	A	252	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GU4	B	1	2	27,27,28	0.91	1 (3%)	32,43,45	1.81	7 (21%)
2	YYJ	B	2	2	27,28,28	0.93	0	33,46,46	1.94	6 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GU4	B	1	2	-	0/21/38/41	0/1/1/1
2	YYJ	B	2	2	-	3/23/42/42	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	GU4	O2-C2	-2.40	1.43	1.47

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	YYJ	C3-O3-S3	-7.15	108.85	117.69
2	B	2	YYJ	C4-O4-S4	-4.32	108.68	119.04
2	B	1	GU4	C3-O3-S3	-4.16	109.05	119.04
2	B	1	GU4	C4-O4-S4	-3.93	109.61	119.04
2	B	2	YYJ	O3S1-S1-O2S1	3.36	125.21	112.24
2	B	2	YYJ	O3S6-S6-O2S6	3.24	124.74	112.24
2	B	1	GU4	O26-S4-O25	3.23	124.68	112.24
2	B	2	YYJ	O3S4-S4-O2S4	3.22	124.66	112.24
2	B	1	GU4	O12-S2-O11	3.22	124.65	112.24
2	B	1	GU4	O27-S3-O28	3.21	124.62	112.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	YYJ	O3S3-S3-O2S3	3.19	124.55	112.24
2	B	1	GU4	O2-C2-C3	3.16	110.15	106.65
2	B	1	GU4	O23-S6-O22	3.15	124.38	112.24

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2	YYJ	O1-C1-C2-C3
2	B	2	YYJ	O1-C1-C2-O2
2	B	2	YYJ	O1-C1-C2-O5

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	FUC	A	1278	1	10,10,11	0.46	0	14,14,16	0.96	1 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	1277	1	14,14,15	0.49	0	17,19,21	1.11	2 (11%)
3	PCF	A	1275	-	49,49,49	0.93	2 (4%)	55,57,57	1.20	8 (14%)
4	NAG	A	1276	1	14,14,15	0.40	0	17,19,21	1.18	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	FUC	A	1278	1	-	-	0/1/1/1
4	NAG	A	1277	1	-	0/6/23/26	0/1/1/1
3	PCF	A	1275	-	-	26/53/53/53	-
4	NAG	A	1276	1	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1275	PCF	O21-C21	3.98	1.45	1.34
3	A	1275	PCF	O31-C31	3.68	1.44	1.33

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1275	PCF	O21-C21-C22	4.39	120.98	111.48
3	A	1275	PCF	C3-C2-C1	-3.15	104.45	111.78
3	A	1275	PCF	C2-O21-C21	-2.55	111.70	117.80
4	A	1276	NAG	C4-C3-C2	-2.54	107.29	111.02
5	A	1278	FUC	C6-C5-C4	-2.34	108.79	113.08
3	A	1275	PCF	O31-C31-C32	2.30	118.86	111.83
4	A	1277	NAG	C8-C7-N2	2.27	119.89	116.12
4	A	1276	NAG	C8-C7-N2	2.18	119.73	116.12
3	A	1275	PCF	O31-C31-O32	-2.17	118.19	123.63
4	A	1277	NAG	C4-C3-C2	-2.17	107.84	111.02
3	A	1275	PCF	C11-C12-N	-2.10	109.08	115.82
3	A	1275	PCF	C33-C32-C31	-2.10	106.01	113.69
3	A	1275	PCF	O21-C21-O22	-2.08	118.85	123.70

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1275	PCF	C11-C12-N-C14
3	A	1275	PCF	C11-C12-N-C15
3	A	1275	PCF	C22-C21-O21-C2
3	A	1275	PCF	O22-C21-O21-C2
3	A	1275	PCF	C32-C31-O31-C3
3	A	1275	PCF	O32-C31-O31-C3
3	A	1275	PCF	C11-C12-N-C13
3	A	1275	PCF	C41-C42-C43-C44
3	A	1275	PCF	C33-C34-C35-C36
3	A	1275	PCF	C30-C47-C48-C49
3	A	1275	PCF	C47-C48-C49-C50
3	A	1275	PCF	C40-C41-C42-C43
3	A	1275	PCF	O21-C2-C3-O31
3	A	1275	PCF	C26-C27-C28-C29
3	A	1275	PCF	C34-C35-C36-C37
3	A	1275	PCF	C23-C24-C25-C26
3	A	1275	PCF	C25-C26-C27-C28
3	A	1275	PCF	C27-C28-C29-C30
3	A	1275	PCF	C24-C25-C26-C27
3	A	1275	PCF	O11-C1-C2-O21
3	A	1275	PCF	O11-C1-C2-C3
3	A	1275	PCF	C32-C33-C34-C35
3	A	1275	PCF	C2-C1-O11-P
3	A	1275	PCF	O21-C21-C22-C23
3	A	1275	PCF	C1-C2-C3-O31
3	A	1275	PCF	C22-C23-C24-C25


There are no ring outliers.

1 monomer is involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1275	PCF	18	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	241/324 (74%)	1.96	95 (39%) 1 1	104, 211, 329, 466	0

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	96	GLN	9.4
1	A	202	ASP	8.9
1	A	258	TYR	7.6
1	A	188	ASN	7.0
1	A	152	VAL	6.5
1	A	76	ARG	6.4
1	A	169	GLN	6.2
1	A	94	ALA	5.9
1	A	43	HIS	5.7
1	A	266	PRO	5.1
1	A	153	ASP	5.1
1	A	34	GLY	4.9
1	A	196	ARG	4.7
1	A	75	GLN	4.6
1	A	72	LYS	4.6
1	A	154	VAL	4.5
1	A	217	ARG	4.3
1	A	227	PRO	4.3
1	A	147	VAL	4.3
1	A	148	ALA	4.3
1	A	267	GLY	4.2
1	A	103	PHE	4.2
1	A	183	PRO	4.2
1	A	215	THR	4.1
1	A	189	GLY	4.1
1	A	168	PRO	4.1
1	A	167	THR	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	79	ALA	4.0
1	A	191	PHE	3.8
1	A	170	ASN	3.7
1	A	260	GLY	3.6
1	A	181	GLU	3.6
1	A	166	LYS	3.6
1	A	102	GLU	3.5
1	A	145	ASP	3.5
1	A	164	ILE	3.4
1	A	77	MET	3.4
1	A	210	GLU	3.4
1	A	143	LYS	3.4
1	A	165	LEU	3.4
1	A	199	GLU	3.4
1	A	60	GLU	3.3
1	A	100	PHE	3.3
1	A	114	MET	3.3
1	A	265	PRO	3.3
1	A	99	TYR	3.3
1	A	95	GLY	3.2
1	A	171	ALA	3.2
1	A	48	LEU	3.2
1	A	106	LEU	3.2
1	A	226	THR	3.1
1	A	149	ALA	3.1
1	A	113	ILE	3.1
1	A	178	GLN	3.0
1	A	192	CYS	3.0
1	A	221	GLY	3.0
1	A	51	PHE	2.9
1	A	257	PHE	2.9
1	A	163	THR	2.9
1	A	177	CYS	2.9
1	A	261	LYS	2.9
1	A	219	MET	2.9
1	A	87	MET	2.9
1	A	249	THR	2.8
1	A	98	GLU	2.8
1	A	69	ASP	2.8
1	A	229	PHE	2.7
1	A	150	PHE	2.7
1	A	225	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	259	PRO	2.6
1	A	53	GLU	2.6
1	A	224	CYS	2.6
1	A	236	PHE	2.5
1	A	243	LYS	2.5
1	A	155	ILE	2.4
1	A	195	ARG	2.4
1	A	190	GLY	2.4
1	A	207	PRO	2.4
1	A	247	SER	2.4
1	A	160	GLU	2.4
1	A	174	PHE	2.3
1	A	272	GLN	2.3
1	A	187	ARG	2.3
1	A	138	PHE	2.3
1	A	184	GLY	2.3
1	A	56	LEU	2.2
1	A	126	THR	2.2
1	A	239	VAL	2.2
1	A	111	LYS	2.2
1	A	175	LYS	2.1
1	A	144	GLN	2.1
1	A	151	GLU	2.1
1	A	36	LEU	2.1
1	A	97	ALA	2.1
1	A	204	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	YYJ	B	2	28/28	0.47	0.17	518,546,585,594	0
2	GU4	B	1	27/28	0.63	0.14	492,519,553,566	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	FUC	A	1278	10/11	0.60	0.26	287,299,315,315	0
4	NAG	A	1277	14/15	0.80	0.35	311,339,357,372	0
3	PCF	A	1275	50/50	0.88	0.32	102,185,344,362	0
4	NAG	A	1276	14/15	0.90	0.33	253,263,276,279	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.