



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 12:30 PM EDT

PDB ID : 2YHV / pdb_00002yhv
Title : Structure of L1196M Mutant Anaplastic Lymphoma Kinase
Authors : McTigue, M.; Deng, Y.; Liu, W.; Brooun, A.
Deposited on : 2011-05-06
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2493 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALK TYROSINE KINASE RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	292	Total	C	N	O	S	0	0	1
			2296	1464	391	419	22			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1070	MET	-	expression tag	UNP Q9UM73
A	1071	ALA	-	expression tag	UNP Q9UM73
A	1072	HIS	-	expression tag	UNP Q9UM73
A	1073	HIS	-	expression tag	UNP Q9UM73
A	1074	HIS	-	expression tag	UNP Q9UM73
A	1075	HIS	-	expression tag	UNP Q9UM73
A	1076	HIS	-	expression tag	UNP Q9UM73
A	1077	HIS	-	expression tag	UNP Q9UM73
A	1078	ASP	-	expression tag	UNP Q9UM73
A	1079	TYR	-	expression tag	UNP Q9UM73
A	1080	GLY	-	expression tag	UNP Q9UM73
A	1081	ILE	-	expression tag	UNP Q9UM73
A	1082	PRO	-	expression tag	UNP Q9UM73
A	1083	THR	-	expression tag	UNP Q9UM73
A	1084	THR	-	expression tag	UNP Q9UM73
A	1085	GLU	-	expression tag	UNP Q9UM73
A	1086	ASN	-	expression tag	UNP Q9UM73
A	1087	LEU	-	expression tag	UNP Q9UM73
A	1088	TYR	-	expression tag	UNP Q9UM73
A	1089	PHE	-	expression tag	UNP Q9UM73
A	1090	GLN	-	expression tag	UNP Q9UM73
A	1091	GLY	-	expression tag	UNP Q9UM73
A	1092	SER	-	expression tag	UNP Q9UM73
A	1196	MET	LEU	engineered mutation	UNP Q9UM73


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	197	Total 197	O 197	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ALK TYROSINE KINASE RECEPTOR



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.10Å 57.76Å 105.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.60 – 1.90 52.59 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (52.60-1.90) 99.7 (52.59-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.87 (at 1.90Å)	Xtriage
Refinement program	CNS 2005	Depositor
R, R_{free}	0.206 , 0.238 0.203 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	27.3	Xtriage
Anisotropy	0.361	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2493	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	1/2350 (0.0%)	0.66	2/3183 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1400	GLU	C-N	-5.74	1.25	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1261	GLY	CA-C-N	6.41	126.91	119.47
1	A	1261	GLY	C-N-CA	6.41	126.91	119.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2296	0	2279	59	0
2	A	197	0	0	4	0
All	All	2493	0	2279	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1101:LYS:HE3	1:A:1102:THR:H	1.17	1.07
1:A:1105:ILE:HD12	1:A:1165:LEU:HD11	1.51	0.92
1:A:1197:GLU:HG2	1:A:1199:MET:HE3	1.55	0.88
1:A:1147:VAL:HG12	1:A:1183:ILE:HD13	1.59	0.85
1:A:1101:LYS:HE3	1:A:1102:THR:N	1.92	0.85
1:A:1105:ILE:N	1:A:1105:ILE:HD13	1.95	0.82
1:A:1199:MET:HA	1:A:1199:MET:HE2	1.65	0.78
1:A:1165:LEU:O	1:A:1169:LEU:HD13	1.89	0.73
1:A:1279:ARG:HA	1:A:1279:ARG:HE	1.54	0.71
1:A:1093:ASN:HB2	1:A:1105:ILE:HG12	1.75	0.68
1:A:1387:THR:HG21	2:A:2094:HOH:O	1.93	0.68
1:A:1279:ARG:HA	1:A:1279:ARG:NE	2.08	0.67
1:A:1383:ILE:O	1:A:1387:THR:HG23	1.97	0.64
1:A:1248:ARG:CZ	1:A:1277:ILE:HG21	2.27	0.64
1:A:1211:THR:HG23	1:A:1221:LEU:HD22	1.80	0.64
1:A:1105:ILE:CD1	1:A:1165:LEU:HD21	2.30	0.61
1:A:1365:CYS:O	1:A:1373:ARG:HD2	2.01	0.60
1:A:1367:GLN:HB2	1:A:1373:ARG:HG2	1.82	0.60
1:A:1105:ILE:HD13	1:A:1105:ILE:H	1.64	0.60
1:A:1214:ARG:HG3	1:A:1215:PRO:HD2	1.83	0.59
1:A:1382:ARG:HD3	2:A:2176:HOH:O	2.03	0.59
1:A:1180:VAL:HG13	1:A:1197:GLU:HB3	1.85	0.58
1:A:1155:VAL:O	1:A:1155:VAL:HG23	2.03	0.58
1:A:1171:ILE:HG12	1:A:1196:MET:HE1	1.86	0.58
1:A:1105:ILE:N	1:A:1105:ILE:CD1	2.66	0.56
1:A:1247:HIS:O	1:A:1248:ARG:HB2	2.07	0.55
1:A:1147:VAL:CG1	1:A:1183:ILE:HD13	2.35	0.54
1:A:1105:ILE:HD11	1:A:1165:LEU:HD21	1.90	0.53
1:A:1108:LEU:HD11	1:A:1169:LEU:CD1	2.39	0.52
1:A:1214:ARG:HE	1:A:1215:PRO:HD2	1.73	0.52
1:A:1298:PRO:HA	1:A:1301:PHE:CZ	2.44	0.52
1:A:1214:ARG:NE	1:A:1400:GLU:O	2.43	0.51
1:A:1214:ARG:NE	1:A:1215:PRO:HD2	2.28	0.48
1:A:1278:TYR:O	1:A:1279:ARG:HB2	2.13	0.48
1:A:1097:CYS:HB3	2:A:2027:HOH:O	2.13	0.48
1:A:1214:ARG:HE	1:A:1215:PRO:CD	2.26	0.48
1:A:1208:LEU:O	1:A:1212:ARG:HG3	2.14	0.48
1:A:1367:GLN:O	1:A:1373:ARG:HD3	2.13	0.47
1:A:1325:LEU:N	1:A:1325:LEU:HD22	2.30	0.47
1:A:1108:LEU:HD11	1:A:1169:LEU:HD12	1.97	0.46
1:A:1147:VAL:HG12	1:A:1183:ILE:CD1	2.39	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1199:MET:HE1	2:A:2075:HOH:O	2.15	0.46
1:A:1105:ILE:HD12	1:A:1165:LEU:CD1	2.36	0.46
1:A:1369:GLN:HB3	1:A:1371:GLU:OE2	2.17	0.45
1:A:1198:LEU:O	1:A:1199:MET:HE2	2.16	0.45
1:A:1114:LYS:HG3	1:A:1115:ASN:ND2	2.32	0.44
1:A:1331:PRO:O	1:A:1332:SER:OG	2.34	0.44
1:A:1371:GLU:CD	1:A:1371:GLU:H	2.25	0.43
1:A:1217:GLN:HB3	1:A:1220:SER:HB3	2.01	0.43
1:A:1328:MET:SD	1:A:1329:PRO:HD2	2.59	0.42
1:A:1171:ILE:HD11	1:A:1180:VAL:O	2.20	0.42
1:A:1108:LEU:HD21	1:A:1169:LEU:HD12	2.02	0.41
1:A:1382:ARG:HG2	1:A:1382:ARG:HH11	1.86	0.41
1:A:1399:ILE:O	1:A:1401:TYR:N	2.54	0.41
1:A:1309:LYS:HA	1:A:1309:LYS:HD3	1.88	0.40
1:A:1275:ARG:C	1:A:1275:ARG:HD2	2.45	0.40
1:A:1328:MET:HA	1:A:1329:PRO:HD3	1.98	0.40
1:A:1115:ASN:N	1:A:1115:ASN:HD22	2.19	0.40
1:A:1199:MET:HE1	1:A:1258:THR:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/342 (83%)	279 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	255/298 (86%)	249 (98%)	6 (2%)	43 38

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1101	LYS
1	A	1105	ILE
1	A	1118	LEU
1	A	1214	ARG
1	A	1221	LEU
1	A	1360	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1093	ASN
1	A	1115	ASN
1	A	1188	GLN
1	A	1217	GLN
1	A	1238	GLN
1	A	1243	ASN
1	A	1244	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	292/342 (85%)	0.75	41 (14%) 6 6	16, 28, 53, 70	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1127	PHE	5.5
1	A	1126	ALA	5.4
1	A	1286	GLY	5.3
1	A	1401	TYR	4.6
1	A	1144	PRO	4.4
1	A	1101	LYS	4.1
1	A	1100	GLY	4.0
1	A	1302	MET	3.8
1	A	1277	ILE	3.7
1	A	1122	LEU	3.6
1	A	1105	ILE	3.6
1	A	1279	ARG	3.5
1	A	1301	PHE	3.5
1	A	1097	CYS	3.3
1	A	1278	TYR	3.2
1	A	1099	ALA	3.1
1	A	1093	ASN	3.1
1	A	1235	CYS	3.0
1	A	1217	GLN	3.0
1	A	1215	PRO	2.8
1	A	1214	ARG	2.8
1	A	1094	PRO	2.8
1	A	1287	GLY	2.7
1	A	1162	LEU	2.7
1	A	1216	SER	2.7
1	A	1400	GLU	2.7
1	A	1211	THR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1221	LEU	2.5
1	A	1288	CYS	2.4
1	A	1155	VAL	2.4
1	A	1106	SER	2.4
1	A	1128	GLY	2.4
1	A	1308	SER	2.3
1	A	1399	ILE	2.3
1	A	1218	PRO	2.2
1	A	1171	ILE	2.2
1	A	1095	ASN	2.2
1	A	1273	MET	2.2
1	A	1303	GLU	2.2
1	A	1219	SER	2.1
1	A	1369	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.