



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:36 PM UTC

PDB ID : 4YHJ / pdb_00004yhj
Title : Structure and Function of the Hypertension Variant A486V of G Protein-coupled Receptor Kinase 4 (GRK4)
Authors : Allen, S.J.; Parthasarathy, G.; Soisson, S.; Munshi, S.
Deposited on : 2015-02-27
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	584	
1	B	584	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8222 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called G protein-coupled receptor kinase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	501	Total	C	N	O	S	0	0	0
			4067	2580	708	752	27			
1	B	490	Total	C	N	O	S	0	0	0
			3978	2521	696	734	27			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	563	SER	CYS	engineered mutation	UNP P32298
A	578	SER	CYS	engineered mutation	UNP P32298
A	579	GLU	-	expression tag	UNP P32298
A	580	ASN	-	expression tag	UNP P32298
A	581	LEU	-	expression tag	UNP P32298
A	582	TYR	-	expression tag	UNP P32298
A	583	PHE	-	expression tag	UNP P32298
A	584	GLN	-	expression tag	UNP P32298
B	563	SER	CYS	engineered mutation	UNP P32298
B	578	SER	CYS	engineered mutation	UNP P32298
B	579	GLU	-	expression tag	UNP P32298
B	580	ASN	-	expression tag	UNP P32298
B	581	LEU	-	expression tag	UNP P32298
B	582	TYR	-	expression tag	UNP P32298
B	583	PHE	-	expression tag	UNP P32298
B	584	GLN	-	expression tag	UNP P32298

- Molecule 2 is AMP PHOSPHORAMIDATE (CCD ID: AN2) (formula: $C_{10}H_{16}N_6O_9P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	6	9	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	6	9	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	62	Total	O	0	0
			62	62		
3	B	61	Total	O	0	0
			61	61		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: G protein-coupled receptor kinase 4



• Molecule 1: G protein-coupled receptor kinase 4



LEU	ASP	LEU	ASP	LYS	ASN	ILE	HIS	THR	PRO	VAL	SER	ARG	PRO	ASN	ARG	GLY	PHE	PHE	TYR	ARG	LEU	PHE	ARG	ARG	GLY	GLY	SER	LEU	THR	MET	VAL	PRO	SER	GLU	GLY	LYS	GLU	VAL	GLU	PRO	LYS	GLN	SER	GLU	ASN	LEU	TYR	PHE	GLN
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.84Å 104.84Å 221.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.00 – 2.60 47.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.5 (47.00-2.60) 99.5 (47.00-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 2.48Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.11.5	Depositor
R, R_{free}	0.197 , 0.248 0.200 , 0.253	Depositor DCC
R_{free} test set	2523 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	46.6	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 36.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8222	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AN2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.85	2/4158 (0.0%)	1.43	22/5599 (0.4%)
1	B	0.84	0/4067	1.42	26/5474 (0.5%)
All	All	0.84	2/8225 (0.0%)	1.42	48/11073 (0.4%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	510	ILE	CA-CB	7.76	1.58	1.54
1	A	481	ILE	CA-CB	5.42	1.61	1.54

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	477	ASP	CA-C-N	9.04	137.97	121.70
1	A	477	ASP	C-N-CA	9.04	137.97	121.70
1	A	482	GLU	CA-C-N	7.56	135.98	121.54
1	A	482	GLU	C-N-CA	7.56	135.98	121.54
1	B	412	LYS	N-CA-C	-6.85	103.60	112.23
1	A	28	LYS	N-CA-C	-6.49	104.14	111.14
1	A	321	ASP	CA-CB-CG	6.22	118.82	112.60
1	B	284	PHE	CA-CB-CG	6.15	119.95	113.80
1	B	321	ASP	CA-CB-CG	6.12	118.72	112.60
1	A	483	GLN	CA-C-N	6.00	130.25	121.31
1	A	483	GLN	C-N-CA	6.00	130.25	121.31
1	A	482	GLU	CB-CA-C	-5.92	108.08	115.89
1	B	405	ASP	CA-CB-CG	5.80	118.40	112.60
1	B	309	VAL	N-CA-CB	5.75	117.94	111.21
1	A	105	ILE	CA-C-N	5.70	127.92	120.28
1	A	105	ILE	C-N-CA	5.70	127.92	120.28
1	B	268	ASN	CA-CB-CG	5.69	118.29	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	395	TRP	CA-C-N	5.67	128.66	120.38
1	B	395	TRP	C-N-CA	5.67	128.66	120.38
1	B	259	ALA	N-CA-C	5.52	117.37	109.14
1	B	296	LEU	CA-C-N	5.52	127.68	120.28
1	B	296	LEU	C-N-CA	5.52	127.68	120.28
1	B	280	GLY	CA-C-N	5.50	126.97	120.09
1	B	280	GLY	C-N-CA	5.50	126.97	120.09
1	B	186	THR	N-CA-C	-5.43	106.16	112.89
1	A	97	ASP	CA-CB-CG	5.35	117.95	112.60
1	A	370	TRP	CA-C-N	5.31	125.88	119.98
1	A	370	TRP	C-N-CA	5.31	125.88	119.98
1	A	166	SER	CA-C-N	5.25	127.32	120.28
1	A	166	SER	C-N-CA	5.25	127.32	120.28
1	A	177	LEU	CA-C-N	5.24	127.83	120.28
1	A	177	LEU	C-N-CA	5.24	127.83	120.28
1	A	285	ASP	CA-C-N	5.24	127.56	120.38
1	A	285	ASP	C-N-CA	5.24	127.56	120.38
1	B	56	LEU	N-CA-C	5.23	117.89	111.82
1	B	101	CYS	CA-C-N	5.23	125.74	119.94
1	B	101	CYS	C-N-CA	5.23	125.74	119.94
1	A	482	GLU	N-CA-C	5.21	117.78	109.02
1	B	160	GLU	CA-C-N	5.20	127.25	120.28
1	B	160	GLU	C-N-CA	5.20	127.25	120.28
1	B	145	GLU	CA-C-N	5.17	127.16	120.44
1	B	145	GLU	C-N-CA	5.17	127.16	120.44
1	B	134	LYS	N-CA-C	-5.15	105.35	111.69
1	A	160	GLU	CB-CG-CD	5.10	121.27	112.60
1	B	328	ILE	N-CA-C	-5.08	103.03	109.58
1	B	471	HIS	CA-C-N	5.07	127.97	120.82
1	B	471	HIS	C-N-CA	5.07	127.97	120.82
1	B	452	ILE	N-CA-C	5.02	115.81	108.58

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4067	0	4025	36	0
1	B	3978	0	3932	30	0
2	A	27	0	14	0	0
2	B	27	0	14	0	0
3	A	62	0	0	2	0
3	B	61	0	0	0	0
All	All	8222	0	7985	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:GLN:HE21	1:A:513:GLN:HE22	1.24	0.84
1:B:60:GLN:HE21	1:B:513:GLN:HE22	1.25	0.79
1:B:344:ARG:HH21	1:B:360:ASN:HD21	1.36	0.73
3:A:708:HOH:O	1:B:155:ARG:HD2	1.91	0.69
1:B:204:CYS:SG	1:B:215:CYS:HB2	2.32	0.69
1:A:487:VAL:HA	1:A:490:ILE:HD12	1.77	0.66
1:A:344:ARG:HD2	1:A:360:ASN:HD21	1.60	0.65
1:A:388:LYS:HE3	1:A:388:LYS:H	1.61	0.65
1:A:345:GLY:H	1:A:360:ASN:HD22	1.47	0.63
1:A:211:LYS:HD2	1:A:515:GLU:HG3	1.84	0.59
1:B:43:SER:O	1:B:47:HIS:HD2	1.86	0.58
1:B:253:ALA:O	1:B:506:GLY:HA2	2.03	0.58
1:A:256:THR:HG22	1:A:258:ASP:H	1.68	0.58
1:A:345:GLY:H	1:A:360:ASN:ND2	2.02	0.57
1:A:446:HIS:ND1	1:A:448:VAL:HG22	2.20	0.57
1:A:356:GLU:HG2	1:A:429:PRO:HG3	1.90	0.54
1:B:79:ARG:HD3	1:B:117:PRO:HB3	1.91	0.53
1:A:479:LEU:CB	1:A:481:ILE:HD12	2.40	0.52
1:B:522:PHE:O	1:B:526:ASN:HB3	2.09	0.52
1:A:204:CYS:SG	1:A:215:CYS:HB2	2.50	0.51
1:A:79:ARG:HD3	1:A:117:PRO:HB3	1.93	0.51
1:A:346:ARG:HG3	1:A:358:VAL:HA	1.92	0.51
1:A:252:TYR:HB2	1:A:263:VAL:HB	1.95	0.49
1:A:221:LYS:HD3	1:A:487:VAL:HB	1.95	0.48
1:A:356:GLU:CG	1:A:429:PRO:HG3	2.43	0.48
1:B:346:ARG:HG3	1:B:358:VAL:HA	1.96	0.48
1:A:479:LEU:HB3	1:A:481:ILE:HD12	1.96	0.47
1:B:311:ARG:HH21	1:B:335:THR:HG21	1.81	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:GLY:HA2	1:A:478:VAL:HG21	1.98	0.46
1:B:346:ARG:NH2	1:B:353:MET:HE1	2.29	0.46
1:B:438:GLU:HB2	1:B:442:GLY:HA3	1.98	0.45
1:A:393:VAL:HG21	1:A:401:ARG:HH12	1.80	0.45
1:B:198:PHE:HE1	1:B:226:ARG:HB2	1.81	0.45
1:A:250:LEU:HA	1:A:264:LEU:HD23	1.99	0.45
1:B:218:LEU:HB3	1:B:223:ILE:HD11	1.99	0.45
1:B:267:MET:HE2	1:B:327:ARG:HB2	1.99	0.45
1:B:356:GLU:HG2	1:B:429:PRO:HG3	1.98	0.45
1:B:453:ASN:CG	1:B:456:ARG:HG3	2.42	0.45
1:B:242:VAL:HG21	1:B:303:LEU:HD23	1.98	0.45
1:B:396:GLU:H	1:B:396:GLU:HG2	1.46	0.45
1:B:383:HIS:CD2	1:B:388:LYS:HA	2.52	0.44
1:A:107:ASP:HA	1:A:111:ASN:ND2	2.33	0.44
1:B:346:ARG:HH21	1:B:353:MET:HE1	1.83	0.44
1:B:433:LEU:HG	1:B:443:VAL:HG22	2.00	0.44
1:B:276:ILE:HD11	1:B:379:MET:HG2	1.99	0.44
1:B:353:MET:HE3	1:B:353:MET:HB2	1.80	0.44
1:A:267:MET:HE2	1:A:327:ARG:HB2	1.99	0.43
1:A:375:LEU:HD12	1:A:375:LEU:HA	1.86	0.43
1:A:482:GLU:HB3	1:A:483:GLN:H	1.29	0.43
1:A:151:HIS:O	1:A:155:ARG:HB2	2.19	0.43
1:A:477:ASP:HA	1:A:478:VAL:HB	2.01	0.42
1:B:234:ASN:O	1:B:238:ILE:HG12	2.19	0.42
1:B:449:PHE:HA	1:B:452:ILE:HD12	2.01	0.42
1:B:446:HIS:ND1	1:B:448:VAL:HG22	2.35	0.42
1:A:30:TRP:HA	1:A:33:ILE:HD12	2.02	0.42
1:A:311:ARG:HD2	3:A:734:HOH:O	2.19	0.42
1:A:479:LEU:HB2	1:A:481:ILE:HD12	2.02	0.41
1:A:428:ASN:HA	1:A:429:PRO:HD2	1.94	0.41
1:A:46:ARG:HB2	1:A:172:LEU:HD13	2.02	0.41
1:A:481:ILE:HG12	1:A:482:GLU:HG3	2.03	0.41
1:B:133:LEU:HD11	1:B:143:PHE:CZ	2.56	0.41
1:A:433:LEU:HG	1:A:443:VAL:HG22	2.02	0.41
1:B:90:TYR:CE1	1:B:98:ARG:HG2	2.56	0.41
1:B:355:PRO:HG2	1:B:427:LYS:HA	2.02	0.41
1:A:388:LYS:HZ1	1:A:401:ARG:HH22	1.68	0.40
1:A:122:PRO:HA	1:A:123:PRO:HD3	1.97	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	499/584 (85%)	472 (95%)	23 (5%)	4 (1%)	16	34
1	B	486/584 (83%)	466 (96%)	19 (4%)	1 (0%)	43	66
All	All	985/1168 (84%)	938 (95%)	42 (4%)	5 (0%)	24	46

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	483	GLN
1	A	478	VAL
1	A	137	ASN
1	B	391	GLU
1	A	481	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	441/515 (86%)	399 (90%)	42 (10%)	8	18
1	B	430/515 (84%)	395 (92%)	35 (8%)	11	24
All	All	871/1030 (85%)	794 (91%)	77 (9%)	9	21

All (77) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	51	LYS
1	A	59	LYS
1	A	65	ARG
1	A	103	LEU
1	A	106	LEU
1	A	113	LYS
1	A	114	LEU
1	A	131	LEU
1	A	135	GLU
1	A	157	GLU
1	A	191	ARG
1	A	217	LYS
1	A	219	GLN
1	A	222	ARG
1	A	233	LEU
1	A	236	LYS
1	A	237	ARG
1	A	238	ILE
1	A	249	SER
1	A	256	THR
1	A	257	LYS
1	A	279	LEU
1	A	285	ASP
1	A	286	GLU
1	A	296	LEU
1	A	311	ARG
1	A	313	LEU
1	A	322	ASP
1	A	326	ILE
1	A	375	LEU
1	A	388	LYS
1	A	396	GLU
1	A	397	GLU
1	A	406	THR
1	A	411	GLU
1	A	431	LYS
1	A	448	VAL
1	A	463	GLU
1	A	476	LYS
1	A	481	ILE
1	A	483	GLN
1	B	28	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	32	GLU
1	B	41	GLN
1	B	49	ILE
1	B	66	LEU
1	B	69	GLN
1	B	99	SER
1	B	103	LEU
1	B	106	LEU
1	B	113	LYS
1	B	136	GLU
1	B	147	THR
1	B	177	LEU
1	B	219	GLN
1	B	227	LYS
1	B	236	LYS
1	B	257	LYS
1	B	268	ASN
1	B	287	GLN
1	B	311	ARG
1	B	313	LEU
1	B	335	THR
1	B	356	GLU
1	B	393	VAL
1	B	396	GLU
1	B	408	GLU
1	B	411	GLU
1	B	415	GLU
1	B	438	GLU
1	B	451	ASP
1	B	455	ARG
1	B	456	ARG
1	B	475	CYS
1	B	498	ASP
1	B	507	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
1	A	47	HIS
1	A	60	GLN
1	A	69	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	163	GLN
1	A	304	GLN
1	A	360	ASN
1	B	47	HIS
1	B	60	GLN
1	B	111	ASN
1	B	189	HIS
1	B	278	ASN
1	B	304	GLN
1	B	360	ASN
1	B	383	HIS
1	B	514	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AN2	B	601	-	28,29,29	2.45	4 (14%)	40,45,45	2.16	11 (27%)
2	AN2	A	601	-	28,29,29	1.35	2 (7%)	40,45,45	2.20	9 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AN2	B	601	-	-	0/13/32/32	0/3/3/3
2	AN2	A	601	-	-	3/13/32/32	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	AN2	PB-O1B	10.75	1.62	1.46
2	B	601	AN2	PB-O3A	3.94	1.64	1.59
2	A	601	AN2	PA-O3A	3.63	1.63	1.59
2	B	601	AN2	PA-O3A	3.50	1.63	1.59
2	A	601	AN2	PB-O3A	3.35	1.63	1.59
2	B	601	AN2	PB-O2B	-3.15	1.48	1.56

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	AN2	C5-C4-N3	-6.24	118.13	126.72
2	A	601	AN2	C5-C4-N3	-5.78	118.77	126.72
2	A	601	AN2	PB-O3A-PA	-5.17	115.56	132.10
2	B	601	AN2	N3-C4-N9	4.84	135.40	127.17
2	A	601	AN2	N3-C4-N9	4.75	135.24	127.17
2	A	601	AN2	N1-C2-N3	-4.71	121.45	128.58
2	B	601	AN2	N1-C2-N3	-4.26	122.13	128.58
2	B	601	AN2	PB-O3A-PA	-4.22	118.60	132.10
2	B	601	AN2	C2-N3-C4	4.16	122.00	111.83
2	A	601	AN2	C5-N7-C8	4.10	109.90	103.45
2	A	601	AN2	C2-N3-C4	4.10	121.83	111.83
2	B	601	AN2	C5-N7-C8	3.88	109.55	103.45
2	B	601	AN2	C4-C5-N7	-3.54	106.54	110.58
2	A	601	AN2	C4-C5-N7	-3.31	106.80	110.58
2	A	601	AN2	O4'-C1'-N9	3.27	114.38	108.09
2	B	601	AN2	O4'-C1'-N9	2.88	113.61	108.09
2	A	601	AN2	N9-C8-N7	-2.84	109.91	113.94
2	B	601	AN2	N9-C8-N7	-2.51	110.37	113.94
2	B	601	AN2	O2A-PA-O3A	2.43	113.85	107.27
2	B	601	AN2	PA-O5'-C5'	-2.20	108.76	121.35

There are no chirality outliers.


All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	AN2	C5'-O5'-PA-O3A
2	A	601	AN2	C5'-O5'-PA-O2A
2	A	601	AN2	C5'-O5'-PA-O1A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	501/584 (85%)	0.01	15 (2%)	52	47	26, 45, 79, 109	0
1	B	490/584 (83%)	0.03	14 (2%)	53	48	27, 47, 83, 135	0
All	All	991/1168 (84%)	0.02	29 (2%)	53	48	26, 46, 81, 135	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	475	CYS	5.2
1	B	473	VAL	5.1
1	B	25	GLY	3.8
1	B	527	LYS	3.6
1	A	483	GLN	3.5
1	B	472	ALA	3.4
1	B	474	TYR	3.3
1	B	471	HIS	3.3
1	A	138	PRO	2.9
1	A	525	ILE	2.9
1	A	28	LYS	2.7
1	B	137	ASN	2.7
1	A	390	LYS	2.7
1	B	470	PRO	2.7
1	A	471	HIS	2.6
1	A	486	VAL	2.5
1	B	489	GLY	2.5
1	A	484	PHE	2.5
1	B	490	ILE	2.4
1	A	389	TYR	2.4
1	B	390	LYS	2.4
1	A	39	VAL	2.4
1	A	388	LYS	2.4
1	A	485	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	482	GLU	2.3
1	B	491	TYR	2.2
1	A	47	HIS	2.2
1	A	406	THR	2.1
1	B	389	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	AN2	A	601	27/27	0.89	0.11	48,69,88,89	0
2	AN2	B	601	27/27	0.90	0.10	35,66,103,105	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around AN2 A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around AN2 B 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.