



wwPDB EM Validation Summary Report ⓘ

Mar 8, 2026 – 05:19 AM UTC

PDB ID : 7YHS / pdb_00007yhs
EMDB ID : EMD-33837
Title : Structure of Csy-AcrIF4-dsDNA
Authors : Feng, Y.; Zhang, L.X.
Deposited on : 2022-07-14
Resolution : 3.37 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	14287 (2.87 - 3.87)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	434	16% 86% 11% ••
2	C	342	9% 68% 16% • 14%
2	D	342	9% 78% 15% • 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	342	
2	G	342	
2	H	342	
2	I	342	
3	E	327	
4	J	100	
5	L	187	
6	M	60	
7	N	54	
8	O	54	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 24596 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type I-F CRISPR-associated protein Csy1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	423	Total	C	N	O	S	0	0
			3039	1916	574	547	2		

- Molecule 2 is a protein called CRISPR-associated protein Csy3.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	293	Total	C	N	O	S	0	0
			2240	1408	403	427	2		
2	D	324	Total	C	N	O	S	0	0
			2425	1519	441	463	2		
2	F	334	Total	C	N	O	S	0	0
			2548	1602	462	482	2		
2	G	335	Total	C	N	O	S	0	0
			2529	1585	457	485	2		
2	H	325	Total	C	N	O	S	0	0
			2490	1565	454	469	2		
2	I	329	Total	C	N	O	S	0	0
			2514	1578	459	475	2		

- Molecule 3 is a protein called CRISPR type I-F/YPEST-associated protein Csy2.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	E	305	Total	C	N	O	S	0	0
			2373	1505	438	426	4		

- Molecule 4 is a protein called AcrIF4.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	J	94	Total	C	N	O	S	0	0
			737	464	131	138	4		

- Molecule 5 is a protein called Csy4.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	L	187	Total	C	N	O	S	0	0
			1428	892	285	248	3		

- Molecule 6 is a RNA chain called RNA (58-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
6	M	58	Total	C	N	O	P	0	0
			1226	549	213	407	57		

- Molecule 7 is a DNA chain called DNA (39-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
7	N	39	Total	C	N	O	P	0	0
			803	378	156	230	39		


- Molecule 8 is a DNA chain called DNA (5'-D(P*AP*GP*CP*AP*GP*CP*TP*GP*CP*AP*CP*C)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
8	O	12	Total	C	N	O	P	0	0
			244	115	47	70	12		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Type I-F CRISPR-associated protein Csy1




- Molecule 2: CRISPR-associated protein Csy3




- Molecule 2: CRISPR-associated protein Csy3




• Molecule 2: CRISPR-associated protein Csy3




• Molecule 2: CRISPR-associated protein Csy3





• Molecule 2: CRISPR-associated protein Csy3








• Molecule 2: CRISPR-associated protein Csy3




• Molecule 3: CRISPR type I-F/YPEST-associated protein Csy2




• Molecule 4: AcrIF4



• Molecule 5: Csy4





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	117510	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1700	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	2.392	Depositor
Minimum map value	-0.053	Depositor
Average map value	0.004	Depositor
Map value standard deviation	0.042	Depositor
Recommended contour level	0.06	Depositor
Map size (Å)	232.8, 232.8, 232.8	wwPDB
Map dimensions	240, 240, 240	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.97, 0.97, 0.97	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/3107	0.56	0/4241
2	C	0.23	0/2280	0.52	1/3097 (0.0%)
2	D	0.20	0/2466	0.46	0/3354
2	F	0.24	0/2594	0.50	0/3525
2	G	0.23	0/2573	0.46	0/3497
2	H	0.21	0/2533	0.47	0/3438
2	I	0.21	0/2559	0.46	0/3474
3	E	0.23	0/2430	0.53	0/3311
4	J	0.17	0/752	0.41	0/1018
5	L	0.43	0/1459	1.01	16/1973 (0.8%)
6	M	0.28	0/1368	0.58	0/2128
7	N	0.31	0/902	0.54	0/1390
8	O	0.18	0/273	0.36	0/418
All	All	0.25	0/25296	0.54	17/34864 (0.0%)

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	L	39	ARG	N-CA-C	11.15	123.18	111.14
5	L	17	PRO	N-CA-C	8.19	129.35	112.47
2	C	168	TRP	N-CA-C	7.17	118.89	111.14
5	L	13	PRO	CA-C-N	6.64	129.18	120.28
5	L	13	PRO	C-N-CA	6.64	129.18	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3039	0	2716	30	0
2	C	2240	0	2179	32	0
2	D	2425	0	2333	26	0
2	F	2548	0	2499	36	0
2	G	2529	0	2474	32	0
2	H	2490	0	2465	30	0
2	I	2514	0	2465	33	0
3	E	2373	0	2356	39	0
4	J	737	0	710	8	0
5	L	1428	0	1366	60	0
6	M	1226	0	625	15	0
7	N	803	0	435	7	0
8	O	244	0	134	0	0
All	All	24596	0	22757	296	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 296 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:L:40:ILE:CG2	5:L:62:ALA:HB2	1.75	1.14
5:L:40:ILE:HG21	5:L:62:ALA:HB2	1.28	1.06
5:L:40:ILE:HG21	5:L:62:ALA:CB	1.87	1.04
5:L:35:GLN:NE2	5:L:70:LEU:CG	2.31	0.94
5:L:35:GLN:HE21	5:L:70:LEU:HD12	1.33	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	421/434 (97%)	378 (90%)	43 (10%)	0	100	100
2	C	287/342 (84%)	267 (93%)	20 (7%)	0	100	100
2	D	320/342 (94%)	304 (95%)	16 (5%)	0	100	100
2	F	332/342 (97%)	302 (91%)	30 (9%)	0	100	100
2	G	333/342 (97%)	313 (94%)	20 (6%)	0	100	100
2	H	321/342 (94%)	302 (94%)	19 (6%)	0	100	100
2	I	325/342 (95%)	307 (94%)	18 (6%)	0	100	100
3	E	301/327 (92%)	270 (90%)	31 (10%)	0	100	100
4	J	90/100 (90%)	88 (98%)	2 (2%)	0	100	100
5	L	185/187 (99%)	150 (81%)	33 (18%)	2 (1%)	11	37
All	All	2915/3100 (94%)	2681 (92%)	232 (8%)	2 (0%)	49	77

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	L	17	PRO
5	L	18	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/365 (70%)	239 (94%)	16 (6%)	16	42
2	C	223/274 (81%)	207 (93%)	16 (7%)	13	39
2	D	235/274 (86%)	215 (92%)	20 (8%)	10	33
2	F	254/274 (93%)	229 (90%)	25 (10%)	7	27
2	G	252/274 (92%)	234 (93%)	18 (7%)	13	39
2	H	251/274 (92%)	235 (94%)	16 (6%)	16	42
2	I	251/274 (92%)	242 (96%)	9 (4%)	31	56
3	E	246/270 (91%)	227 (92%)	19 (8%)	12	37

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	J	73/77 (95%)	67 (92%)	6 (8%)	10	36
5	L	136/160 (85%)	127 (93%)	9 (7%)	15	41
All	All	2176/2516 (86%)	2022 (93%)	154 (7%)	15	39

5 of 154 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	57	GLU
4	J	63	THR
2	H	95	LEU
2	I	86	LEU
5	L	52	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 53 such sidechains are listed below:

Mol	Chain	Res	Type
2	F	310	GLN
2	G	163	ASN
5	L	29	HIS
2	F	316	GLN
2	G	94	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
6	M	57/60 (95%)	33 (57%)	3 (5%)

5 of 33 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
6	M	3	A
6	M	9	U
6	M	10	U
6	M	11	C
6	M	14	G

All (3) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
6	M	33	U
6	M	44	C
6	M	47	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-33837. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 120




Y Index: 120




Z Index: 120


6.2.2 Raw map



X Index: 120



Y Index: 120



Z Index: 120


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 121




Y Index: 141




Z Index: 165


6.3.2 Raw map



X Index: 118



Y Index: 151




Z Index: 87


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

This section was not generated.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 169 nm^3 ; this corresponds to an approximate mass of 153 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.297 $\text{\AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.297 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.37	-	-
Author-provided FSC curve	3.33	3.90	3.39
Unmasked-calculated*	3.60	4.20	3.76

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-33837 and PDB model 7YHS. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)

This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 81% of all backbone atoms, 78% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.06) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7790	 0.4040
A	 0.7270	 0.3580
C	 0.7410	 0.3300
D	 0.7230	 0.3220
E	 0.8320	 0.4440
F	 0.8270	 0.4300
G	 0.8590	 0.5070
H	 0.8680	 0.5290
I	 0.7910	 0.4550
J	 0.7180	 0.4060
L	 0.6630	 0.2500
M	 0.8300	 0.3930
N	 0.6950	 0.3470
O	 0.3400	 0.0900

