



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 06:41 AM UTC

PDB ID : 8YH0 / pdb_00008yh0
EMDB ID : EMD-39278
Title : A3R-Gi complex bound to NECA
Authors : Oshima, H.S.; Shihoya, W.; Nureki, O.
Deposited on : 2024-02-27
Resolution : 2.86 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	12017 (2.36 - 3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	R	794	
2	A	433	
2	G	433	
3	B	375	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	S	260	18% 73% 16% 11%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8693 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HA tag,Adenosine receptor A3,LgBiT,eGFP chimera.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	271	Total	C	N	O	S	0	0
			2198	1486	341	351	20		

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	-8	ASP	-	linker	UNP P03435
R	-7	TYR	-	linker	UNP P03435
R	-6	LYS	-	linker	UNP P03435
R	-5	ASP	-	linker	UNP P03435
R	-4	ASP	-	linker	UNP P03435
R	-3	ASP	-	linker	UNP P03435
R	-2	ASP	-	linker	UNP P03435
R	-1	ALA	-	linker	UNP P03435
R	0	MET	-	linker	UNP P03435
R	1	GLY	-	linker	UNP P03435
R	318	GLY	-	linker	UNP W5QED6
R	319	SER	-	linker	UNP W5QED6
R	320	GLY	-	linker	UNP W5QED6
R	321	GLY	-	linker	UNP W5QED6
R	322	GLY	-	linker	UNP W5QED6
R	323	GLY	-	linker	UNP W5QED6
R	324	SER	-	linker	UNP W5QED6
R	325	GLY	-	linker	UNP W5QED6
R	326	GLY	-	linker	UNP W5QED6
R	327	SER	-	linker	UNP W5QED6
R	328	SER	-	linker	UNP W5QED6
R	329	SER	-	linker	UNP W5QED6
R	330	GLY	-	linker	UNP W5QED6
R	331	GLY	-	linker	UNP W5QED6
R	490	GLY	-	linker	UNP W5QED6
R	491	GLY	-	linker	UNP W5QED6
R	492	SER	-	linker	UNP W5QED6
R	493	GLY	-	linker	UNP W5QED6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
R	494	GLY	-	linker	UNP W5QED6
R	495	GLY	-	linker	UNP W5QED6
R	496	GLY	-	linker	UNP W5QED6
R	497	SER	-	linker	UNP W5QED6
R	498	GLY	-	linker	UNP W5QED6
R	499	GLY	-	linker	UNP W5QED6
R	500	SER	-	linker	UNP W5QED6
R	501	SER	-	linker	UNP W5QED6
R	502	SER	-	linker	UNP W5QED6
R	503	GLY	-	linker	UNP W5QED6
R	504	GLY	-	linker	UNP W5QED6
R	505	LEU	-	linker	UNP W5QED6
R	506	GLU	-	linker	UNP W5QED6
R	507	VAL	-	linker	UNP W5QED6
R	508	LEU	-	linker	UNP W5QED6
R	509	PHE	-	linker	UNP W5QED6
R	510	GLN	-	linker	UNP W5QED6
R	511	GLY	-	linker	UNP W5QED6
R	512	PRO	-	linker	UNP W5QED6
R	513	GLY	-	linker	UNP W5QED6
R	514	SER	-	linker	UNP W5QED6
R	515	ALA	-	linker	UNP W5QED6
R	516	ALA	-	linker	UNP W5QED6
R	517	ALA	-	linker	UNP W5QED6
R	518	ALA	-	linker	UNP W5QED6
R	757	SER	-	expression tag	UNP A0A5P9VSM6
R	758	GLY	-	expression tag	UNP A0A5P9VSM6
R	759	LEU	-	expression tag	UNP A0A5P9VSM6
R	760	ARG	-	expression tag	UNP A0A5P9VSM6
R	761	SER	-	expression tag	UNP A0A5P9VSM6
R	762	HIS	-	expression tag	UNP A0A5P9VSM6
R	763	HIS	-	expression tag	UNP A0A5P9VSM6
R	764	HIS	-	expression tag	UNP A0A5P9VSM6
R	765	HIS	-	expression tag	UNP A0A5P9VSM6
R	766	HIS	-	expression tag	UNP A0A5P9VSM6
R	767	HIS	-	expression tag	UNP A0A5P9VSM6
R	768	HIS	-	expression tag	UNP A0A5P9VSM6
R	769	HIS	-	expression tag	UNP A0A5P9VSM6

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2, Guanine nucleotide-binding protein G(i) subunit alpha-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	A	211	Total	C	N	O	S	0	0
			1700	1082	286	320	12		
2	G	53	Total	C	N	O	S	0	0
			407	257	70	77	3		

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	GLY	-	linker	UNP P59768
A	-6	SER	-	linker	UNP P59768
A	-5	ALA	-	linker	UNP P59768
A	-4	GLY	-	linker	UNP P59768
A	-3	SER	-	linker	UNP P59768
A	-2	ALA	-	linker	UNP P59768
A	-1	GLY	-	linker	UNP P59768
A	0	SER	-	linker	UNP P59768
A	1	ALA	-	linker	UNP P59768
A	2	MET	-	linker	UNP P59768
G	72	GLY	-	linker	UNP P59768
G	73	SER	-	linker	UNP P59768
G	74	ALA	-	linker	UNP P59768
G	75	GLY	-	linker	UNP P59768
G	76	SER	-	linker	UNP P59768
G	77	ALA	-	linker	UNP P59768
G	78	GLY	-	linker	UNP P59768
G	79	SER	-	linker	UNP P59768
G	80	ALA	-	linker	UNP P59768
G	81	MET	-	linker	UNP P59768

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	B	336	Total	C	N	O	S	0	0
			2583	1593	465	504	21		

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	MET	-	initiating methionine	UNP P62871
B	-3	GLY	-	expression tag	UNP P62871
B	-2	SER	-	expression tag	UNP P62871
B	-1	LEU	-	expression tag	UNP P62871

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	LEU	-	expression tag	UNP P62871
B	1	GLN	-	expression tag	UNP P62871
B	341	GLY	-	expression tag	UNP P62871
B	342	ALA	-	expression tag	UNP P62871
B	343	SER	-	expression tag	UNP P62871
B	344	GLY	-	expression tag	UNP P62871
B	345	GLY	-	expression tag	UNP P62871
B	346	GLY	-	expression tag	UNP P62871
B	347	SER	-	expression tag	UNP P62871
B	348	GLY	-	expression tag	UNP P62871
B	349	GLY	-	expression tag	UNP P62871
B	350	ASN	-	expression tag	UNP P62871
B	351	SER	-	expression tag	UNP P62871
B	352	GLY	-	expression tag	UNP P62871
B	353	SER	-	expression tag	UNP P62871
B	354	SER	-	expression tag	UNP P62871
B	355	GLY	-	expression tag	UNP P62871
B	356	GLY	-	expression tag	UNP P62871
B	357	SER	-	expression tag	UNP P62871
B	358	SER	-	expression tag	UNP P62871
B	359	GLY	-	expression tag	UNP P62871
B	360	VAL	-	expression tag	UNP P62871
B	361	SER	-	expression tag	UNP P62871
B	362	GLY	-	expression tag	UNP P62871
B	363	TRP	-	expression tag	UNP P62871
B	364	ARG	-	expression tag	UNP P62871
B	365	LEU	-	expression tag	UNP P62871
B	366	PHE	-	expression tag	UNP P62871
B	367	LYS	-	expression tag	UNP P62871
B	368	LYS	-	expression tag	UNP P62871
B	369	ILE	-	expression tag	UNP P62871
B	370	SER	-	expression tag	UNP P62871

- Molecule 4 is a protein called scfv16.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	S	232	Total	C	N	O	S	0	0
			1783	1131	295	347	10		

- Molecule 5 is N-ETHYL-5'-CARBOXAMIDO ADENOSINE (CCD ID: NEC) (formula: $C_{12}H_{16}N_6O_4$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
5	R	1	22	12	6	4	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HA tag,Adenosine receptor A3,LgBiT,eGFP chimera



- Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2,Guanine nucleotide-binding protein G(i) subunit alpha-1




- Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2, Guanine nucleotide-binding protein G(i) subunit alpha-1





- Molecule 3: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1





• Molecule 4: scfv16



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	210192	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1600	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	33.666	Depositor
Minimum map value	-15.369	Depositor
Average map value	-0.004	Depositor
Map value standard deviation	1.047	Depositor
Recommended contour level	7.7	Depositor
Map size (Å)	232.38, 232.38, 232.38	wwPDB
Map dimensions	180, 180, 180	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.291, 1.291, 1.291	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NEC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	R	0.14	0/2254	0.35	0/3070
2	A	0.18	0/1728	0.36	0/2313
2	G	0.14	0/413	0.40	1/558 (0.2%)
3	B	0.15	0/2630	0.35	0/3566
4	S	0.14	0/1827	0.35	0/2477
All	All	0.15	0/8852	0.35	1/11984 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	53	PRO	N-CA-C	6.59	121.41	111.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	2198	0	2328	36	0
2	A	1700	0	1700	25	0
2	G	407	0	418	8	0
3	B	2583	0	2488	60	0
4	S	1783	0	1717	32	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	R	22	0	16	0	0
All	All	8693	0	8667	150	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 150 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:14:GLU:OE1	4:S:57:THR:OG1	2.10	0.69
4:S:166:LEU:HB2	4:S:176:LEU:HD11	1.75	0.69
4:S:61:ALA:O	4:S:65:LYS:N	2.27	0.67
3:B:86:THR:O	3:B:87:THR:OG1	2.08	0.66
4:S:83:MET:HE1	4:S:117:LEU:HD22	1.77	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	R	267/794 (34%)	258 (97%)	9 (3%)	0	100	100
2	A	205/433 (47%)	199 (97%)	6 (3%)	0	100	100
2	G	51/433 (12%)	49 (96%)	2 (4%)	0	100	100
3	B	334/375 (89%)	323 (97%)	11 (3%)	0	100	100
4	S	228/260 (88%)	215 (94%)	12 (5%)	1 (0%)	30	48
All	All	1085/2295 (47%)	1044 (96%)	40 (4%)	1 (0%)	49	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	S	68	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	R	247/685 (36%)	247 (100%)	0	100	100
2	A	187/366 (51%)	187 (100%)	0	100	100
2	G	43/366 (12%)	43 (100%)	0	100	100
3	B	279/305 (92%)	279 (100%)	0	100	100
4	S	197/209 (94%)	197 (100%)	0	100	100
All	All	953/1931 (49%)	953 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
4	S	157	ASN
4	S	159	ASN
2	A	322	HIS
2	A	331	ASN
2	A	333	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NEC	R	801	-	24,24,24	3.07	8 (33%)	34,35,35	4.07	14 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NEC	R	801	-	-	2/11/27/27	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	R	801	NEC	C5-N7	7.51	1.52	1.39
5	R	801	NEC	C8-N9	-6.85	1.25	1.37
5	R	801	NEC	C5-C4	5.21	1.48	1.39
5	R	801	NEC	C5'-N5'	5.10	1.45	1.33
5	R	801	NEC	C6-N6	4.43	1.45	1.34

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	R	801	NEC	C4-N9-C8	14.87	121.35	105.74
5	R	801	NEC	C4-C5-N7	-8.84	100.47	110.58
5	R	801	NEC	N3-C4-N9	6.25	137.80	127.17
5	R	801	NEC	C5-C4-N3	-5.95	118.52	126.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	R	801	NEC	C6-C5-N7	5.88	143.43	132.09

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	R	801	NEC	O4'-C4'-C5'-N5'
5	R	801	NEC	O4'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-39278. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 90



Y Index: 90




Z Index: 90


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 97



Y Index: 93




Z Index: 73


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 7.7. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 44 nm^3 ; this corresponds to an approximate mass of 40 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.350 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-39278 and PDB model 8YH0. Per-residue inclusion information can be found in section [3](#) on page [9](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 7.7 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (7.7).

9.4 Atom inclusion [i](#)



At the recommended contour level, 76% of all backbone atoms, 62% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (7.7) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6200	0.4420
A	0.5400	0.4130
B	0.6890	0.4750
G	0.2860	0.2890
R	0.6520	0.4250
S	0.6360	0.4770

