



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 07:54 PM UTC

PDB ID : 1YI6 / pdb_00001yi6
Title : C-term tail segment of human tyrosine kinase (258-533)
Authors : Fleury, D.; Sarubbi, E.; Courjaud, A.; Guitton, J.D.; Ducruix, A.
Deposited on : 2005-01-11
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	276	 87% 11% •
1	B	276	 81% 12% 5% •

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4995 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	P	S	0	0	0
			2230	1422	373	417	2	16			
1	B	276	Total	C	N	O	P	S	0	0	0
			2230	1422	373	417	2	16			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	335	PTR	TYR	modified residue	UNP P12931
A	416	PTR	TYR	modified residue	UNP P12931
B	335	PTR	TYR	modified residue	UNP P12931
B	416	PTR	TYR	modified residue	UNP P12931

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	304	Total	O	0	0
			304	304		
2	B	231	Total	O	0	0
			231	231		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.98Å 63.42Å 76.69Å 90.00° 109.54° 90.00°	Depositor
Resolution (Å)	19.00 – 2.00	Depositor
% Data completeness (in resolution range)	98.0 (19.00-2.00)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	0.07	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.216 , 0.252	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4995	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/2248	0.87	4/3041 (0.1%)
1	B	0.42	0/2248	1.13	18/3041 (0.6%)
All	All	0.41	0/4496	1.01	22/6082 (0.4%)

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	414	ASN	N-CA-C	15.46	128.22	111.36
1	B	301	THR	N-CA-C	14.48	126.75	110.97
1	B	298	LYS	CA-C-N	10.91	133.47	119.84
1	B	298	LYS	C-N-CA	10.91	133.47	119.84
1	B	277	CYS	N-CA-C	9.31	130.63	110.80
1	A	342	SER	N-CA-C	8.84	122.03	111.33
1	B	424	PHE	CA-C-N	7.75	129.52	119.84
1	B	424	PHE	C-N-CA	7.75	129.52	119.84
1	B	342	SER	N-CA-C	7.68	121.58	111.75
1	B	298	LYS	CB-CA-C	7.53	120.22	109.38
1	B	278	PHE	N-CA-C	-7.29	95.28	110.80
1	B	299	PRO	N-CA-C	7.11	127.12	112.47
1	B	300	GLY	N-CA-C	6.89	129.52	113.18
1	A	330	SER	N-CA-C	5.95	120.15	113.01
1	A	283	MET	N-CA-C	-5.70	99.95	109.24
1	B	297	LEU	CA-C-N	-5.70	113.69	121.61
1	B	297	LEU	C-N-CA	-5.70	113.69	121.61
1	A	522	SER	N-CA-C	5.52	118.48	111.69
1	B	519	TYR	N-CA-C	5.33	120.11	111.37
1	B	413	ASP	CA-C-N	5.32	127.84	120.29
1	B	413	ASP	C-N-CA	5.32	127.84	120.29
1	B	415	GLU	N-CA-C	5.17	125.47	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2230	0	2198	14	0
1	B	2230	0	2198	35	0
2	A	304	0	0	0	0
2	B	231	0	0	1	0
All	All	4995	0	4396	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:423:LYS:HE2	1:A:423:LYS:H	1.53	0.74
1:A:329:VAL:HB	1:A:335:PTR:HB3	1.71	0.71
1:B:308:LEU:HD21	1:B:334:ILE:HD11	1.77	0.66
1:B:418:ALA:C	1:B:420:GLN:H	2.05	0.65
1:B:418:ALA:HB3	1:B:420:GLN:HE21	1.62	0.64
1:B:275:GLN:O	1:B:275:GLN:HG3	1.99	0.61
1:B:525:PRO:HB2	1:B:526:GLN:HE21	1.66	0.61
1:B:280:GLU:HG3	1:B:298:LYS:HZ2	1.65	0.61
1:A:319:HIS:HB3	1:A:322:LEU:HG	1.82	0.60
1:B:525:PRO:HB2	1:B:526:GLN:NE2	2.22	0.54
1:A:277:CYS:O	1:A:301:THR:HG21	2.07	0.54
1:B:418:ALA:C	1:B:420:GLN:N	2.67	0.53
1:A:353:GLU:O	1:A:356:LYS:HG2	2.09	0.52
1:A:438:ARG:HD3	1:B:397:ASN:OD1	2.10	0.52
1:B:424:PHE:CD2	1:B:424:PHE:O	2.63	0.51
1:A:413:ASP:HB3	1:A:416:PTR:HB2	1.92	0.51
1:B:381:ASN:HB3	2:B:642:HOH:O	2.12	0.50
1:B:268:ARG:HH11	1:B:268:ARG:HB3	1.78	0.49
1:B:308:LEU:CD2	1:B:334:ILE:HD11	2.43	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:416:PTR:CG	1:B:417:THR:N	2.76	0.48
1:B:426:ILE:HD11	1:B:472:LEU:HD13	1.95	0.48
1:A:500:ARG:O	1:A:506:ARG:HD2	2.14	0.48
1:A:423:LYS:H	1:A:423:LYS:CE	2.23	0.47
1:B:323:VAL:HG21	1:B:393:LEU:HD12	1.95	0.47
1:B:329:VAL:O	1:B:334:ILE:HG13	2.13	0.47
1:A:334:ILE:O	1:A:335:PTR:HD1	2.15	0.47
1:B:477:ARG:O	1:B:477:ARG:HG3	2.14	0.47
1:B:297:LEU:HB2	1:B:307:PHE:CE1	2.50	0.46
1:B:280:GLU:HG3	1:B:298:LYS:NZ	2.31	0.46
1:B:275:GLN:OE1	1:B:277:CYS:SG	2.75	0.45
1:B:311:ALA:O	1:B:315:LYS:HG3	2.17	0.45
1:B:419:ARG:HH11	1:B:419:ARG:HG3	1.82	0.44
1:B:278:PHE:O	1:B:297:LEU:HD12	2.18	0.43
1:B:423:LYS:HE3	1:B:423:LYS:H	1.83	0.43
1:B:424:PHE:O	1:B:426:ILE:N	2.52	0.42
1:B:375:ALA:O	1:B:379:ARG:HG3	2.19	0.42
1:A:346:LEU:HD22	1:A:350:LEU:HD22	2.01	0.41
1:B:298:LYS:HA	1:B:299:PRO:HD3	1.80	0.41
1:A:278:PHE:CD1	1:A:302:MET:HE2	2.55	0.41
1:B:523:THR:C	1:B:525:PRO:HD3	2.45	0.41
1:B:280:GLU:CD	1:B:298:LYS:HZ3	2.29	0.41
1:A:272:LYS:HG2	1:A:282:TRP:CE2	2.55	0.41
1:B:275:GLN:NE2	1:B:280:GLU:OE1	2.54	0.41
1:B:362:GLN:O	1:B:366:MET:HG3	2.21	0.41
1:A:375:ALA:O	1:A:379:ARG:HG3	2.21	0.41
1:B:360:LEU:HD22	1:B:364:VAL:HG23	2.03	0.41
1:B:361:PRO:HA	1:B:520:PHE:CE2	2.56	0.40
1:B:466:MET:HE1	1:B:479:TYR:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/276 (99%)	264 (97%)	7 (3%)	1 (0%)	30	27
1	B	272/276 (99%)	252 (93%)	15 (6%)	5 (2%)	6	3
All	All	544/552 (99%)	516 (95%)	22 (4%)	6 (1%)	11	7

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	277	CYS
1	B	299	PRO
1	B	425	PRO
1	B	418	ALA
1	A	404	ASP
1	B	417	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/236 (100%)	225 (95%)	11 (5%)	23	22
1	B	236/236 (100%)	220 (93%)	16 (7%)	14	11
All	All	472/472 (100%)	445 (94%)	27 (6%)	18	15

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	265	GLU
1	A	272	LYS
1	A	297	LEU
1	A	346	LEU
1	A	350	LEU
1	A	398	LEU
1	A	414	ASN
1	A	423	LYS
1	A	426	ILE
1	A	491	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	533	LEU
1	B	268	ARG
1	B	275	GLN
1	B	278	PHE
1	B	280	GLU
1	B	299	PRO
1	B	301	THR
1	B	304	PRO
1	B	346	LEU
1	B	350	LEU
1	B	360	LEU
1	B	420	GLN
1	B	423	LYS
1	B	424	PHE
1	B	425	PRO
1	B	426	ILE
1	B	477	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	287	ASN
1	A	381	ASN
1	A	468	ASN
1	A	526	GLN
1	A	528	GLN
1	B	312	GLN
1	B	324	GLN
1	B	420	GLN
1	B	526	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PTR	A	416	1	15,16,17	1.28	2 (13%)	17,22,24	1.17	1 (5%)
1	PTR	B	416	1	15,16,17	1.00	0	17,22,24	1.25	1 (5%)
1	PTR	B	335	1	15,16,17	1.10	0	17,22,24	1.34	2 (11%)
1	PTR	A	335	1	15,16,17	0.97	1 (6%)	17,22,24	1.21	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	416	1	-	1/10/11/13	0/1/1/1
1	PTR	B	416	1	-	2/10/11/13	0/1/1/1
1	PTR	B	335	1	-	2/10/11/13	0/1/1/1
1	PTR	A	335	1	-	4/10/11/13	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	416	PTR	P-OH	2.37	1.64	1.59
1	A	416	PTR	OH-CZ	2.14	1.45	1.40
1	A	335	PTR	P-O2P	-2.05	1.47	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	335	PTR	CD2-CE2-CZ	-2.65	116.70	119.73
1	B	335	PTR	CD2-CE2-CZ	-2.65	116.71	119.73
1	B	416	PTR	CD2-CE2-CZ	-2.59	116.78	119.73
1	A	416	PTR	CD2-CE2-CZ	-2.26	117.15	119.73
1	B	335	PTR	CE1-CD1-CG	-2.07	118.28	121.00

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	416	PTR	C-CA-CB-CG
1	A	335	PTR	CA-CB-CG-CD1
1	A	335	PTR	CA-CB-CG-CD2
1	A	335	PTR	N-CA-CB-CG
1	B	335	PTR	CA-CB-CG-CD1
1	B	335	PTR	CA-CB-CG-CD2
1	A	335	PTR	C-CA-CB-CG
1	A	416	PTR	C-CA-CB-CG
1	B	416	PTR	CZ-OH-P-O2P

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	416	PTR	1	0
1	B	416	PTR	1	0
1	A	335	PTR	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.