



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:54 PM UTC

PDB ID : 6YIH / pdb_00006yih
Title : Structure of Chromosomal Passenger Complex (CPC) bound to phosphorylated Histone 3 peptide at 2.6 Å.
Authors : Serena, M.; Elliott, P.R.; Barr, F.A.
Deposited on : 2020-04-01
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	144	24% 84% 10% 6%
2	B	67	34% 79% 10% 10%
3	C	76	22% 46% • 50%
4	D	12	25% 50% 8% 42%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3704 atoms, of which 1770 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Baculoviral IAP repeat-containing protein 5.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	135	Total	C	H	N	O	S	0	0	0
			2009	669	962	177	194	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP O15392
A	0	PRO	-	expression tag	UNP O15392

- Molecule 2 is a protein called Borealin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	60	Total	C	H	N	O	S	0	0	0
			958	311	471	82	93	1			

- Molecule 3 is a protein called Inner centromere protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	38	Total	C	H	N	O	S	0	0	0
			573	188	278	49	54	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	5	GLY	ALA	conflict	UNP Q9NQS7


- Molecule 4 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	D	7	Total	C	H	N	O	P	0	0	0
			116	31	59	12	13	1			

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	16	Total	O	0	0
			16	16		
7	B	24	Total	O	0	0
			24	24		
7	C	1	Total	O	0	0
			1	1		
7	D	1	Total	O	0	0
			1	1		

4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	99.36Å 99.36Å 56.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.07 – 2.55 47.07 – 2.55	Depositor EDS
% Data completeness (in resolution range)	94.0 (47.07-2.55) 94.3 (47.07-2.55)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.54Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.234 , 0.259 0.239 , 0.262	Depositor DCC
R_{free} test set	494 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	53.2	Xtriage
Anisotropy	0.061	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.053 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	3704	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, TPO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.20	0/1077	0.40	0/1459
2	B	0.17	0/493	0.35	0/665
3	C	0.19	0/299	0.44	0/404
4	D	0.11	0/44	0.29	0/55
All	All	0.19	0/1913	0.39	0/2583

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1047	962	960	12	0
2	B	487	471	470	6	0
3	C	295	278	276	2	0
4	D	57	59	59	0	0
5	A	1	0	0	0	0
6	A	5	0	0	0	0
7	A	16	0	0	1	0
7	B	24	0	0	1	0
7	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	D	1	0	0	0	0
All	All	1934	1770	1765	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:ILE:CD1	2:B:50:VAL:HG21	2.26	0.66
3:C:33:TRP:CE3	3:C:34:LEU:HD23	2.31	0.66
1:A:21:THR:HG23	1:A:22:PHE:CD1	2.33	0.64
2:B:66:ARG:NH2	7:B:103:HOH:O	2.38	0.57
1:A:132:ARG:NH2	7:A:301:HOH:O	2.41	0.53
1:A:124:PHE:HD1	3:C:31:LEU:HD12	1.75	0.52
1:A:14:LEU:HD22	1:A:94:GLU:HG3	1.95	0.49
1:A:25:TRP:CD2	1:A:26:PRO:HD2	2.51	0.46
2:B:25:LEU:HD23	2:B:25:LEU:N	2.32	0.45
1:A:18:ARG:C	1:A:21:THR:HG22	2.43	0.43
1:A:6:LEU:HD11	2:B:71:LEU:CD2	2.48	0.42
2:B:63:LYS:O	2:B:67:GLU:HG3	2.19	0.42
1:A:36:GLU:H	1:A:36:GLU:CD	2.28	0.42
1:A:132:ARG:O	1:A:136:GLU:N	2.52	0.42
1:A:18:ARG:O	1:A:21:THR:HG22	2.21	0.41
1:A:125:GLU:HG3	2:B:36:ILE:HD11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/144 (92%)	131 (98%)	2 (2%)	0	100	100
2	B	58/67 (87%)	58 (100%)	0	0	100	100
3	C	36/76 (47%)	35 (97%)	1 (3%)	0	100	100
4	D	4/12 (33%)	4 (100%)	0	0	100	100
All	All	231/299 (77%)	228 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	105/124 (85%)	105 (100%)	0	100	100
2	B	50/63 (79%)	50 (100%)	0	100	100
3	C	30/73 (41%)	30 (100%)	0	100	100
4	D	4/8 (50%)	4 (100%)	0	100	100
All	All	189/268 (70%)	189 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	TPO	D	3	4	8,10,11	1.67	1 (12%)	10,14,16	1.61	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TPO	D	3	4	-	1/9/11/13	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	3	TPO	P-O1P	3.60	1.61	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	3	TPO	P-OG1-CB	-4.25	111.77	123.33

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	3	TPO	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	SO4	A	202	-	4,4,4	0.25	0	6,6,6	0.06	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	135/144 (93%)	1.36	35 (25%)	1 1	36, 71, 199, 237	0
2	B	60/67 (89%)	2.02	23 (38%)	1 0	36, 70, 226, 303	0
3	C	38/76 (50%)	1.87	17 (44%)	0 0	45, 112, 171, 181	0
4	D	6/12 (50%)	2.34	3 (50%)	0 0	72, 108, 193, 236	0
All	All	239/299 (79%)	1.63	78 (32%)	1 0	36, 74, 204, 303	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	21	LEU	6.7
1	A	4	PRO	5.4
2	B	19	ARG	5.1
2	B	25	LEU	5.0
4	D	7	ALA	4.9
1	A	138	LEU	4.9
2	B	32	VAL	4.8
2	B	22	ALA	4.7
2	B	17	ARG	4.4
2	B	28	PHE	4.3
4	D	6	THR	4.2
1	A	6	LEU	4.1
1	A	5	THR	4.0
2	B	68	MET	3.9
2	B	18	ARG	3.8
3	C	31	LEU	3.8
3	C	44	MET	3.8
3	C	41	ALA	3.7
2	B	24	PHE	3.7
1	A	49	GLU	3.6
2	B	40	GLU	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	36	GLU	3.5
2	B	37	LYS	3.5
1	A	135	ILE	3.5
3	C	30	ASP	3.4
2	B	26	LYS	3.4
2	B	33	GLU	3.3
3	C	28	ASN	3.2
3	C	37	ILE	3.2
1	A	124	PHE	3.1
2	B	34	ILE	3.1
2	B	30	ARG	3.0
1	A	23	LYS	3.0
1	A	137	GLN	3.0
1	A	128	ALA	2.9
3	C	35	GLU	2.9
3	C	26	MET	2.9
1	A	132	ARG	2.9
1	A	117	THR	2.9
1	A	136	GLU	2.9
1	A	133	ARG	2.8
1	A	134	ALA	2.8
1	A	131	VAL	2.8
2	B	29	ASP	2.8
2	B	67	GLU	2.7
3	C	27	ASP	2.7
1	A	50	ASN	2.7
1	A	130	LYS	2.7
3	C	42	GLU	2.6
1	A	53	ASP	2.6
1	A	22	PHE	2.6
2	B	75	ALA	2.6
1	A	116	GLU	2.5
1	A	61	PHE	2.5
1	A	89	VAL	2.4
3	C	43	ARG	2.4
1	A	52	PRO	2.4
1	A	81	SER	2.4
1	A	83	GLY	2.3
2	B	43	ARG	2.3
3	C	24	CYS	2.3
1	A	29	GLU	2.3
1	A	127	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	74	ILE	2.3
1	A	14	LEU	2.3
2	B	20	LYS	2.2
2	B	31	GLU	2.2
1	A	93	PHE	2.2
4	D	5	GLN	2.2
3	C	40	GLU	2.2
1	A	46	CYS	2.1
3	C	29	LYS	2.1
1	A	118	ASN	2.0
3	C	34	LEU	2.0
1	A	90	LYS	2.0
1	A	70	ASP	2.0
2	B	66	ARG	2.0
3	C	32	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	TPO	D	3	11/12	0.91	0.11	65,79,97,114	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	SO4	A	202	5/5	0.81	0.14	104,105,105,105	0
5	ZN	A	201	1/1	0.99	0.02	51,51,51,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.