



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:33 AM UTC

PDB ID : 6YIO / pdb_00006yio
Title : CRYSTAL STRUCTURE OF FAB RG6292 IN COMPLEX WITH CD25 ECD
Authors : Benz, J.; Koll, H.; Leibrock, L.
Deposited on : 2020-04-01
Resolution : 1.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1296 (1.84-1.84)
Clashscore	190562	1329 (1.84-1.84)
Ramachandran outliers	187476	1318 (1.84-1.84)
Sidechain outliers	187428	1318 (1.84-1.84)
RSRZ outliers	180081	1296 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	203	
2	H	227	
3	L	214	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-2 receptor subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	134	Total	C	N	O	S	0	2	0
			1082	671	194	199	18			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	193	ALA	-	expression tag	UNP P01589
B	194	HIS	-	expression tag	UNP P01589
B	195	HIS	-	expression tag	UNP P01589
B	196	HIS	-	expression tag	UNP P01589
B	197	HIS	-	expression tag	UNP P01589
B	198	HIS	-	expression tag	UNP P01589
B	199	HIS	-	expression tag	UNP P01589
B	200	HIS	-	expression tag	UNP P01589
B	201	HIS	-	expression tag	UNP P01589
B	202	HIS	-	expression tag	UNP P01589
B	203	HIS	-	expression tag	UNP P01589

- Molecule 2 is a protein called FAB FRAGMENT HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	216	Total	C	N	O	S	0	4	0
			1603	1011	265	318	9			

- Molecule 3 is a protein called FAB FRAGMENT LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	213	Total	C	N	O	S	0	2	0
			1650	1036	274	335	5			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	108	Total 108	O 108	0	0
4	H	246	Total 246	O 246	0	0
4	L	235	Total 235	O 235	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-2 receptor subunit alpha



- Molecule 2: FAB FRAGMENT HEAVY CHAIN



- Molecule 3: FAB FRAGMENT LIGHT CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.38Å 44.32Å 140.09Å 90.00° 92.41° 90.00°	Depositor
Resolution (Å)	69.98 – 1.83 69.98 – 1.83	Depositor EDS
% Data completeness (in resolution range)	99.9 (69.98-1.83) 99.9 (69.98-1.83)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 1.83Å)	Xtriage
Refinement program	BUSTER 2.11.7 (6-FEB-2020)	Depositor
R, R_{free}	0.203 , 0.231 0.205 , 0.237	Depositor DCC
R_{free} test set	2889 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	25.8	Xtriage
Anisotropy	0.437	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 36.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.022 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4924	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.62	0/1111	0.94	0/1501
2	H	0.72	0/1643	0.92	2/2236 (0.1%)
3	L	0.72	0/1689	0.91	0/2294
All	All	0.69	0/4443	0.92	2/6031 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	64	PHE	CA-CB-CG	-6.84	106.95	113.80
2	H	129	SER	N-CA-C	-5.06	101.46	109.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1082	0	1021	5	0
2	H	1603	0	1589	6	0
3	L	1650	0	1611	5	0
4	B	108	0	0	0	0
4	H	246	0	0	0	0
4	L	235	0	0	0	0
All	All	4924	0	4221	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:131:CYS:HG	1:B:163:CYS:HG	0.88	0.83
2:H:173:HIS:CD2	3:L:137:ASN:HD21	2.06	0.72
2:H:173:HIS:HD2	3:L:137:ASN:HD21	1.39	0.71
3:L:39:LYS:HB2	3:L:42:LYS:HD2	1.82	0.61
3:L:4:MET:HE3	3:L:23:CYS:SG	2.53	0.48
2:H:97:ALA:HB1	2:H:109:PHE:HB3	1.98	0.45
2:H:36:TRP:CE2	2:H:81:MET:HB2	2.52	0.43
1:B:16:LYS:HG3	2:H:54:ILE:HG22	1.99	0.43
1:B:16:LYS:HD3	2:H:55:PHE:HA	2.01	0.43
1:B:136:ARG:HD2	3:L:32:TRP:CZ2	2.54	0.42
1:B:13:ALA:HB2	1:B:163:CYS:SG	2.60	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	132/203 (65%)	130 (98%)	2 (2%)	0	100	100
2	H	216/227 (95%)	210 (97%)	5 (2%)	1 (0%)	24	12
3	L	213/214 (100%)	208 (98%)	5 (2%)	0	100	100
All	All	561/644 (87%)	548 (98%)	12 (2%)	1 (0%)	43	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	102	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	120/181 (66%)	118 (98%)	2 (2%)	53	37
2	H	182/188 (97%)	180 (99%)	2 (1%)	65	54
3	L	189/188 (100%)	187 (99%)	2 (1%)	65	54
All	All	491/557 (88%)	485 (99%)	6 (1%)	63	51

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	37	ILE
1	B	133	GLN
2	H	23	LYS
2	H	215	LYS
3	L	105	GLU
3	L	147	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	12	HIS
1	B	57	ASN
1	B	60	GLN
1	B	151	HIS
2	H	208	ASN
3	L	92	ASN
3	L	137	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	134/203 (66%)	1.05	27 (20%) 3 2	16, 40, 76, 86	2 (1%)
2	H	216/227 (95%)	-0.08	3 (1%) 73 81	11, 26, 39, 59	4 (1%)
3	L	213/214 (99%)	0.06	2 (0%) 81 87	17, 30, 41, 51	2 (0%)
All	All	563/644 (87%)	0.24	32 (5%) 29 31	11, 29, 57, 86	8 (1%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	1	GLN	4.8
1	B	34	PHE	4.6
1	B	37	ILE	4.2
1	B	92	MET	3.8
1	B	4	ASP	3.5
1	B	150	THR	3.3
1	B	61	CYS	3.2
1	B	39	SER	3.2
1	B	151	HIS	3.2
1	B	32	ARG	3.1
2	H	223	LYS	3.1
1	B	152	GLY	3.0
1	B	31	LYS	3.0
1	B	33	GLY	2.9
2	H	222	PRO	2.8
1	B	40	GLY	2.5
1	B	164	THR	2.4
1	B	100	LEU	2.4
1	B	35	ARG	2.4
1	B	149	MET	2.4
1	B	95	VAL	2.3
1	B	59	CYS	2.3
3	L	154	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1	GLU	2.2
1	B	38	LYS	2.2
1	B	3	CYS	2.2
1	B	114	ALA	2.2
1	B	36	ARG	2.2
3	L	77	SER	2.1
1	B	94	PRO	2.1
1	B	115	THR	2.1
1	B	52	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.