



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 08:52 AM UTC

PDB ID : 4YJ1 / pdb_00004yj1
Title : Crystal structure of T. brucei MRB1590-ADP bound to poly-U RNA
Authors : Schumacher, M.A.
Deposited on : 2015-03-03
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	615	6% 82% 14% .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9716 atoms, of which 4581 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	600	9154	2866	4569	819	874	26	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ARG	deletion	UNP Q57ZF2

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	P		
2	A	1	39	10	12	5	10	2	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total 2	Mg 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	521	Total 521	O 521	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	215.43Å 215.43Å 100.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.80 – 2.05 41.80 – 2.05	Depositor EDS
% Data completeness (in resolution range)	95.6 (41.80-2.05) 94.8 (41.80-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.43 (at 2.05Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.173 , 0.203 (Not available) , 0.184	Depositor DCC
R_{free} test set	3711 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	24.6	Xtriage
Anisotropy	0.425	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9716	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.13	7/4670 (0.1%)	1.32	11/6332 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	215	VAL	CA-CB	8.33	1.58	1.54
1	A	238	HIS	CA-C	6.14	1.60	1.52
1	A	242	VAL	CA-CB	5.83	1.61	1.54
1	A	485	ALA	CA-CB	5.24	1.61	1.53
1	A	558	ILE	N-CA	5.22	1.52	1.46
1	A	312	ALA	CA-CB	5.04	1.60	1.53
1	A	443	ALA	CA-CB	5.01	1.61	1.53

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	578	GLU	CA-C-N	45.35	176.53	119.84
1	A	578	GLU	C-N-CA	45.35	176.53	119.84
1	A	578	GLU	C-N-CD	-11.78	76.70	125.00
1	A	579	PRO	CA-N-CD	-7.31	101.77	112.00
1	A	633	THR	CA-C-N	5.49	125.58	119.32
1	A	633	THR	C-N-CA	5.49	125.58	119.32
1	A	242	VAL	CG1-CB-CG2	-5.47	98.76	110.80
1	A	105	CYS	CA-C-N	-5.09	114.71	119.85
1	A	105	CYS	C-N-CA	-5.09	114.71	119.85
1	A	337	ASP	N-CA-C	5.05	117.52	111.71
1	A	327	GLU	CB-CA-C	-5.01	102.48	110.79

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4585	4569	4550	74	0
2	A	27	12	12	4	0
3	A	2	0	0	0	0
4	A	521	0	0	15	1
All	All	5135	4581	4562	74	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:525:ARG:HB3	4:A:827:HOH:O	1.46	1.12
1:A:640:ARG:HH11	1:A:640:ARG:HG3	1.30	0.96
1:A:183:GLN:N	4:A:1290:HOH:O	2.05	0.90
1:A:621:CYS:SG	4:A:1299:HOH:O	2.31	0.88
1:A:584:GLU:OE2	4:A:1276:HOH:O	1.95	0.85
1:A:418:MET:HA	1:A:418:MET:HE2	1.71	0.72
1:A:575:TYR:CD2	1:A:575:TYR:C	2.69	0.70
1:A:665:THR:O	4:A:1212:HOH:O	2.10	0.70
1:A:281:GLN:OE1	4:A:1256:HOH:O	2.12	0.68
1:A:324:ARG:NE	4:A:801:HOH:O	2.14	0.66
1:A:606:ARG:O	4:A:1218:HOH:O	2.14	0.65
1:A:199:ARG:HG3	1:A:199:ARG:O	1.99	0.63
1:A:522:THR:O	1:A:525:ARG:HG2	2.00	0.62
1:A:419:TYR:CE2	1:A:460:SER:HB3	2.34	0.62
1:A:522:THR:HG23	1:A:523:THR:N	2.14	0.62
1:A:579:PRO:HD2	1:A:580:ARG:N	2.14	0.62
1:A:263:PRO:HG3	2:A:701:ADP:C6	2.35	0.60
1:A:526:ARG:HH11	1:A:526:ARG:HG3	1.67	0.60
1:A:606:ARG:HD2	4:A:1303:HOH:O	2.02	0.59
1:A:126:CYS:HB2	1:A:601:VAL:HG22	1.83	0.59
1:A:522:THR:CG2	1:A:523:THR:N	2.66	0.59
1:A:316:PHE:H	2:A:701:ADP:PB	2.27	0.58
1:A:391:GLN:HE22	1:A:418:MET:HE3	1.68	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:ASP:HB2	1:A:94:PRO:HD2	1.85	0.57
1:A:593:THR:O	1:A:604:ILE:HD12	2.05	0.57
1:A:248:LEU:HD22	1:A:606:ARG:HD3	1.88	0.56
1:A:582:VAL:HB	1:A:583:PRO:HD3	1.88	0.56
1:A:52:SER:N	1:A:53:PRO:CD	2.70	0.55
1:A:462:GLN:NE2	4:A:805:HOH:O	2.40	0.54
1:A:257:ALA:CB	1:A:299:ILE:HD11	2.38	0.54
1:A:291:THR:HG22	1:A:298:THR:OG1	2.08	0.54
1:A:522:THR:OG1	1:A:525:ARG:NH2	2.42	0.53
1:A:575:TYR:C	1:A:575:TYR:HD2	2.15	0.53
1:A:263:PRO:HG3	2:A:701:ADP:C5	2.45	0.52
1:A:632:ARG:NH2	4:A:806:HOH:O	2.42	0.51
1:A:225:LEU:HD12	1:A:225:LEU:C	2.35	0.51
1:A:640:ARG:HG3	1:A:640:ARG:NH1	2.08	0.51
1:A:588:LYS:HE2	1:A:623:HIS:HB2	1.93	0.51
1:A:579:PRO:HD2	1:A:580:ARG:H	1.75	0.50
1:A:158:LEU:HD23	1:A:158:LEU:C	2.37	0.49
1:A:126:CYS:HB2	1:A:601:VAL:CG2	2.44	0.48
1:A:199:ARG:O	1:A:199:ARG:CG	2.61	0.48
1:A:579:PRO:CD	1:A:580:ARG:N	2.77	0.47
1:A:225:LEU:HD12	1:A:225:LEU:O	2.15	0.47
1:A:257:ALA:HB1	1:A:299:ILE:HD11	1.97	0.47
1:A:324:ARG:CD	4:A:801:HOH:O	2.60	0.47
1:A:578:GLU:OE1	4:A:802:HOH:O	2.20	0.47
1:A:471:LEU:HD13	1:A:478:ILE:HD11	1.97	0.47
1:A:526:ARG:HH11	1:A:526:ARG:CG	2.27	0.47
1:A:575:TYR:CD2	1:A:575:TYR:O	2.68	0.47
1:A:544:ARG:HH11	1:A:544:ARG:HG2	1.80	0.46
1:A:659:ARG:HD3	4:A:1109:HOH:O	2.16	0.46
1:A:429:PRO:HD2	1:A:432:GLN:OE1	2.15	0.45
1:A:421:ASP:C	1:A:421:ASP:OD1	2.60	0.45
1:A:262:LEU:N	1:A:263:PRO:HD2	2.32	0.44
1:A:578:GLU:O	1:A:578:GLU:HG2	2.13	0.44
1:A:368:ILE:HD12	1:A:389:THR:HG22	2.00	0.44
1:A:239:ASP:CG	1:A:292:LEU:HB3	2.43	0.43
1:A:316:PHE:HA	2:A:701:ADP:O3A	2.19	0.43
1:A:540:THR:HA	1:A:564:VAL:HG21	2.01	0.43
1:A:147:ILE:N	1:A:148:PRO:CD	2.82	0.43
1:A:371:LEU:HD23	1:A:434:PRO:HB3	2.00	0.43
1:A:522:THR:OG1	1:A:525:ARG:CZ	2.67	0.43
1:A:265:ASP:C	1:A:265:ASP:OD1	2.60	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:525:ARG:CA	4:A:827:HOH:O	2.63	0.42
1:A:52:SER:O	1:A:53:PRO:C	2.63	0.42
1:A:494:LEU:HA	1:A:495:PRO:HD2	1.75	0.42
1:A:391:GLN:HE22	1:A:418:MET:CE	2.31	0.41
1:A:262:LEU:N	1:A:263:PRO:CD	2.84	0.41
1:A:55:MET:HG2	1:A:59:HIS:CE1	2.56	0.41
1:A:257:ALA:HB2	1:A:299:ILE:HD11	2.01	0.41
1:A:511:VAL:HB	1:A:611:SER:HB3	2.03	0.41
1:A:418:MET:HE2	1:A:418:MET:CA	2.45	0.41
1:A:316:PHE:CD1	1:A:316:PHE:N	2.87	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:874:HOH:O	4:A:874:HOH:O[15_545]	1.69	0.51

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	592/615 (96%)	576 (97%)	14 (2%)	2 (0%)	36	30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	579	PRO
1	A	458	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	506/518 (98%)	490 (97%)	16 (3%)	34	29

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	120	LEU
1	A	174	LYS
1	A	268	ASN
1	A	368	ILE
1	A	416	ASN
1	A	417	LEU
1	A	420	ARG
1	A	430	ARG
1	A	473	MET
1	A	523	THR
1	A	533	THR
1	A	546	SER
1	A	574	LEU
1	A	578	GLU
1	A	602	CYS
1	A	640	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	65	ASN
1	A	116	GLN
1	A	462	GLN
1	A	512	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ADP	A	701	3	28,29,29	1.64	4 (14%)	43,45,45	2.16	15 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	701	3	-	2/16/32/32	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	ADP	C5-C4	5.29	1.48	1.39
2	A	701	ADP	C5-C6	3.32	1.50	1.41
2	A	701	ADP	PA-O3A	3.25	1.63	1.59
2	A	701	ADP	C8-N7	2.78	1.37	1.31

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	ADP	O2A-PA-O3A	-4.79	94.32	107.27
2	A	701	ADP	C5-C4-N3	-4.41	120.64	126.72
2	A	701	ADP	N3-C2-N1	-4.29	122.09	128.58
2	A	701	ADP	O2A-PA-O1A	3.83	130.25	112.44
2	A	701	ADP	N3-C4-N9	3.73	133.52	127.17
2	A	701	ADP	C2-N1-C6	3.53	124.53	118.73
2	A	701	ADP	C2-N3-C4	3.47	120.30	111.83
2	A	701	ADP	O4'-C1'-N9	3.35	114.53	108.09
2	A	701	ADP	C4-N9-C8	3.27	109.17	105.74
2	A	701	ADP	C4-C5-N7	-3.07	107.08	110.58
2	A	701	ADP	O2B-PB-O3A	-2.42	96.52	104.64
2	A	701	ADP	C6-C5-N7	2.37	136.66	132.09
2	A	701	ADP	C5-N7-C8	2.27	107.02	103.45
2	A	701	ADP	N9-C8-N7	-2.16	110.87	113.94
2	A	701	ADP	O3B-PB-O2B	2.01	115.33	107.80

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	701	ADP	PA-O3A-PB-O2B
2	A	701	ADP	PA-O3A-PB-O3B

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	ADP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	600/615 (97%)	-0.07	39 (6%) 25 24	15, 30, 63, 114	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	532	GLY	6.6
1	A	533	THR	6.0
1	A	495	PRO	5.4
1	A	95	PHE	5.0
1	A	604	ILE	4.4
1	A	176	HIS	4.3
1	A	276	ASN	4.0
1	A	316	PHE	4.0
1	A	268	ASN	3.8
1	A	494	LEU	3.8
1	A	525	ARG	3.8
1	A	524	VAL	3.8
1	A	602	CYS	3.6
1	A	605	GLN	3.6
1	A	266	ALA	3.4
1	A	526	ARG	3.4
1	A	501	GLY	3.4
1	A	579	PRO	3.3
1	A	502	ASP	3.3
1	A	375	LYS	3.2
1	A	419	TYR	3.2
1	A	265	ASP	2.9
1	A	426	MET	2.8
1	A	523	THR	2.8
1	A	606	ARG	2.7
1	A	603	GLU	2.7
1	A	50	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	575	TYR	2.6
1	A	197	HIS	2.5
1	A	373	PHE	2.4
1	A	199	ARG	2.4
1	A	242	VAL	2.4
1	A	608	LYS	2.4
1	A	376	THR	2.4
1	A	374	GLY	2.3
1	A	274	ARG	2.2
1	A	150	ARG	2.2
1	A	416	ASN	2.1
1	A	267	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	A	702	1/1	0.89	0.09	50,50,50,50	0
2	ADP	A	701	27/27	0.92	0.11	31,44,62,74	0
3	MG	A	703	1/1	0.95	0.12	39,39,39,39	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ADP A 701:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.