



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:22 PM UTC

PDB ID : 5YJ7 / pdb_00005yj7
Title : Structural insight into the beta-GH1 glucosidase BGLN1 from oleaginous microalgae *Nannochloropsis*
Authors : Dong, S.; Liu, Y.J.; Zhou, H.X.; Xiao, Y.; Xu, J.; Cui, Q.; Wang, X.Q.; Feng, Y.G.
Deposited on : 2017-10-09
Resolution : 1.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6728 (1.64-1.60)
Clashscore	190562	7023 (1.64-1.60)
Ramachandran outliers	187476	6898 (1.64-1.60)
Sidechain outliers	187428	6896 (1.64-1.60)
RSRZ outliers	180081	6727 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	518	% 92% 7%
1	B	518	2% 93% 6%
1	C	518	% 93% 5%
1	D	518	93% 7%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 33400 atoms, of which 14858 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycoside hydrolase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	483	Total	C	H	N	O	S	0	2	0
			7549	2478	3683	645	724	19			
1	B	488	Total	C	H	N	O	S	0	6	0
			7643	2505	3734	654	730	20			
1	C	490	Total	C	H	N	O	S	0	3	0
			7645	2507	3730	654	735	19			
1	D	483	Total	C	H	N	O	S	0	1	0
			7539	2474	3679	645	722	19			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		
2	C	1	Total	Ca	0	0
			1	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	C	1	Total	C	H	O	0	0
			14	3	8	3		
3	D	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	781	Total	O	0	0
			781	781		
4	B	791	Total	O	0	0
			791	791		
4	C	652	Total	O	0	0
			652	652		
4	D	742	Total	O	0	0
			742	742		

- Molecule 1: Glycoside hydrolase



- [illegible]

- [illegible]

- MET D21 K140 E422 S503 GLY ASP THR ALA PRO PRO ASP ASP ALA GLU ALA VAL ALA ARG ASP ALA VAL LYS ALA THR PRO SER GLU ALA PRO LEU GLU HIS HIS HIS HIS HIS

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	70.72Å 74.75Å 102.94Å 105.29° 96.10° 90.32°	Depositor
Resolution (Å)	27.31 – 1.61 27.31 – 1.61	Depositor EDS
% Data completeness (in resolution range)	82.7 (27.31-1.61) 80.9 (27.31-1.61)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.33 (at 1.61Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.134 , 0.151 0.137 , 0.152	Depositor DCC
R_{free} test set	2000 reflections (0.92%)	wwPDB-VP
Wilson B-factor (Å ²)	13.5	Xtriage
Anisotropy	0.257	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 51.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	33400	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/3982	0.68	0/5404
1	B	0.43	1/4056 (0.0%)	0.68	0/5505
1	C	0.43	1/4043 (0.0%)	0.67	2/5489 (0.0%)
1	D	0.40	0/3973	0.66	0/5392
All	All	0.42	2/16054 (0.0%)	0.67	2/21790 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	136	ALA	C-O	-6.22	1.16	1.23
1	B	107	GLU	C-O	-5.60	1.17	1.24

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	507	ALA	CA-C-N	-5.12	115.10	120.38
1	C	507	ALA	C-N-CA	-5.12	115.10	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3866	3683	3694	3	0
1	B	3909	3734	3715	2	0
1	C	3915	3730	3729	5	0
1	D	3860	3679	3688	1	0
2	A	1	0	0	0	0
2	C	1	0	0	0	0
3	A	6	8	8	0	0
3	B	6	8	8	0	0
3	C	6	8	8	0	0
3	D	6	8	8	0	0
4	A	781	0	0	0	4
4	B	791	0	0	0	5
4	C	652	0	0	0	3
4	D	742	0	0	1	6
All	All	18542	14858	14858	10	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:140:LYS:NZ	4:D:709:HOH:O	2.41	0.44
1:C:345:ASN:HB3	1:C:367:ILE:HG23	2.00	0.43
1:C:507:ALA:N	1:C:508:PRO:CD	2.81	0.43
1:A:371:PRO:HB3	1:C:25:TRP:CG	2.54	0.43
1:A:68:ALA:HB1	1:A:71:GLU:CG	2.49	0.42
1:C:111:LEU:HD21	1:C:477:VAL:HG21	2.02	0.41
1:C:480:ASP:O	1:C:484:GLN:N	2.53	0.41
1:A:120:TYR:CE2	1:A:122:PHE:HB3	2.56	0.40
1:B:111:LEU:HD23	1:B:111:LEU:C	2.46	0.40
1:B:480:ASP:O	1:B:484:GLN:N	2.53	0.40

All (9) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:1074:HOH:O	4:D:804:HOH:O[1_545]	1.85	0.35
4:A:1216:HOH:O	4:B:1264:HOH:O[1_655]	1.99	0.21
4:A:1318:HOH:O	4:D:1189:HOH:O[1_455]	1.99	0.21
4:B:1133:HOH:O	4:D:1162:HOH:O[1_456]	2.04	0.16

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1381:HOH:O	4:B:1441:HOH:O[1_665]	2.05	0.15
4:C:1335:HOH:O	4:D:1368:HOH:O[1_545]	2.06	0.14
4:B:1450:HOH:O	4:D:1431:HOH:O[1_445]	2.12	0.08
4:C:1255:HOH:O	4:D:1433:HOH:O[1_545]	2.12	0.08
4:A:716:HOH:O	4:B:705:HOH:O[1_565]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	483/518 (93%)	471 (98%)	12 (2%)	0	100	100
1	B	492/518 (95%)	477 (97%)	15 (3%)	0	100	100
1	C	491/518 (95%)	476 (97%)	15 (3%)	0	100	100
1	D	482/518 (93%)	469 (97%)	13 (3%)	0	100	100
All	All	1948/2072 (94%)	1893 (97%)	55 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	394/418 (94%)	394 (100%)	0	100	100
1	B	401/418 (96%)	401 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	400/418 (96%)	400 (100%)	0	100	100
1	D	393/418 (94%)	393 (100%)	0	100	100
All	All	1588/1672 (95%)	1588 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	409	ASN
1	B	196	GLN
1	B	409	ASN
1	B	478	GLN
1	C	325	GLN
1	C	409	ASN
1	D	103	HIS
1	D	196	GLN
1	D	325	GLN
1	D	328	GLN
1	D	409	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	D	601	-	5,5,5	0.44	0	5,5,5	0.55	0
3	GOL	B	601	-	5,5,5	0.47	0	5,5,5	0.54	0
3	GOL	A	602	-	5,5,5	0.43	0	5,5,5	0.49	0
3	GOL	C	602	-	5,5,5	0.42	0	5,5,5	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	D	601	-	-	0/4/4/4	-
3	GOL	B	601	-	-	0/4/4/4	-
3	GOL	A	602	-	-	0/4/4/4	-
3	GOL	C	602	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	483/518 (93%)	-0.74	3 (0%) 85 89	7, 17, 39, 98	2 (0%)
1	B	488/518 (94%)	-0.70	8 (1%) 70 75	8, 17, 44, 93	3 (0%)
1	C	490/518 (94%)	-0.47	7 (1%) 73 78	9, 22, 50, 87	2 (0%)
1	D	483/518 (93%)	-0.74	2 (0%) 88 91	9, 18, 38, 75	1 (0%)
All	All	1944/2072 (93%)	-0.66	20 (1%) 79 83	7, 18, 44, 98	8 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	509	PRO	4.8
1	A	503	SER	4.4
1	B	22	VAL	4.1
1	B	507	ALA	4.0
1	B	420	THR	3.9
1	B	421	PRO	3.1
1	C	21	ASP	3.0
1	D	422	GLU	2.9
1	B	508	PRO	2.6
1	C	501[A]	LYS	2.5
1	D	21	ASP	2.5
1	C	352	GLU	2.5
1	C	22	VAL	2.4
1	A	352	GLU	2.4
1	C	366	TRP	2.4
1	B	501[A]	LYS	2.4
1	A	21	ASP	2.3
1	C	348	GLN	2.2
1	B	422	GLU	2.2
1	C	503	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	D	601	6/6	0.92	0.10	20,30,38,38	0
3	GOL	B	601	6/6	0.95	0.13	16,22,32,32	0
3	GOL	A	602	6/6	0.95	0.12	17,23,33,33	0
3	GOL	C	602	6/6	0.97	0.07	19,26,33,33	0
2	CA	C	601	1/1	0.97	0.11	26,26,26,26	0
2	CA	A	601	1/1	0.98	0.04	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.