



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 05:42 AM UTC

PDB ID : 8YJM / pdb_00008yjm
Title : Structure of human SPT16 MD-CTD and MCM2 HBD chaperoning a histone
H3-H4 tetramer and a single chain H2B-H2A chimera
Authors : Gan, S.L.; Yang, W.S.; Xu, R.M.
Deposited on : 2024-03-02
Resolution : 4.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1077 (4.50-3.82)
Clashscore	190562	1019 (4.48-3.84)
Ramachandran outliers	187476	1035 (4.50-3.82)
Sidechain outliers	187428	1022 (4.50-3.82)
RSRZ outliers	180081	1074 (4.50-3.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	350	
2	B	93	
3	C	81	
3	E	81	
4	D	103	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	103	 63% 9% 28%
5	G	213	 73% 12% 15%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6879 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FACT complex subunit SPT16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	310	Total	C	N	O	S	0	0	0
			2549	1622	422	493	12			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	639	GLY	-	expression tag	UNP Q9Y5B9
A	640	PRO	-	expression tag	UNP Q9Y5B9
A	641	LEU	-	expression tag	UNP Q9Y5B9
A	642	GLY	-	expression tag	UNP Q9Y5B9
A	643	SER	-	expression tag	UNP Q9Y5B9

- Molecule 2 is a protein called DNA replication licensing factor MCM2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	58	Total	C	N	O	S	0	0	0
			455	268	81	104	2			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	62	SER	-	expression tag	UNP P49736

- Molecule 3 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	79	Total	C	N	O	S	0	0	0
			641	405	119	113	4			
3	E	80	Total	C	N	O	S	0	0	0
			646	408	120	114	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	55	MET	-	initiating methionine	UNP P68431
E	55	MET	-	initiating methionine	UNP P68431

- Molecule 4 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	71	Total	C	N	O	S	0	0	0
			572	359	113	99	1			
4	F	74	Total	C	N	O	S	0	0	0
			591	374	115	101	1			




- Molecule 5 is a protein called Histone H2B 1/2/3/4/6,Histone H2A type 1-D.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	G	182	Total	C	N	O	S	0	2	0
			1425	898	264	261	2			

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
G	12	MET	-	initiating methionine	UNP P0C1H3
G	13	GLY	-	expression tag	UNP P0C1H3
G	14	SER	-	expression tag	UNP P0C1H3
G	15	SER	-	expression tag	UNP P0C1H3
G	16	HIS	-	expression tag	UNP P0C1H3
G	17	HIS	-	expression tag	UNP P0C1H3
G	18	HIS	-	expression tag	UNP P0C1H3
G	19	HIS	-	expression tag	UNP P0C1H3
G	20	HIS	-	expression tag	UNP P0C1H3
G	21	HIS	-	expression tag	UNP P0C1H3
G	22	SER	-	expression tag	UNP P0C1H3
G	23	SER	-	expression tag	UNP P0C1H3
G	24	GLY	-	expression tag	UNP P0C1H3
G	25	LEU	-	expression tag	UNP P0C1H3
G	26	VAL	-	expression tag	UNP P0C1H3
G	27	PRO	-	expression tag	UNP P0C1H3
G	28	ARG	-	expression tag	UNP P0C1H3
G	29	GLY	-	expression tag	UNP P0C1H3
G	30	SER	-	expression tag	UNP P0C1H3
G	31	HIS	-	expression tag	UNP P0C1H3
G	32	MET	-	expression tag	UNP P0C1H3

- Molecule 1: FACT complex subunit SPT16






● Molecule 4: Histone H4



● Molecule 5: Histone H2B 1/2/3/4/6,Histone H2A type 1-D



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	191.34Å 191.34Å 80.09Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	165.70 – 4.15 165.70 – 4.15	Depositor EDS
% Data completeness (in resolution range)	99.8 (165.70-4.15) 96.1 (165.70-4.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	497.77 (at 4.15Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.263 , 0.286 (Not available) , 0.261	Depositor DCC
R_{free} test set	1269 reflections (9.87%)	wwPDB-VP
Wilson B-factor (Å ²)	154.0	Xtriage
Anisotropy	0.155	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 189.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.041 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6879	wwPDB-VP
Average B, all atoms (Å ²)	200.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.11	0/2607	0.29	0/3522
2	B	0.11	0/457	0.31	0/612
3	C	0.11	0/648	0.28	0/868
3	E	0.11	0/653	0.30	0/875
4	D	0.10	0/577	0.26	0/773
4	F	0.12	0/597	0.26	0/800
5	G	0.12	0/1451	0.27	0/1954
All	All	0.11	0/6990	0.28	0/9404

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2549	0	2463	33	0
2	B	455	0	411	13	0
3	C	641	0	674	9	0
3	E	646	0	679	7	0
4	D	572	0	615	9	0
4	F	591	0	638	10	0
5	G	1425	0	1497	20	0
All	All	6879	0	6977	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 77 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:660:PRO:HG2	1:A:682:ALA:HB3	1.53	0.88
3:C:59:GLU:HB3	4:D:40:ARG:HH22	1.47	0.76
1:A:750:ASP:HB3	4:D:45:ARG:HG3	1.77	0.67
4:D:92:ARG:HH22	5:G:101:LEU:HD23	1.61	0.65
4:D:74:GLU:OE2	5:G:99:ARG:NH1	2.32	0.62

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/350 (68%)	230 (96%)	9 (4%)	0	100	100
2	B	56/93 (60%)	55 (98%)	1 (2%)	0	100	100
3	C	77/81 (95%)	77 (100%)	0	0	100	100
3	E	35/81 (43%)	34 (97%)	1 (3%)	0	100	100
4	D	69/103 (67%)	69 (100%)	0	0	100	100
4	F	72/103 (70%)	69 (96%)	3 (4%)	0	100	100
5	G	158/213 (74%)	156 (99%)	2 (1%)	0	100	100
All	All	706/1024 (69%)	690 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/315 (90%)	285 (100%)	0	100	100
2	B	44/75 (59%)	44 (100%)	0	100	100
3	C	69/70 (99%)	69 (100%)	0	100	100
3	E	69/70 (99%)	69 (100%)	0	100	100
4	D	60/79 (76%)	60 (100%)	0	100	100
4	F	61/79 (77%)	61 (100%)	0	100	100
5	G	152/176 (86%)	152 (100%)	0	100	100
All	All	740/864 (86%)	740 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
5	G	1038	ASN
5	G	1024	GLN
3	E	108	ASN
3	C	85	GLN
5	G	95	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	310/350 (88%)	-0.36	0 100 100	116, 188, 262, 306	0
2	B	58/93 (62%)	-0.29	0 100 100	157, 251, 369, 409	0
3	C	79/81 (97%)	-0.44	0 100 100	141, 177, 228, 247	0
3	E	80/81 (98%)	-0.41	0 100 100	125, 167, 234, 244	0
4	D	71/103 (68%)	-0.47	0 100 100	153, 195, 226, 236	0
4	F	74/103 (71%)	-0.16	0 100 100	135, 187, 240, 262	0
5	G	182/213 (85%)	-0.32	0 100 100	131, 218, 258, 301	2 (1%)
All	All	854/1024 (83%)	-0.35	0 100 100	116, 200, 272, 409	2 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.