



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 19, 2026 – 07:31 AM UTC

PDB ID : 7YKE / pdb_00007yke
Title : Crystal structure of chondroitin ABC lyase I in complex with chondroitin disaccharide 4,6-sulfate
Authors : Takashima, M.; Watanabe, I.; Miyanaga, A.; Eguchi, T.
Deposited on : 2022-07-22
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1220 (1.88-1.88)
Clashscore	190562	1234 (1.88-1.88)
Ramachandran outliers	187476	1222 (1.88-1.88)
Sidechain outliers	187428	1222 (1.88-1.88)
RSRZ outliers	180081	1220 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1021	4% 84% 9% 5%
2	B	2	100%
2	C	2	50% 50%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 8530 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chondroitin sulfate ABC endolyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	966	Total	C	N	O	S	0	0	0
			7720	4913	1317	1469	21			

- Molecule 2 is an oligosaccharide called 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-4,6-di-O-sulfo-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	S	0	0	0
			34	14	1	17	2			
2	C	2	Total	C	N	O	S	0	0	0
			34	14	1	17	2			

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	741	Total	O	0	0
			741	741		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chondroitin sulfate ABC endolyase



- Molecule 2: 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-4,6-di-O-sulfo-beta-D-galactopyranose



- Molecule 2: 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-4,6-di-O-sulfo-beta-D-galactopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.07Å 94.50Å 229.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.07 – 1.88 49.07 – 1.88	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.07-1.88) 100.0 (49.07-1.88)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 1.88Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.169 , 0.213 0.180 , 0.219	Depositor DCC
R_{free} test set	4380 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	22.4	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8530	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GCD, MG, 8EX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.18	12/7909 (0.2%)	1.36	20/10726 (0.2%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	831	GLN	C-O	8.90	1.34	1.23
1	A	307	HIS	C-O	6.71	1.31	1.23
1	A	389	HIS	CE1-NE2	6.59	1.39	1.32
1	A	389	HIS	CG-ND1	6.57	1.45	1.38
1	A	148	ASP	C-O	-6.05	1.17	1.23
1	A	833	GLU	C-O	5.88	1.31	1.24
1	A	938	VAL	C-O	5.70	1.29	1.24
1	A	750	PRO	C-O	-5.66	1.17	1.24
1	A	837	VAL	C-O	5.33	1.29	1.24
1	A	372	HIS	CE1-NE2	5.21	1.37	1.32
1	A	790	ILE	C-O	5.10	1.29	1.24
1	A	45	ALA	C-O	5.09	1.29	1.23

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	906	ASP	CA-CB-CG	7.63	120.23	112.60
1	A	484	VAL	CA-C-O	7.41	123.27	119.12
1	A	332	ASN	CA-CB-CG	-7.10	105.50	112.60
1	A	599	ASP	CA-CB-CG	-7.00	105.59	112.60
1	A	131	GLU	O-C-N	6.72	130.87	123.27
1	A	485	PRO	O-C-N	6.31	124.21	121.31
1	A	599	ASP	CB-CA-C	-5.78	103.77	111.63
1	A	626	ASN	CA-CB-CG	-5.68	106.92	112.60
1	A	634	ARG	NE-CZ-NH2	5.64	124.28	119.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	252	GLU	CB-CG-CD	5.61	122.14	112.60
1	A	634	ARG	NE-CZ-NH1	-5.55	115.94	121.50
1	A	131	GLU	CB-CA-C	5.45	118.75	109.75
1	A	612	THR	CA-CB-OG1	-5.25	101.72	109.60
1	A	830	THR	CB-CA-C	5.22	117.84	109.07
1	A	338	ASN	N-CA-C	-5.19	105.53	111.14
1	A	81	GLY	CA-C-O	-5.14	116.89	121.41
1	A	35	ASN	CA-C-N	5.12	127.40	120.38
1	A	35	ASN	C-N-CA	5.12	127.40	120.38
1	A	205	GLN	CA-C-O	-5.12	115.93	121.36
1	A	134	ILE	CA-C-O	-5.03	116.41	121.49

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7720	0	7558	39	0
2	B	34	0	5	0	0
2	C	34	0	5	2	0
3	A	1	0	0	0	0
4	A	741	0	0	5	0
All	All	8530	0	7568	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:276:ASN:ND2	2:C:1:8EX:O1S	1.99	0.95
1:A:812:THR:HG22	1:A:836:ASN:ND2	2.09	0.67
1:A:976:ASN:HB3	4:A:1743:HOH:O	1.95	0.65
1:A:57:LYS:HG2	1:A:58:ASN:ND2	2.14	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:812:THR:HG22	1:A:836:ASN:HD22	1.64	0.62
1:A:388:HIS:ND1	1:A:389:HIS:ND1	2.41	0.59
1:A:276:ASN:CG	2:C:1:8EX:O1S	2.46	0.58
1:A:999:ASP:HB3	4:A:1339:HOH:O	2.04	0.57
1:A:282:ASN:OD1	1:A:284:SER:HB2	2.07	0.55
1:A:450:SER:HB2	1:A:519:LEU:HD11	1.89	0.54
1:A:829:ILE:HD13	1:A:835:VAL:HG21	1.93	0.51
1:A:32:ASP:O	1:A:35:ASN:N	2.39	0.50
1:A:806:ASN:HB2	4:A:1521:HOH:O	2.10	0.49
1:A:58:ASN:HB2	1:A:83:SER:HB2	1.94	0.49
1:A:79:TRP:CE3	1:A:199:ALA:HB1	2.48	0.48
1:A:97:LYS:HG2	4:A:1512:HOH:O	2.13	0.47
1:A:129:PHE:O	1:A:140:GLN:HB3	2.15	0.46
1:A:273:LYS:HD2	1:A:273:LYS:HA	1.72	0.46
1:A:388:HIS:HD1	1:A:389:HIS:CE1	2.29	0.46
1:A:734:MET:HA	1:A:758:VAL:O	2.16	0.45
1:A:257:ILE:HG21	1:A:466:ARG:HD2	1.99	0.45
1:A:56:ASP:OD1	1:A:56:ASP:C	2.59	0.44
1:A:798:TRP:O	1:A:819:LEU:HA	2.18	0.44
1:A:995:GLN:NE2	1:A:996:VAL:O	2.48	0.43
1:A:26:THR:O	1:A:89:LYS:NZ	2.52	0.43
1:A:555:LEU:N	1:A:556:PRO:CD	2.82	0.42
1:A:451:ARG:HD2	1:A:518:GLN:OE1	2.20	0.42
1:A:115:LEU:HA	1:A:209:TYR:O	2.19	0.41
1:A:826:GLY:O	1:A:877:VAL:HA	2.20	0.41
1:A:283:ILE:CD1	1:A:349:ALA:HB1	2.50	0.41
1:A:832:ALA:C	4:A:1599:HOH:O	2.63	0.41
1:A:103:TRP:HE3	1:A:107:SER:OG	2.04	0.41
1:A:201:SER:C	1:A:203:VAL:H	2.28	0.41
1:A:33:PRO:HD3	1:A:92:ILE:HD13	2.03	0.41
1:A:622:PHE:CG	1:A:733:GLY:HA3	2.56	0.41
1:A:769:GLY:HA3	1:A:873:TYR:CE2	2.56	0.40
1:A:323:ASN:OD1	1:A:323:ASN:C	2.63	0.40
1:A:499:TRP:CH2	1:A:964:MET:HE1	2.56	0.40
1:A:57:LYS:CG	1:A:58:ASN:ND2	2.83	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	956/1021 (94%)	917 (96%)	37 (4%)	2 (0%)	43	34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	202	ASN
1	A	48	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	847/890 (95%)	821 (97%)	26 (3%)	35	18

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	LYS
1	A	36	LEU
1	A	50	LEU
1	A	54	SER
1	A	57	LYS
1	A	58	ASN
1	A	62	THR
1	A	72	ASN
1	A	90	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	101	LYS
1	A	127	ILE
1	A	140	GLN
1	A	144	LYS
1	A	188	LEU
1	A	191	LYS
1	A	204	SER
1	A	212	ARG
1	A	230	LYS
1	A	234	SER
1	A	273	LYS
1	A	332	ASN
1	A	405	SER
1	A	451	ARG
1	A	599	ASP
1	A	741	TYR
1	A	819	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	GLN
1	A	46	GLN
1	A	58	ASN
1	A	140	GLN
1	A	296	HIS
1	A	325	GLN
1	A	416	GLN
1	A	464	GLN
1	A	468	ASN
1	A	573	GLN
1	A	636	GLN
1	A	648	ASN
1	A	731	GLN
1	A	744	ASN
1	A	771	ASN
1	A	823	ASN
1	A	836	ASN
1	A	840	GLN
1	A	841	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	8EX	B	1	2	23,23,23	1.95	7 (30%)	31,35,35	3.40	11 (35%)
2	GCD	B	2	2	10,11,12	2.28	3 (30%)	12,15,17	1.88	4 (33%)
2	8EX	C	1	2	23,23,23	1.73	5 (21%)	31,35,35	3.68	9 (29%)
2	GCD	C	2	2	10,11,12	2.45	2 (20%)	12,15,17	2.56	6 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8EX	B	1	2	-	9/15/35/35	0/1/1/1
2	GCD	B	2	2	-	0/4/17/20	0/1/1/1
2	8EX	C	1	2	-	4/15/35/35	0/1/1/1
2	GCD	C	2	2	-	4/4/17/20	0/1/1/1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	GCD	O5-C5	6.05	1.45	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	GCD	O5-C5	5.38	1.44	1.37
2	C	1	8EX	O5-C5	4.59	1.55	1.44
2	C	1	8EX	C3-C2	4.14	1.60	1.53
2	B	1	8EX	O1S-S1	4.13	1.63	1.45
2	B	1	8EX	O1-C1	-3.91	1.27	1.39
2	B	1	8EX	O3S-S1	3.72	1.61	1.45
2	B	2	GCD	C4-C5	3.49	1.39	1.33
2	C	2	GCD	C5-C6	-3.35	1.41	1.48
2	B	1	8EX	C3-C2	-2.84	1.47	1.53
2	B	1	8EX	O3-C3	-2.74	1.36	1.43
2	B	2	GCD	C3-C4	2.68	1.53	1.50
2	C	1	8EX	O3S-S1	2.64	1.56	1.45
2	C	1	8EX	O6-S1	-2.50	1.50	1.56
2	B	1	8EX	O5-C1	-2.46	1.36	1.42
2	B	1	8EX	C1-C2	2.22	1.55	1.52
2	C	1	8EX	O6S-S2	2.19	1.54	1.45

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	8EX	C1-O5-C5	-11.23	91.93	113.65
2	C	1	8EX	C1-C2-C3	-11.08	95.44	110.54
2	B	1	8EX	C1-C2-C3	-9.74	97.26	110.54
2	B	1	8EX	O6-S1-O3S	-7.69	83.45	106.92
2	B	1	8EX	O2S-S1-O6	7.43	123.44	106.37
2	B	1	8EX	O2S-S1-O3S	-7.01	84.03	108.56
2	C	1	8EX	O1-C1-C2	5.88	121.44	109.22
2	C	1	8EX	O2S-S1-O3S	5.44	127.60	108.56
2	C	1	8EX	O6-C6-C5	-5.28	98.16	107.57
2	C	1	8EX	O1-C1-O5	4.78	124.60	110.41
2	B	1	8EX	C1-O5-C5	-4.35	105.24	113.65
2	C	2	GCD	C1-C2-C3	4.31	115.92	109.64
2	C	2	GCD	C4-C5-C6	-4.21	114.44	123.56
2	B	1	8EX	O2S-S1-O1S	3.90	122.21	108.56
2	B	1	8EX	C1-C2-N2	-3.49	106.68	110.73
2	C	2	GCD	O5-C5-C6	3.40	118.10	111.85
2	C	2	GCD	O6A-C6-C5	-3.38	112.28	120.57
2	C	1	8EX	O3-C3-C2	3.37	116.30	109.58
2	B	2	GCD	O6B-C6-C5	3.20	121.32	114.06
2	B	1	8EX	O5-C1-C2	-2.99	106.51	109.52
2	B	2	GCD	O2-C2-C1	-2.96	102.46	109.22
2	C	1	8EX	O2S-S1-O6	-2.95	99.59	106.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	GCD	O5-C5-C4	-2.65	122.51	124.94
2	C	2	GCD	O6B-C6-O6A	2.60	130.08	123.90
2	C	2	GCD	O5-C5-C4	2.59	127.31	124.94
2	B	1	8EX	O3S-S1-O1S	-2.57	102.34	112.24
2	B	1	8EX	O6-S1-O1S	2.43	114.34	106.92
2	B	1	8EX	O3-C3-C4	2.27	115.75	109.94
2	B	2	GCD	C1-C2-C3	2.07	112.66	109.64
2	C	1	8EX	O6-S1-O3S	-2.00	100.80	106.92

There are no chirality outliers.

All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1	8EX	C4-O4-S2-O6S
2	B	1	8EX	C4-O4-S2-O4S
2	C	1	8EX	C4-C5-C6-O6
2	C	1	8EX	O5-C5-C6-O6
2	C	2	GCD	C4-C5-C6-O6A
2	C	2	GCD	C4-C5-C6-O6B
2	C	2	GCD	O5-C5-C6-O6A
2	C	2	GCD	O5-C5-C6-O6B
2	B	1	8EX	C8-C7-N2-C2
2	B	1	8EX	O7-C7-N2-C2
2	B	1	8EX	C4-O4-S2-O5S
2	B	1	8EX	C6-O6-S1-O2S
2	B	1	8EX	C6-O6-S1-O3S
2	B	1	8EX	C5-C6-O6-S1
2	C	1	8EX	C1-C2-N2-C7
2	C	1	8EX	C6-O6-S1-O3S
2	B	1	8EX	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	8EX	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	966/1021 (94%)	-0.02	43 (4%)	38 41	14, 23, 50, 72	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	77	TRP	4.0
1	A	203	VAL	3.9
1	A	51	ALA	3.5
1	A	392	TYR	3.2
1	A	204	SER	3.2
1	A	201	SER	3.2
1	A	63	LEU	3.1
1	A	299	ALA	3.1
1	A	188	LEU	2.9
1	A	61	LEU	2.8
1	A	34	LYS	2.8
1	A	189	GLY	2.8
1	A	64	SER	2.7
1	A	134	ILE	2.7
1	A	202	ASN	2.7
1	A	985	SER	2.6
1	A	273	LYS	2.6
1	A	137	SER	2.6
1	A	33	PRO	2.6
1	A	50	LEU	2.6
1	A	85	PHE	2.5
1	A	79	TRP	2.5
1	A	301	GLY	2.5
1	A	832	ALA	2.5
1	A	80	LYS	2.5
1	A	951	ASP	2.4
1	A	49	PRO	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	62	THR	2.3
1	A	121	ILE	2.3
1	A	191	LYS	2.3
1	A	35	ASN	2.2
1	A	136	THR	2.2
1	A	205	GLN	2.2
1	A	135	SER	2.2
1	A	206	GLY	2.2
1	A	60	ILE	2.2
1	A	192	VAL	2.2
1	A	76	LEU	2.2
1	A	57	LYS	2.1
1	A	45	ALA	2.1
1	A	806	ASN	2.0
1	A	93	VAL	2.0
1	A	133	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	8EX	B	1	23/23	0.87	0.11	32,40,55,63	0
2	8EX	C	1	23/23	0.88	0.10	22,37,56,63	0
2	GCD	C	2	11/12	0.89	0.09	36,40,51,56	0
2	GCD	B	2	11/12	0.96	0.06	23,29,31,32	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	A	1201	1/1	0.92	0.11	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.