



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:48 PM UTC

PDB ID : 1YLV / pdb_00001ylv
Title : SCHIFF-BASE COMPLEX OF YEAST 5-AMINOLAEVULINIC ACID DEHYDRATASE WITH LAEVULINIC ACID
Authors : Erskine, P.T.; Newbold, R.; Roper, J.; Coker, A.; Warren, M.J.; Shoolingin-Jordan, P.M.; Wood, S.P.; Cooper, J.B.
Deposited on : 1999-02-22
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3111 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROTEIN (5-AMINOLAEVULINIC ACID DEHYDRATASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	341	Total	C	N	O	S	0	1	0
			2650	1685	459	491	15			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is LAEVULINIC ACID (CCD ID: SHF) (formula: C₅H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	1
			14	10	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	446	Total 446	O 446	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROTEIN (5-AMINOLAEVULINIC ACID DEHYDRATASE)



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	103.40Å 103.40Å 168.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.15 25.00 – 2.15	Depositor EDS
% Data completeness (in resolution range)	96.8 (25.00-2.15) 96.7 (25.00-2.15)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.31 (at 2.15Å)	Xtriage
Refinement program	CCP4	Depositor
R, R_{free}	0.207 , 0.265 0.184 , 0.229	Depositor DCC
R_{free} test set	1236 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtriage
Anisotropy	0.374	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 95.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3111	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SHF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	1/2709 (0.0%)	1.63	47/3674 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	207	TYR	CA-C	5.62	1.59	1.53

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	PHE	CA-C-N	9.89	134.91	120.87
1	A	54	PHE	C-N-CA	9.89	134.91	120.87
1	A	227	PRO	CA-C-N	9.53	136.39	122.86
1	A	227	PRO	C-N-CA	9.53	136.39	122.86
1	A	224	CYS	CA-C-N	8.95	137.75	122.26
1	A	224	CYS	C-N-CA	8.95	137.75	122.26
1	A	17	VAL	N-CA-C	8.79	119.85	107.37
1	A	110	GLY	CA-C-N	8.60	128.18	119.24
1	A	110	GLY	C-N-CA	8.60	128.18	119.24
1	A	66	ILE	N-CA-C	7.18	118.63	108.36
1	A	326	ILE	CA-C-N	-6.96	113.67	122.37
1	A	326	ILE	C-N-CA	-6.96	113.67	122.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	237	LEU	CA-C-N	-6.89	114.70	119.66
1	A	237	LEU	C-N-CA	-6.89	114.70	119.66
1	A	236	GLN	CB-CG-CD	6.63	123.88	112.60
1	A	203	PHE	CA-C-N	-6.34	114.49	123.11
1	A	203	PHE	C-N-CA	-6.34	114.49	123.11
1	A	49	ASP	CA-CB-CG	-6.25	106.35	112.60
1	A	207	TYR	O-C-N	6.08	129.70	122.46
1	A	305	VAL	CA-C-O	-6.08	115.68	121.64
1	A	70	ARG	CA-C-N	6.05	128.67	120.44
1	A	70	ARG	C-N-CA	6.05	128.67	120.44
1	A	48	SER	CA-C-N	5.87	129.23	120.31
1	A	48	SER	C-N-CA	5.87	129.23	120.31
1	A	53	ASP	CA-C-N	5.79	132.59	121.54
1	A	53	ASP	C-N-CA	5.79	132.59	121.54
1	A	235	TYR	CA-C-N	-5.78	113.16	121.42
1	A	235	TYR	C-N-CA	-5.78	113.16	121.42
1	A	60	LEU	CA-C-N	5.77	127.05	119.84
1	A	60	LEU	C-N-CA	5.77	127.05	119.84
1	A	250	GLU	CA-CB-CG	5.72	125.55	114.10
1	A	230	GLY	N-CA-C	5.61	126.48	113.18
1	A	214	ASN	CA-CB-CG	5.61	118.21	112.60
1	A	128	ILE	N-CA-C	5.54	116.08	107.99
1	A	286	ALA	CA-C-N	-5.43	115.40	123.11
1	A	286	ALA	C-N-CA	-5.43	115.40	123.11
1	A	207	TYR	CA-C-O	-5.29	116.28	122.37
1	A	289	VAL	N-CA-C	5.27	118.11	109.78
1	A	221	ASP	CA-C-N	5.24	128.27	120.31
1	A	221	ASP	C-N-CA	5.24	128.27	120.31
1	A	207	TYR	CB-CA-C	-5.14	104.01	112.03
1	A	1	MET	CB-CG-SD	-5.11	97.36	112.70
1	A	292	GLU	CA-C-N	5.11	127.39	120.44
1	A	292	GLU	C-N-CA	5.11	127.39	120.44
1	A	129	ILE	CA-C-N	-5.08	115.83	122.99
1	A	129	ILE	C-N-CA	-5.08	115.83	122.99
1	A	324	ARG	NE-CZ-NH1	-5.04	116.46	121.50

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	263[B]	LYS	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2650	0	2638	85	1
2	A	1	0	0	0	0
3	A	14	0	14	0	0
4	A	446	0	0	11	5
All	All	3111	0	2652	85	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:228:SER:O	1:A:229:ASN:HB2	1.68	0.91
1:A:228:SER:O	1:A:229:ASN:CB	2.22	0.87
1:A:250:GLU:HG3	4:A:545:HOH:O	1.76	0.85
1:A:65:ARG:HD2	4:A:859:HOH:O	1.77	0.83
1:A:208:ALA:CB	1:A:260:ILE:HD12	2.11	0.80
1:A:253:MET:HG3	1:A:282:LEU:HD11	1.66	0.78
1:A:224:CYS:SG	4:A:931:HOH:O	2.41	0.77
1:A:208:ALA:HB2	1:A:260:ILE:CD1	2.17	0.74
1:A:57:ILE:HD13	1:A:222:ALA:CB	2.21	0.70
1:A:216:TYR:CE1	1:A:263[A]:LYS:HE2	2.27	0.70
1:A:113:ILE:O	1:A:117:LYS:HG3	1.92	0.68
1:A:99:ASP:HB2	1:A:100:PRO:CD	2.25	0.67
1:A:235:TYR:CD1	1:A:235:TYR:N	2.63	0.67
1:A:57:ILE:HG21	1:A:222:ALA:HB2	1.78	0.65
1:A:216:TYR:CE1	1:A:263[B]:LYS:HE2	2.32	0.65
1:A:208:ALA:HB3	1:A:260:ILE:HG23	1.77	0.65
1:A:181:MET:HE2	1:A:237:LEU:HG	1.79	0.65
1:A:49:ASP:HB2	1:A:93:LEU:H	1.60	0.64
1:A:208:ALA:CB	1:A:260:ILE:CD1	2.74	0.64
1:A:212:SER:HA	1:A:236:GLN:HE21	1.63	0.63
1:A:227:PRO:HD3	1:A:232:ARG:HH21	1.64	0.62
1:A:158:VAL:HG13	1:A:185:ARG:HA	1.82	0.62
1:A:179:SER:HB3	1:A:207:TYR:CE1	2.38	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:334:PHE:O	1:A:338:LEU:HG	2.04	0.57
1:A:177:ALA:HA	1:A:205:LEU:O	2.04	0.57
1:A:57:ILE:HD13	1:A:222:ALA:HB1	1.86	0.57
1:A:277:GLU:HB3	4:A:689:HOH:O	2.05	0.57
1:A:208:ALA:HB3	1:A:260:ILE:HD12	1.86	0.56
1:A:92:PRO:HB2	1:A:97:THR:HG21	1.88	0.56
1:A:208:ALA:HB2	1:A:260:ILE:HD13	1.87	0.56
1:A:155:GLU:O	1:A:159:SER:HB2	2.06	0.55
1:A:298:ALA:HA	1:A:301:GLU:HG2	1.88	0.55
1:A:125:GLU:HG2	4:A:920:HOH:O	2.08	0.54
1:A:227:PRO:HG3	1:A:232:ARG:HE	1.73	0.53
1:A:250:GLU:CB	4:A:545:HOH:O	2.57	0.53
1:A:316:THR:HG21	4:A:755:HOH:O	2.08	0.52
1:A:232:ARG:HD3	1:A:235:TYR:OH	2.11	0.51
1:A:250:GLU:CG	4:A:545:HOH:O	2.47	0.50
1:A:34:ARG:HH11	1:A:34:ARG:HG2	1.77	0.50
1:A:162:ALA:HA	1:A:189:ILE:HG12	1.94	0.50
1:A:216:TYR:HE1	1:A:263[A]:LYS:HE2	1.73	0.50
1:A:213:GLY:HA3	4:A:835:HOH:O	2.10	0.50
1:A:49:ASP:CB	1:A:93:LEU:H	2.24	0.50
1:A:99:ASP:HB2	1:A:100:PRO:HD3	1.91	0.50
1:A:75:LEU:O	1:A:79:VAL:HG23	2.11	0.49
1:A:179:SER:HB3	1:A:207:TYR:CZ	2.47	0.49
1:A:207:TYR:HA	1:A:261:ILE:HB	1.94	0.49
1:A:142:HIS:NE2	1:A:229:ASN:HB3	2.27	0.49
1:A:72:LYS:HE3	1:A:76:LYS:NZ	2.28	0.49
1:A:68:VAL:HG23	4:A:574:HOH:O	2.13	0.48
1:A:210:LYS:HG3	1:A:263[B]:LYS:HD3	1.96	0.48
1:A:49:ASP:CG	1:A:92:PRO:HA	2.38	0.48
1:A:227:PRO:CD	1:A:232:ARG:HH21	2.26	0.48
1:A:333:GLU:OE1	1:A:333:GLU:N	2.46	0.47
1:A:270:ASP:OD1	1:A:270:ASP:N	2.43	0.46
1:A:200:HIS:CE1	1:A:201:LYS:HD2	2.51	0.46
1:A:51:PRO:O	1:A:69:ASN:HB2	2.16	0.46
1:A:250:GLU:HB2	4:A:545:HOH:O	2.15	0.46
1:A:220:ARG:CG	1:A:220:ARG:HH11	2.29	0.46
1:A:72:LYS:HB2	1:A:118:PHE:HZ	1.81	0.45
1:A:158:VAL:CG1	1:A:188:ASP:HB2	2.46	0.45
1:A:152:ILE:HD12	1:A:235:TYR:CD1	2.52	0.45
1:A:263[A]:LYS:HA	1:A:264:PRO:C	2.40	0.45
1:A:39:ASN:HD22	1:A:39:ASN:C	2.25	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:ILE:HD13	1:A:222:ALA:HB2	1.98	0.45
1:A:260:ILE:HG22	1:A:261:ILE:N	2.31	0.44
1:A:182:ILE:O	1:A:185:ARG:HD2	2.17	0.44
1:A:216:TYR:HE1	1:A:263[B]:LYS:HE2	1.82	0.44
1:A:273:ARG:NH1	1:A:276:SER:OG	2.50	0.44
1:A:218:PRO:HB2	1:A:290:SER:HB2	2.00	0.43
1:A:302:LYS:O	1:A:302:LYS:HG3	2.19	0.43
1:A:265:SER:HB2	1:A:318:PHE:CE1	2.53	0.43
1:A:140:HIS:O	1:A:228:SER:HB2	2.19	0.42
1:A:185:ARG:HG2	1:A:186:ILE:N	2.34	0.42
1:A:260:ILE:CG2	1:A:261:ILE:N	2.82	0.42
1:A:78:LEU:HD23	1:A:78:LEU:HA	1.86	0.42
1:A:166:VAL:O	1:A:170:LYS:HG3	2.20	0.42
1:A:54:PHE:CD1	1:A:66:ILE:HG12	2.55	0.41
1:A:94:ILE:HG22	1:A:95:PRO:O	2.20	0.41
1:A:210:LYS:HA	1:A:263[A]:LYS:O	2.21	0.41
1:A:249:LEU:CD2	1:A:260:ILE:HG21	2.51	0.41
1:A:253:MET:HG3	1:A:282:LEU:CD1	2.45	0.41
1:A:340:GLU:H	1:A:340:GLU:HG3	1.52	0.41
1:A:43:PHE:HA	1:A:44:PRO:HD3	1.85	0.41
1:A:324:ARG:HH11	1:A:324:ARG:HD2	1.62	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:677:HOH:O	4:A:677:HOH:O[16_555]	0.76	1.44
1:A:125:GLU:OE1	4:A:610:HOH:O[5_555]	1.91	0.29
4:A:643:HOH:O	4:A:643:HOH:O[5_555]	1.97	0.23
4:A:888:HOH:O	4:A:888:HOH:O[7_555]	2.04	0.16
4:A:684:HOH:O	4:A:935:HOH:O[7_555]	2.10	0.10

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/342 (99%)	330 (97%)	8 (2%)	2 (1%)	21	15

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	229	ASN
1	A	54	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/282 (100%)	272 (96%)	10 (4%)	32	32

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	GLU
1	A	39	ASN
1	A	49	ASP
1	A	70	ARG
1	A	71	LEU
1	A	117	LYS
1	A	201	LYS
1	A	220	ARG
1	A	225	SER
1	A	235	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	A	69	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	200	HIS
1	A	236	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SHF	A	363[B]	1	6,6,7	1.10	0	6,6,8	2.09	3 (50%)
3	SHF	A	363[A]	1	6,6,7	0.82	0	6,6,8	1.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SHF	A	363[B]	1	-	3/4/4/5	-
3	SHF	A	363[A]	1	-	1/4/4/5	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	363[B]	SHF	OH1-C1-C2	3.12	123.87	114.00
3	A	363[B]	SHF	C3-C2-C1	2.93	122.15	114.51
3	A	363[B]	SHF	OH1-C1-O1	-2.14	117.83	123.33

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	363[A]	SHF	C2-C3-C4-C5
3	A	363[B]	SHF	C2-C3-C4-C5
3	A	363[B]	SHF	OH1-C1-C2-C3
3	A	363[B]	SHF	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	341/342 (99%)	-0.25	11 (3%) 50 54	2, 11, 50, 127	1 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	234	CYS	5.0
1	A	230	GLY	4.8
1	A	229	ASN	4.3
1	A	9	THR	3.9
1	A	231	ASP	3.1
1	A	228	SER	3.1
1	A	225	SER	2.9
1	A	226	ALA	2.8
1	A	341	GLU	2.7
1	A	235	TYR	2.3
1	A	232	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SHF	A	363[A]	7/8	0.91	0.09	1,2,6,19	7
3	SHF	A	363[B]	7/8	0.91	0.09	4,11,18,28	7
2	ZN	A	400	1/1	0.99	0.07	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.