



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:44 PM UTC

PDB ID : 2YLW / pdb_00002ylw
Title : SNAPSHOTS OF ENZYMATIC BAEYER-VILLIGER CATALYSIS: OXY-
GEN ACTIVATION AND INTERMEDIATE STABILIZATION: Arg337Lys
MUTANT IN COMPLEX WITH MES
Authors : Orru, R.; Dudek, H.M.; Martinoli, C.; Torres Pazmino, D.E.; Royant, A.;
Weik, M.; Fraaije, M.W.; Mattevi, A.
Deposited on : 2011-06-06
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	542	 71% 23% ...

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4392 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PHENYLACETONE MONOOXYGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	532	4216	2690	723	794	9	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	337	LYS	ARG	engineered mutation	UNP Q47PU3

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 48	C 21	N 7	O 17	P 3	0	0

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	63	Total 63	O 63	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PHENYLACETONE MONOOXYGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	107.73Å 107.73Å 106.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.70 – 2.90 42.70 – 2.90	Depositor EDS
% Data completeness (in resolution range)	97.7 (42.70-2.90) 97.5 (42.70-2.90)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.68 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.180 , 0.266 0.184 , 0.266	Depositor DCC
R_{free} test set	757 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.201	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 23.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4392	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, MES, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.27	8/4328 (0.2%)	1.30	30/5892 (0.5%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	217	ARG	CA-C	9.52	1.58	1.52
1	A	507	VAL	CA-CB	7.42	1.61	1.53
1	A	361	ALA	CA-C	6.49	1.59	1.52
1	A	252	THR	CA-C	6.35	1.58	1.52
1	A	540	VAL	CA-CB	5.75	1.61	1.53
1	A	223	VAL	CA-CB	5.52	1.59	1.55
1	A	407	VAL	CA-CB	5.31	1.60	1.54
1	A	223	VAL	CA-C	5.10	1.56	1.52

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	242	TYR	N-CA-C	8.23	119.87	111.07
1	A	490	ALA	N-CA-C	-8.03	102.53	111.28
1	A	359	LEU	N-CA-C	-6.86	103.03	111.40
1	A	113	ILE	N-CA-C	6.53	117.43	107.77
1	A	261	GLY	CA-C-N	6.40	125.97	119.19
1	A	261	GLY	C-N-CA	6.40	125.97	119.19
1	A	416	GLY	CA-C-N	6.35	127.77	119.84
1	A	416	GLY	C-N-CA	6.35	127.77	119.84
1	A	56	TYR	N-CA-C	-6.31	104.29	112.68
1	A	513	VAL	CB-CA-C	-6.15	100.06	110.71
1	A	448	VAL	N-CA-C	-6.08	104.92	110.82
1	A	301	ALA	N-CA-C	-6.06	104.59	111.07
1	A	375	GLU	N-CA-C	5.88	120.50	112.04

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	459	ASP	N-CA-C	-5.83	105.01	111.36
1	A	518	VAL	CB-CA-C	-5.82	103.90	111.88
1	A	367	THR	CA-C-N	-5.80	113.78	119.64
1	A	367	THR	C-N-CA	-5.80	113.78	119.64
1	A	92	SER	N-CA-C	-5.78	103.30	110.41
1	A	258	ARG	N-CA-C	5.75	118.67	110.10
1	A	518	VAL	N-CA-C	5.48	117.44	112.29
1	A	179	HIS	N-CA-C	5.40	119.03	112.23
1	A	146	LEU	CA-C-N	-5.38	115.88	122.93
1	A	146	LEU	C-N-CA	-5.38	115.88	122.93
1	A	12	GLN	N-CA-C	5.33	115.72	109.60
1	A	380	LEU	N-CA-C	5.29	118.05	109.06
1	A	58	ASN	N-CA-C	5.24	117.72	109.39
1	A	443	LEU	N-CA-C	-5.21	106.77	113.02
1	A	498	THR	N-CA-C	5.11	117.57	109.50
1	A	515	MET	N-CA-C	5.03	119.42	113.28
1	A	89	ARG	N-CA-C	-5.03	105.80	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4216	0	4053	80	0
2	A	53	0	31	4	0
3	A	48	0	25	2	0
4	A	12	0	12	1	0
5	A	63	0	0	2	0
All	All	4392	0	4121	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:ASN:CB	5:A:2020:HOH:O	1.97	1.12
1:A:246:ARG:CG	1:A:246:ARG:HH11	1.75	0.99
1:A:246:ARG:HH11	1:A:246:ARG:HG3	1.42	0.85
1:A:231:ASP:OD1	1:A:232:PRO:HD2	1.77	0.84
1:A:196:SER:HB3	1:A:339:ILE:HG12	1.64	0.79
1:A:67:ILE:HD12	1:A:256:THR:HG21	1.66	0.77
1:A:270:ASP:OD1	1:A:304:ARG:NH2	2.17	0.77
1:A:471:ARG:NH1	1:A:473:GLU:OE1	2.17	0.76
1:A:506:ASN:HD22	1:A:506:ASN:H	1.32	0.75
1:A:246:ARG:CG	1:A:246:ARG:NH1	2.47	0.73
1:A:372:ARG:HG3	1:A:372:ARG:HH11	1.54	0.71
1:A:464:MET:SD	1:A:541:LEU:HD22	2.31	0.70
1:A:152:GLN:HE22	2:A:700:FAD:H5'2	1.57	0.69
1:A:246:ARG:HH11	1:A:246:ARG:HG2	1.60	0.65
1:A:402:ARG:HD2	1:A:406:ASN:HD21	1.63	0.64
1:A:177:TRP:CD1	1:A:178:PRO:HD2	2.34	0.60
1:A:246:ARG:NH1	1:A:246:ARG:HG2	2.17	0.59
1:A:217:ARG:NH1	3:A:701:NAP:O1X	2.35	0.59
1:A:27:SER:HB3	1:A:434:ILE:HG21	1.84	0.59
1:A:335:THR:OG1	1:A:498:THR:HG21	2.03	0.58
1:A:44:VAL:HB	1:A:113:ILE:HG12	1.86	0.57
1:A:506:ASN:HD22	1:A:506:ASN:N	2.02	0.56
1:A:220:HIS:HA	1:A:331:TYR:OH	2.06	0.56
1:A:17:VAL:O	1:A:143:ALA:HA	2.07	0.55
1:A:310:ARG:HG2	1:A:327:VAL:HG13	1.89	0.55
1:A:329:LYS:HD2	5:A:2043:HOH:O	2.06	0.54
1:A:457:VAL:O	1:A:460:HIS:HB3	2.07	0.53
1:A:441:SER:C	1:A:446:MET:HE3	2.36	0.51
1:A:512:ARG:HH11	1:A:512:ARG:HG3	1.75	0.51
1:A:402:ARG:NH2	1:A:408:ALA:HB2	2.25	0.50
1:A:523:ARG:HA	1:A:526:GLN:HE21	1.75	0.50
1:A:421:LEU:HA	1:A:482:TRP:CE2	2.46	0.50
2:A:700:FAD:N1	2:A:700:FAD:O3'	2.39	0.49
1:A:370:GLY:HA3	1:A:378:TYR:O	2.12	0.49
1:A:503:THR:HG22	1:A:505:ALA:H	1.78	0.49
1:A:495:TYR:HB3	1:A:514:PHE:CE1	2.48	0.48
1:A:231:ASP:OD1	1:A:232:PRO:CD	2.57	0.48
1:A:496:PRO:HB2	1:A:497:MET:HE2	1.94	0.48
1:A:402:ARG:HD2	1:A:406:ASN:ND2	2.27	0.48
1:A:206:ALA:O	1:A:350:ARG:NH2	2.42	0.47
1:A:62:GLY:O	1:A:63:ALA:C	2.57	0.47
1:A:124:PHE:CE2	1:A:429:PRO:HG2	2.49	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:THR:O	1:A:220:HIS:HD2	1.97	0.47
1:A:411:GLU:O	1:A:412:LYS:C	2.59	0.46
1:A:78:GLU:OE2	1:A:82:GLN:NE2	2.47	0.45
1:A:423:LEU:HD21	1:A:453:HIS:CD2	2.52	0.45
1:A:76:PHE:C	1:A:76:PHE:CD1	2.95	0.45
1:A:53:GLY:HA2	2:A:700:FAD:O3B	2.16	0.45
1:A:54:VAL:HG13	2:A:700:FAD:O2A	2.17	0.44
1:A:144:ARG:HG2	1:A:145:TYR:CE2	2.53	0.44
1:A:503:THR:O	1:A:503:THR:HG23	2.18	0.44
1:A:372:ARG:HG3	1:A:372:ARG:NH1	2.28	0.44
1:A:402:ARG:CD	1:A:406:ASN:ND2	2.81	0.44
1:A:217:ARG:O	1:A:359:LEU:HG	2.18	0.44
1:A:441:SER:HB3	1:A:516:LEU:HD22	1.99	0.44
1:A:158:LEU:HD12	1:A:173:HIS:CD2	2.53	0.43
1:A:196:SER:OG	3:A:701:NAP:C4N	2.67	0.43
1:A:331:TYR:HB2	1:A:498:THR:CG2	2.48	0.43
1:A:402:ARG:CD	1:A:406:ASN:HD21	2.30	0.43
1:A:506:ASN:H	1:A:506:ASN:ND2	2.09	0.43
1:A:331:TYR:HB2	1:A:498:THR:HG21	2.01	0.43
1:A:517:TYR:CE2	1:A:524:TYR:HB2	2.54	0.43
1:A:152:GLN:N	1:A:152:GLN:CD	2.77	0.43
1:A:142:ARG:HB2	1:A:142:ARG:CZ	2.49	0.42
1:A:327:VAL:HA	1:A:328:PRO:HD3	1.94	0.42
1:A:265:ALA:HB2	1:A:292:TYR:CB	2.49	0.42
1:A:535:GLY:O	1:A:536:TYR:HB2	2.20	0.42
1:A:337:LYS:HE3	4:A:702:MES:H81	2.00	0.42
1:A:23:GLY:O	1:A:28:GLY:HA3	2.20	0.41
1:A:199:ILE:HG23	1:A:344:TYR:CG	2.55	0.41
1:A:35:LEU:HB2	1:A:42:VAL:HG11	2.03	0.41
1:A:303:GLU:OE2	1:A:310:ARG:NH1	2.47	0.41
1:A:221:PHE:CZ	1:A:328:PRO:HB2	2.56	0.41
1:A:404:VAL:HG23	1:A:427:GLY:O	2.21	0.41
1:A:261:GLY:HA2	1:A:262:PRO:HD2	1.92	0.41
1:A:125:ASP:OD2	1:A:127:ALA:HB3	2.20	0.41
1:A:307:GLU:OE1	1:A:310:ARG:HD2	2.21	0.40
1:A:244:GLU:O	1:A:248:GLU:HG3	2.22	0.40
1:A:331:TYR:CZ	1:A:336:LYS:HD2	2.56	0.40
1:A:416:GLY:HA3	1:A:513:VAL:HG21	2.02	0.40
1:A:505:ALA:HB2	1:A:512:ARG:HD2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	530/542 (98%)	507 (96%)	23 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	438/453 (97%)	406 (93%)	32 (7%)	13	38

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	GLU
1	A	42	VAL
1	A	54	VAL
1	A	88	GLU
1	A	158	LEU
1	A	165	LYS
1	A	166	ASP
1	A	217	ARG
1	A	235	LEU
1	A	240	LYS
1	A	242	TYR
1	A	246	ARG
1	A	279	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	282	GLN
1	A	315	ASN
1	A	337	LYS
1	A	360	SER
1	A	372	ARG
1	A	377	GLU
1	A	392	LEU
1	A	398	LYS
1	A	404	VAL
1	A	444	SER
1	A	466	LYS
1	A	484	GLU
1	A	498	THR
1	A	503	THR
1	A	506	ASN
1	A	512	ARG
1	A	513	VAL
1	A	530	GLU
1	A	540	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	ASN
1	A	152	GLN
1	A	216	GLN
1	A	251	ASN
1	A	315	ASN
1	A	406	ASN
1	A	506	ASN
1	A	526	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAP	A	701	-	50,52,52	1.49	9 (18%)	71,80,80	1.98	21 (29%)
2	FAD	A	700	-	58,58,58	1.21	6 (10%)	85,89,89	1.86	21 (24%)
4	MES	A	702	-	12,12,12	2.06	1 (8%)	15,16,16	2.75	8 (53%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	701	-	-	6/35/67/67	0/5/5/5
2	FAD	A	700	-	-	4/34/50/50	0/6/6/6
4	MES	A	702	-	-	4/6/14/14	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	702	MES	C8-S	-6.81	1.68	1.77
3	A	701	NAP	O4D-C1D	4.64	1.47	1.40
2	A	700	FAD	C8A-N7A	3.90	1.39	1.31
3	A	701	NAP	P2B-O1X	3.42	1.61	1.50
2	A	700	FAD	C4X-N5	3.30	1.37	1.30
3	A	701	NAP	C7N-N7N	3.04	1.38	1.33
3	A	701	NAP	C5A-C4A	2.70	1.43	1.39
2	A	700	FAD	C10-N1	2.68	1.38	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	701	NAP	C6A-N6A	2.68	1.41	1.34
3	A	701	NAP	C2N-N1N	2.53	1.37	1.35
3	A	701	NAP	P2B-O2B	2.43	1.63	1.59
3	A	701	NAP	C3D-C4D	2.37	1.59	1.53
3	A	701	NAP	P2B-O3X	2.35	1.63	1.54
2	A	700	FAD	C2A-N3A	2.34	1.38	1.33
2	A	700	FAD	C9A-N10	-2.27	1.37	1.41
2	A	700	FAD	P-O3P	2.12	1.61	1.59

All (50) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	702	MES	C5-N4-C3	6.90	123.70	108.84
3	A	701	NAP	P2B-O2B-C2B	-6.16	106.99	123.43
2	A	700	FAD	N3A-C2A-N1A	-5.39	120.42	128.58
2	A	700	FAD	C5A-C4A-N3A	-5.32	119.39	126.72
3	A	701	NAP	C3N-C7N-N7N	4.94	123.83	117.74
4	A	702	MES	O3S-S-C8	4.48	114.77	106.00
2	A	700	FAD	O4B-C1B-N9A	4.41	116.55	108.09
2	A	700	FAD	N9A-C8A-N7A	-4.31	107.83	113.94
3	A	701	NAP	N3A-C2A-N1A	-3.97	122.57	128.58
2	A	700	FAD	C2A-N3A-C4A	3.91	121.39	111.83
3	A	701	NAP	C4D-O4D-C1D	-3.75	106.49	109.92
2	A	700	FAD	C5A-N7A-C8A	3.68	109.23	103.45
3	A	701	NAP	N3A-C4A-N9A	3.61	133.30	127.17
2	A	700	FAD	C4A-C5A-N7A	-3.58	106.49	110.58
2	A	700	FAD	C4-N3-C2	-3.53	119.37	125.64
3	A	701	NAP	C2B-C1B-N9A	-3.53	107.94	113.75
3	A	701	NAP	C3B-C2B-C1B	-3.42	96.26	102.81
3	A	701	NAP	C5A-C4A-N3A	-3.39	122.04	126.72
2	A	700	FAD	C5A-C4A-N9A	3.36	109.48	105.81
2	A	700	FAD	C4X-C10-N10	3.29	121.19	116.48
2	A	700	FAD	C10-C4X-N5	-3.19	118.28	124.81
2	A	700	FAD	C4X-C10-N1	-3.19	116.76	124.59
3	A	701	NAP	C4A-N9A-C8A	3.03	108.92	105.74
3	A	701	NAP	O2B-C2B-C1B	-3.02	99.42	110.05
4	A	702	MES	O3S-S-O2S	-3.02	103.85	111.40
3	A	701	NAP	C5A-N7A-C8A	3.00	108.16	103.45
3	A	701	NAP	O5B-C5B-C4B	-2.91	99.10	108.99
3	A	701	NAP	C2A-N3A-C4A	2.90	118.91	111.83
3	A	701	NAP	O7N-C7N-C3N	-2.89	116.06	119.60
2	A	700	FAD	O2A-PA-O3P	2.75	114.70	107.27

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	701	NAP	N9A-C8A-N7A	-2.70	110.10	113.94
4	A	702	MES	C7-N4-C3	2.67	118.34	111.24
2	A	700	FAD	C6-C7-C8	-2.59	115.89	119.69
2	A	700	FAD	O5'-C5'-C4'	-2.54	102.59	109.36
4	A	702	MES	C7-N4-C5	2.50	117.89	111.24
3	A	701	NAP	O3B-C3B-C4B	-2.47	103.99	111.08
4	A	702	MES	O1-C6-C5	-2.41	106.58	111.77
2	A	700	FAD	O4-C4-C4X	-2.40	120.20	126.53
3	A	701	NAP	C4A-N9A-C1B	-2.38	121.06	126.63
3	A	701	NAP	O2X-P2B-O2B	2.36	115.04	105.85
2	A	700	FAD	N3A-C4A-N9A	2.31	131.09	127.17
3	A	701	NAP	PN-O5D-C5D	-2.25	108.44	121.35
2	A	700	FAD	C4-C4X-C10	2.17	120.65	116.93
2	A	700	FAD	C4X-C4-N3	2.14	118.70	113.25
3	A	701	NAP	C4A-C5A-N7A	-2.12	108.15	110.58
3	A	701	NAP	C2A-N1A-C6A	2.07	122.13	118.73
2	A	700	FAD	C4-C4X-N5	2.07	121.06	118.21
4	A	702	MES	O2S-S-O1S	-2.02	107.26	113.82
2	A	700	FAD	O3B-C3B-C4B	-2.02	105.28	111.08
4	A	702	MES	O1S-S-C8	2.01	109.76	106.73

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	700	FAD	N10-C1'-C2'-O2'
2	A	700	FAD	N10-C1'-C2'-C3'
3	A	701	NAP	O4D-C1D-N1N-C2N
3	A	701	NAP	O4D-C4D-C5D-O5D
3	A	701	NAP	C3D-C4D-C5D-O5D
4	A	702	MES	C7-C8-S-O3S
3	A	701	NAP	PA-O3-PN-O5D
4	A	702	MES	C7-C8-S-O1S
4	A	702	MES	C7-C8-S-O2S
2	A	700	FAD	C3B-C4B-C5B-O5B
2	A	700	FAD	O4B-C4B-C5B-O5B
4	A	702	MES	C8-C7-N4-C5
3	A	701	NAP	PA-O3-PN-O1N
3	A	701	NAP	O4B-C4B-C5B-O5B


There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	701	NAP	2	0
2	A	700	FAD	4	0
4	A	702	MES	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	532/542 (98%)	-0.54	2 (0%) 88 85	2, 11, 23, 34	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	11	ARG	2.5
1	A	315	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MES	A	702	12/12	0.95	0.11	38,43,49,50	0
2	FAD	A	700	53/53	0.98	0.05	2,4,12,14	0
3	NAP	A	701	48/48	0.99	0.05	2,7,9,10	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FAD A 700:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NAP A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.