



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 01:02 PM EDT

PDB ID : 5YL7 / pdb_00005yl7
Title : Proteases from Pseudoalteromonas arctica PAMC 21717 (Pro21717)
Authors : Lee, J.H.; Lee, C.W.
Deposited on : 2017-10-17
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	 86% 11% ..
2	B	3	 67% 33%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2985 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called *Pseudoalteromonas arctica* PAMC 21717.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	339	Total	C	N	O	S	0	0	0
			2371	1431	423	508	9			

- Molecule 2 is a protein called Copurified unknown peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	0	0
			15	9	3	3			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Ca	0	0
			4	4		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	593	Total	O	0	0
			593	593		
4	B	2	Total	O	0	0
			2	2		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: *Pseudoalteromonas arctica* PAMC 21717

Chain A:  86% 11% ..



- Molecule 2: Copurified unknown peptide

Chain B:  67% 33%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.95Å 74.56Å 83.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.40 30.00 – 1.40	Depositor EDS
% Data completeness (in resolution range)	97.9 (30.00-1.40) 98.0 (30.00-1.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	29.67 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.130 , 0.155 0.130 , 0.156	Depositor DCC
R_{free} test set	2939 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	8.0	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 39.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2985	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.73	24/2414 (1.0%)	1.45	9/3305 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	314	MET	SD-CE	-13.56	1.45	1.79
1	A	303	PRO	C-O	12.96	1.39	1.23
1	A	306	SER	CB-OG	-9.65	1.22	1.42
1	A	464	THR	CA-C	6.34	1.60	1.52
1	A	257	ASN	CA-C	-6.32	1.44	1.53
1	A	423	GLY	CA-C	-6.31	1.45	1.51
1	A	173	THR	CA-CB	6.12	1.66	1.54
1	A	418	TYR	CA-C	-6.01	1.45	1.52
1	A	160	GLU	N-CA	5.96	1.53	1.45
1	A	344	ALA	CA-C	-5.87	1.44	1.52
1	A	347	ASP	CG-OD2	5.69	1.36	1.25
1	A	335	SER	N-CA	5.67	1.53	1.46
1	A	405	THR	CA-C	-5.60	1.45	1.52
1	A	344	ALA	C-O	5.59	1.31	1.24
1	A	406	TYR	CD2-CE2	5.52	1.55	1.38
1	A	204	MET	CA-C	-5.46	1.45	1.52
1	A	438	ILE	CA-C	-5.41	1.46	1.52
1	A	438	ILE	C-O	5.41	1.30	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	202	TYR	CE1-CZ	-5.38	1.25	1.38
1	A	302	ILE	CA-C	5.34	1.56	1.52
1	A	241	SER	C-O	-5.27	1.17	1.23
1	A	396	ALA	C-O	-5.18	1.17	1.23
1	A	196	ALA	C-N	5.09	1.40	1.33
1	A	307	ASN	N-CA	-5.04	1.41	1.46

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	314	MET	CG-SD-CE	-9.10	80.89	100.90
1	A	459	ARG	NE-CZ-NH2	7.58	126.03	119.20
1	A	300	SER	O-C-N	7.29	131.99	122.95
1	A	300	SER	CA-C-O	-6.87	113.09	120.92
1	A	473	VAL	O-C-N	6.13	128.73	122.79
1	A	307	ASN	OD1-CG-ND2	6.11	128.71	122.60
1	A	148	ASN	CA-C-O	5.27	125.80	119.43
1	A	359	CYS	O-C-N	-5.17	116.90	122.79
1	A	350	ASN	CA-C-O	5.11	126.51	120.58

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	459	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2371	0	2203	15	0
2	B	15	0	5	1	0
3	A	4	0	0	0	0
4	A	593	0	0	11	9
4	B	2	0	0	1	0
All	All	2985	0	2208	16	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:ARG:NH1	4:A:902:HOH:O	2.12	0.82
1:A:284:THR:HG23	4:A:1168:HOH:O	1.80	0.81
1:A:278:GLY:O	4:A:901:HOH:O	2.01	0.78
2:B:3:UNK:CB	4:B:102:HOH:O	2.32	0.77
1:A:322:CYS:SG	4:A:1044:HOH:O	2.51	0.68
1:A:414:SER:O	4:A:903:HOH:O	2.17	0.59
1:A:340:VAL:HG23	4:A:979:HOH:O	2.12	0.49
1:A:364:ASN:ND2	4:A:918:HOH:O	2.49	0.46
1:A:314:MET:HB2	1:A:314:MET:HE3	1.96	0.42
1:A:316:LEU:C	1:A:316:LEU:HD12	2.45	0.42
1:A:411:SER:HB2	4:A:1357:HOH:O	2.18	0.42
1:A:157:HIS:CD2	1:A:157:HIS:H	2.38	0.42
1:A:383:ASN:ND2	4:A:918:HOH:O	2.51	0.42
1:A:204:MET:HE1	1:A:302:ILE:CD1	2.50	0.41
1:A:334:ARG:HG2	4:A:979:HOH:O	2.19	0.41
1:A:407:ASN:ND2	4:A:903:HOH:O	2.19	0.41

All (9) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1355:HOH:O	4:A:1459:HOH:O[3_554]	1.87	0.33
4:A:1093:HOH:O	4:A:1148:HOH:O[2_554]	1.95	0.25
4:A:1175:HOH:O	4:A:1232:HOH:O[3_554]	1.97	0.23
4:A:907:HOH:O	4:A:1288:HOH:O[2_555]	2.03	0.17
4:A:972:HOH:O	4:A:1282:HOH:O[2_554]	2.14	0.06
4:A:1064:HOH:O	4:A:1202:HOH:O[2_554]	2.14	0.06
4:A:936:HOH:O	4:A:1204:HOH:O[2_555]	2.16	0.04
4:A:1219:HOH:O	4:A:1419:HOH:O[2_555]	2.18	0.02
4:A:1204:HOH:O	4:A:1424:HOH:O[2_554]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	337/347 (97%)	325 (96%)	12 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/254 (97%)	247 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	313	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	339/347 (97%)	-0.38	1 (0%) 90 92	4, 7, 17, 24	0
2	B	0/3	-	-	-	-
All	All	339/350 (96%)	-0.38	1 (0%) 90 92	4, 7, 17, 24	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	234	GLY	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	A	801	1/1	0.99	0.02	6,6,6,6	0
3	CA	A	802	1/1	1.00	0.02	7,7,7,7	0
3	CA	A	803	1/1	1.00	0.01	5,5,5,5	0
3	CA	A	804	1/1	1.00	0.01	6,6,6,6	0

6.5 Other polymers [i](#)

There are no such residues in this entry.