



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 01:34 AM UTC

PDB ID : 2YNZ / pdb_00002ynz
Title : Salmonella enterica SadA 823-947 fused to a GCN4 adaptor (SadAK5)
Authors : Hartmann, M.D.; Hernandez Alvarez, B.; Lupas, A.N.
Deposited on : 2012-10-20
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	154	4% 67% •• 29%
1	B	154	3% 65% 5% • 29%
1	C	154	4% 64% 5% • 29%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2887 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GENERAL CONTROL PROTEIN GCN4, PUTATIVE INNER MEMBRANE PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	109	Total	C	N	O	S	0	3	0
			831	509	136	185	1			
1	B	109	Total	C	N	O	S	0	5	0
			851	528	140	182	1			
1	C	109	Total	C	N	O	S	0	3	0
			838	515	137	185	1			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	797	ILE	LEU	engineered mutation	UNP P03069
A	801	ILE	VAL	engineered mutation	UNP P03069
A	804	ILE	LEU	engineered mutation	UNP P03069
A	808	ILE	ASN	engineered mutation	UNP P03069
A	811	ILE	LEU	engineered mutation	UNP P03069
A	815	ILE	VAL	engineered mutation	UNP P03069
A	818	ILE	LEU	engineered mutation	UNP P03069
A	822	ILE	VAL	engineered mutation	UNP P03069
B	797	ILE	LEU	engineered mutation	UNP P03069
B	801	ILE	VAL	engineered mutation	UNP P03069
B	804	ILE	LEU	engineered mutation	UNP P03069
B	808	ILE	ASN	engineered mutation	UNP P03069
B	811	ILE	LEU	engineered mutation	UNP P03069
B	815	ILE	VAL	engineered mutation	UNP P03069
B	818	ILE	LEU	engineered mutation	UNP P03069
B	822	ILE	VAL	engineered mutation	UNP P03069
C	797	ILE	LEU	engineered mutation	UNP P03069
C	801	ILE	VAL	engineered mutation	UNP P03069
C	804	ILE	LEU	engineered mutation	UNP P03069
C	808	ILE	ASN	engineered mutation	UNP P03069
C	811	ILE	LEU	engineered mutation	UNP P03069
C	815	ILE	VAL	engineered mutation	UNP P03069

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	818	ILE	LEU	engineered mutation	UNP P03069
C	822	ILE	VAL	engineered mutation	UNP P03069

- Molecule 2 is NITRATE ION (CCD ID: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	N	O	0	0
			4	1	3		
2	A	1	Total	N	O	0	0
			4	1	3		
2	B	1	Total	N	O	0	0
			4	1	3		
2	C	1	Total	N	O	0	0
			4	1	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	146	Total	O	0	0
			146	146		
3	B	107	Total	O	0	0
			107	107		
3	C	98	Total	O	0	0
			98	98		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PUTATIVE INNER MEMBRANE PROTEIN



- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PUTATIVE INNER MEMBRANE PROTEIN



- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PUTATIVE INNER MEMBRANE PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.69Å 60.30Å 135.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.27 – 1.40 19.27 – 1.40	Depositor EDS
% Data completeness (in resolution range)	98.8 (19.27-1.40) 98.8 (19.27-1.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.176 , 0.232 0.182 , 0.233	Depositor DCC
R_{free} test set	3640 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	12.9	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2887	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/847	0.81	0/1148
1	B	0.64	0/874	0.78	0/1184
1	C	0.64	0/854	0.80	0/1158
All	All	0.64	0/2575	0.80	0/3490

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	831	0	798	9	0
1	B	851	0	837	14	0
1	C	838	0	812	18	0
2	A	8	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
3	A	146	0	0	0	2
3	B	107	0	0	1	1
3	C	98	0	0	0	1
All	All	2887	0	2447	23	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:797:ILE:HD13	1:C:797:ILE:CD1	1.72	1.18
1:B:797:ILE:HD13	1:C:797:ILE:HD11	1.32	1.12
1:A:843:ILE:HD11	1:C:839:ILE:HG23	1.55	0.88
1:A:797:ILE:HD13	1:C:797:ILE:HD13	1.60	0.82
1:B:797:ILE:HD13	1:C:797:ILE:HD12	1.66	0.78
1:A:843:ILE:CD1	1:C:839:ILE:HG23	2.24	0.68
1:A:797:ILE:HD13	1:C:797:ILE:CD1	2.24	0.66
1:B:797:ILE:CD1	1:C:797:ILE:HD11	2.20	0.63
1:A:898:MET:O	1:A:902:GLN:HG3	2.02	0.60
1:C:796:GLN:NE2	1:C:800:LYS:HE2	2.18	0.59
1:B:821[B]:LEU:HD23	1:C:822:ILE:HG21	1.88	0.55
1:A:797:ILE:CD1	1:C:797:ILE:HD13	2.33	0.53
1:A:839:ILE:HG23	1:B:843:ILE:CD1	2.39	0.53
1:B:797:ILE:CD1	1:C:797:ILE:CD1	2.66	0.52
1:B:811:ILE:HG23	1:C:815[A]:ILE:CD1	2.43	0.49
1:A:888:VAL:HG22	1:B:888:VAL:HG23	1.95	0.48
1:C:796:GLN:HE22	1:C:800:LYS:HE2	1.80	0.46
1:A:888:VAL:CG2	1:B:888:VAL:HG23	2.47	0.44
1:B:800:LYS:HG2	1:C:801:ILE:HG21	2.00	0.44
1:B:817[B]:ARG:NH1	3:B:2015:HOH:O	2.50	0.43
1:B:888:VAL:CG2	1:C:888:VAL:HG23	2.50	0.42
1:B:811:ILE:HG23	1:C:815[A]:ILE:HD11	2.02	0.41
1:C:894:TYR:CZ	1:C:898:MET:HE3	2.55	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:2107:HOH:O	3:C:2072:HOH:O[3_655]	1.99	0.21
3:A:2012:HOH:O	3:B:2061:HOH:O[2_554]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	110/154 (71%)	110 (100%)	0	0	100	100
1	B	112/154 (73%)	112 (100%)	0	0	100	100
1	C	110/154 (71%)	110 (100%)	0	0	100	100
All	All	332/462 (72%)	332 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	93/128 (73%)	92 (99%)	1 (1%)	65	37
1	B	95/128 (74%)	91 (96%)	4 (4%)	26	4
1	C	94/128 (73%)	90 (96%)	4 (4%)	26	4
All	All	282/384 (73%)	273 (97%)	9 (3%)	33	7

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	797	ILE
1	B	796	GLN
1	B	797	ILE
1	B	800	LYS
1	B	898	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	796	GLN
1	C	797	ILE
1	C	800	LYS
1	C	802	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	831	GLN
1	A	897	ASN
1	B	897	ASN
1	C	796	GLN
1	C	826	ASN
1	C	874	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NO3	B	1905	-	1,3,3	2.94	1 (100%)	0,3,3	-	-
2	NO3	A	1906	-	1,3,3	3.09	1 (100%)	0,3,3	-	-
2	NO3	A	1905	-	1,3,3	3.00	1 (100%)	0,3,3	-	-
2	NO3	C	1905	-	1,3,3	3.12	1 (100%)	0,3,3	-	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1905	NO3	O1-N	3.12	1.39	1.24
2	A	1906	NO3	O1-N	3.09	1.39	1.24
2	A	1905	NO3	O1-N	3.00	1.39	1.24
2	B	1905	NO3	O1-N	2.94	1.38	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	109/154 (70%)	-0.07	6 (5%)	30 32	9, 17, 45, 53	3 (2%)
1	B	109/154 (70%)	-0.02	4 (3%)	45 47	8, 18, 43, 51	5 (4%)
1	C	109/154 (70%)	-0.03	6 (5%)	30 32	8, 18, 47, 56	3 (2%)
All	All	327/462 (70%)	-0.04	16 (4%)	35 37	8, 18, 45, 56	11 (3%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	903	TYR	7.3
1	C	904	ASN	4.2
1	A	797	ILE	4.0
1	B	904	ASN	3.6
1	B	903	TYR	3.6
1	A	904	ASN	2.9
1	B	899	LEU	2.9
1	C	899	LEU	2.9
1	A	903	TYR	2.7
1	C	901	SER	2.7
1	A	834	SER	2.6
1	A	901	SER	2.5
1	C	902	GLN	2.3
1	B	900	ILE	2.1
1	C	797	ILE	2.1
1	A	902	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NO3	A	1906	4/4	0.92	0.09	25,32,37,39	0
2	NO3	C	1905	4/4	0.94	0.09	17,18,19,24	0
2	NO3	A	1905	4/4	0.96	0.07	22,27,31,33	0
2	NO3	B	1905	4/4	0.98	0.05	16,18,18,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.