



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:35 AM UTC

PDB ID : 2YO0 / pdb_00002yo0
Title : Salmonella enterica SadA 1049-1304 fused to GCN4 adaptors (SadAK9-cfl)
Authors : Hartmann, M.D.; Hernandez Alvarez, B.; Lupas, A.N.
Deposited on : 2012-10-20
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	322	14% 62% 20% • 14%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1989 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GENERAL CONTROL PROTEIN GCN4, PUTATIVE INNER MEMBRANE PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	S	0	0	0
			1986	1194	357	430	5			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1023	ILE	LEU	engineered mutation	UNP P03069
A	1027	ILE	VAL	engineered mutation	UNP P03069
A	1030	ILE	LEU	engineered mutation	UNP P03069
A	1034	ILE	ASN	engineered mutation	UNP P03069
A	1037	ILE	LEU	engineered mutation	UNP P03069
A	1041	ILE	VAL	engineered mutation	UNP P03069
A	1044	ILE	LEU	engineered mutation	UNP P03069
A	1048	ILE	VAL	engineered mutation	UNP P03069
A	1308	ILE	LEU	engineered mutation	UNP P03069
A	1312	ILE	VAL	engineered mutation	UNP P03069
A	1315	ILE	LEU	engineered mutation	UNP P03069
A	1319	ILE	ASN	engineered mutation	UNP P03069
A	1322	ILE	LEU	engineered mutation	UNP P03069
A	1326	ILE	VAL	engineered mutation	UNP P03069
A	1329	ILE	LEU	engineered mutation	UNP P03069
A	1333	ILE	VAL	engineered mutation	UNP P03069
A	1334	LYS	-	expression tag	UNP P03069
A	1335	LEU	-	expression tag	UNP P03069
A	1336	HIS	-	expression tag	UNP P03069
A	1337	HIS	-	expression tag	UNP P03069
A	1338	HIS	-	expression tag	UNP P03069
A	1339	HIS	-	expression tag	UNP P03069
A	1340	HIS	-	expression tag	UNP P03069
A	1341	HIS	-	expression tag	UNP P03069

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	O	0	0
			2	2		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PUTATIVE INNER MEMBRANE PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	54.00Å 54.00Å 306.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.53 – 2.80 34.53 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.4 (34.53-2.80) 99.8 (34.53-2.80)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.09 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.221 , 0.297 0.231 , 0.293	Depositor DCC
R_{free} test set	654 reflections (5.27%)	wwPDB-VP
Wilson B-factor (Å ²)	34.9	Xtriage
Anisotropy	0.362	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 18.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.109 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.571 for H,K,L 0.429 for -H-K,K,-L	Depositor
Outliers	6 of 12379 reflections (0.048%)	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	1989	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/1998	0.89	2/2715 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	1103	MET	N-CA-C	5.92	117.97	110.33
1	A	1124	ILE	N-CA-C	5.00	115.85	108.45

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1986	0	1925	52	0
2	A	1	0	0	0	0
3	A	2	0	0	0	0
All	All	1989	0	1925	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1110:ILE:HD13	1:A:1124:ILE:HD11	1.31	1.11
1:A:1110:ILE:HD13	1:A:1124:ILE:CD1	1.88	1.04
1:A:1074:ALA:HB1	1:A:1079:SER:OG	1.60	1.00
1:A:1110:ILE:HB	1:A:1124:ILE:HD12	1.57	0.86
1:A:1073:LEU:HD21	1:A:1075:VAL:CG2	2.09	0.83
1:A:1073:LEU:HD21	1:A:1075:VAL:HG21	1.72	0.70
1:A:1320:TYR:O	1:A:1324:ASN:ND2	2.24	0.70
1:A:1123:SER:O	1:A:1124:ILE:HD13	1.93	0.69
1:A:1073:LEU:HD11	1:A:1075:VAL:HG23	1.76	0.68
1:A:1219:ILE:HG23	1:A:1222:ILE:HD12	1.77	0.67
1:A:1073:LEU:HD21	1:A:1075:VAL:HG23	1.77	0.66
1:A:1077:THR:N	1:A:1091:ASP:OD1	2.28	0.66
1:A:1082:MET:HE3	1:A:1096:ILE:HG12	1.80	0.64
1:A:1197:SER:O	1:A:1201:LEU:HD13	2.00	0.62
1:A:1110:ILE:HG22	1:A:1112:ILE:HG13	1.83	0.61
1:A:1110:ILE:CD1	1:A:1124:ILE:CD1	2.71	0.61
1:A:1128:ASN:HD22	1:A:1164:ARG:NH2	2.01	0.59
1:A:1075:VAL:HG12	1:A:1076:GLY:N	2.18	0.58
1:A:1101:LEU:HD21	1:A:1103:MET:HG3	1.87	0.56
1:A:1252:SER:HB2	1:A:1266:THR:HG23	1.87	0.56
1:A:1190:GLN:O	1:A:1191:VAL:C	2.48	0.55
1:A:1073:LEU:CG	1:A:1075:VAL:HG23	2.38	0.54
1:A:1128:ASN:ND2	1:A:1164:ARG:NH2	2.56	0.54
1:A:1270:ALA:HB2	1:A:1277:SER:HB3	1.90	0.53
1:A:1082:MET:CE	1:A:1096:ILE:HG12	2.38	0.52
1:A:1257:ALA:HB3	1:A:1271:ASP:HA	1.91	0.52
1:A:1145:MET:HA	1:A:1145:MET:HE2	1.92	0.52
1:A:1078:ASP:O	1:A:1092:ALA:HA	2.09	0.52
1:A:1073:LEU:CD1	1:A:1075:VAL:HG23	2.39	0.52
1:A:1073:LEU:CD2	1:A:1075:VAL:HG23	2.40	0.51
1:A:1114:SER:O	1:A:1129:GLY:N	2.43	0.51
1:A:1138:THR:O	1:A:1139:ASP:C	2.53	0.50
1:A:1319:ILE:HA	1:A:1322:ILE:HD12	1.94	0.49
1:A:1110:ILE:CG2	1:A:1112:ILE:HD11	2.42	0.49
1:A:1172:GLY:H	1:A:1183:GLN:NE2	2.10	0.49
1:A:1204:GLN:O	1:A:1208:LEU:HD13	2.14	0.48
1:A:1099:ASN:HB3	1:A:1115:ASN:HD22	1.78	0.48
1:A:1292:ALA:HB1	1:A:1304:GLN:CD	2.39	0.48
1:A:1219:ILE:HG23	1:A:1222:ILE:CD1	2.44	0.47
1:A:1287:ILE:HG22	1:A:1290:VAL:CG2	2.44	0.47
1:A:1140:TYR:CE1	1:A:1142:ALA:HB2	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1086:THR:HG22	1:A:1086:THR:O	2.15	0.46
1:A:1252:SER:CB	1:A:1266:THR:HG23	2.45	0.46
1:A:1172:GLY:H	1:A:1183:GLN:HE22	1.64	0.46
1:A:1128:ASN:HD22	1:A:1128:ASN:N	2.14	0.45
1:A:1128:ASN:ND2	1:A:1128:ASN:O	2.50	0.45
1:A:1217:ASN:HD22	1:A:1217:ASN:N	2.15	0.43
1:A:1147:THR:O	1:A:1148:PRO:C	2.62	0.43
1:A:1250:ILE:HD12	1:A:1264:LEU:HD13	2.00	0.42
1:A:1082:MET:HE3	1:A:1096:ILE:CG1	2.48	0.41
1:A:1248:VAL:HG12	1:A:1249:ALA:N	2.37	0.40
1:A:1073:LEU:CD2	1:A:1075:VAL:CG2	2.92	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/322 (85%)	235 (86%)	34 (12%)	5 (2%)	6	23

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1104	ALA
1	A	1171	ALA
1	A	1219	ILE
1	A	1335	LEU
1	A	1107	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	211/257 (82%)	190 (90%)	21 (10%)	7 24

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1075	VAL
1	A	1077	THR
1	A	1085	LYS
1	A	1086	THR
1	A	1105	ASP
1	A	1107	ILE
1	A	1128	ASN
1	A	1169	VAL
1	A	1199	THR
1	A	1203	THR
1	A	1205	VAL
1	A	1228	THR
1	A	1252	SER
1	A	1259	GLU
1	A	1282	THR
1	A	1288	THR
1	A	1312	ILE
1	A	1314	GLU
1	A	1319	ILE
1	A	1335	LEU
1	A	1336	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1089	ASN
1	A	1099	ASN
1	A	1115	ASN
1	A	1119	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1128	ASN
1	A	1150	ASN
1	A	1168	ASN
1	A	1183	GLN
1	A	1217	ASN
1	A	1267	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/322 (85%)	0.87	46 (16%) 4 3	2, 30, 88, 90	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1232	LYS	5.5
1	A	1248	VAL	5.2
1	A	1262	VAL	5.1
1	A	1256	ALA	5.0
1	A	1269	VAL	4.8
1	A	1231	PHE	4.7
1	A	1285	ARG	4.5
1	A	1260	ASN	4.2
1	A	1227	SER	4.2
1	A	1224	THR	4.1
1	A	1249	ALA	4.1
1	A	1255	ILE	3.9
1	A	1276	VAL	3.8
1	A	1336	HIS	3.7
1	A	1218	GLY	3.7
1	A	1270	ALA	3.5
1	A	1257	ALA	3.4
1	A	1228	THR	3.3
1	A	1233	THR	3.3
1	A	1225	THR	3.2
1	A	1252	SER	3.1
1	A	1230	TYR	3.0
1	A	1290	VAL	3.0
1	A	1287	ILE	2.9
1	A	1264	LEU	2.9
1	A	1296	ASN	2.8
1	A	1277	SER	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1229	LYS	2.7
1	A	1160	GLU	2.7
1	A	1278	VAL	2.7
1	A	1226	GLY	2.6
1	A	1223	VAL	2.6
1	A	1282	THR	2.6
1	A	1238	ALA	2.5
1	A	1302	VAL	2.5
1	A	1251	GLY	2.4
1	A	1219	ILE	2.4
1	A	1307	GLN	2.3
1	A	1286	ARG	2.3
1	A	1205	VAL	2.3
1	A	1236	ASP	2.3
1	A	1235	THR	2.2
1	A	1335	LEU	2.1
1	A	1222	ILE	2.1
1	A	1242	ALA	2.1
1	A	1313	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	A	2337	1/1	0.97	0.10	31,31,31,31	1

6.5 Other polymers [i](#)

There are no such residues in this entry.