



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:08 PM UTC

PDB ID : 2YPT / pdb_00002ypt
Title : Crystal structure of the human nuclear membrane zinc metalloprotease ZMP-STE24 mutant (E336A) in complex with a synthetic CSIM tetrapeptide from the C-terminus of prelamin A
Authors : Pike, A.C.W.; Dong, Y.Y.; Quigley, A.; Dong, L.; Savitsky, P.; Cooper, C.D.O.; Chaikuad, A.; Goubin, S.; Shrestha, L.; Li, Q.; Mukhopadhyay, S.; Yang, J.; Xia, X.; Shintre, C.A.; Barr, A.J.; Berridge, G.; Chalk, R.; Bray, J.E.; von Delft, F.; Bullock, A.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; Burgess-Brown, N.; Carpenter, E.P.
Deposited on : 2012-11-01
Resolution : 3.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1065 (3.96-3.64)
Clashscore	190562	1012 (3.94-3.66)
Ramachandran outliers	187476	1048 (3.96-3.64)
Sidechain outliers	187428	1043 (3.96-3.64)
RSRZ outliers	180081	1064 (3.96-3.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	4% 83% 5% 12%
1	B	482	6% 83% • 14%
1	D	482	5% 80% 6% 14%

Continued on next page...

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	482	3% 82% 14%
2	F	4	25% 50% 25% 25%
2	G	4	25% 100%
2	H	4	25% 75% 25%
2	I	4	25% 50% 25% 25%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12513 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CAAX PRENYL PROTEASE 1 HOMOLOG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	424	Total	C	N	O	S	0	1	0
			3153	2096	489	561	7			
1	B	416	Total	C	N	O	S	0	1	0
			3061	2026	482	546	7			
1	D	415	Total	C	N	O	S	0	1	0
			3156	2101	490	557	8			
1	E	414	Total	C	N	O	S	0	1	0
			3045	2015	473	550	7			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	137	ALA	THR	variant	UNP O75844
A	336	ALA	GLU	engineered mutation	UNP O75844
A	476	ALA	-	expression tag	UNP O75844
A	477	GLU	-	expression tag	UNP O75844
A	478	ASN	-	expression tag	UNP O75844
A	479	LEU	-	expression tag	UNP O75844
A	480	TYR	-	expression tag	UNP O75844
A	481	PHE	-	expression tag	UNP O75844
A	482	GLN	-	expression tag	UNP O75844
B	137	ALA	THR	variant	UNP O75844
B	336	ALA	GLU	engineered mutation	UNP O75844
B	476	ALA	-	expression tag	UNP O75844
B	477	GLU	-	expression tag	UNP O75844
B	478	ASN	-	expression tag	UNP O75844
B	479	LEU	-	expression tag	UNP O75844
B	480	TYR	-	expression tag	UNP O75844
B	481	PHE	-	expression tag	UNP O75844
B	482	GLN	-	expression tag	UNP O75844
D	137	ALA	THR	variant	UNP O75844
D	336	ALA	GLU	engineered mutation	UNP O75844
D	476	ALA	-	expression tag	UNP O75844

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	477	GLU	-	expression tag	UNP O75844
D	478	ASN	-	expression tag	UNP O75844
D	479	LEU	-	expression tag	UNP O75844
D	480	TYR	-	expression tag	UNP O75844
D	481	PHE	-	expression tag	UNP O75844
D	482	GLN	-	expression tag	UNP O75844
E	137	ALA	THR	variant	UNP O75844
E	336	ALA	GLU	engineered mutation	UNP O75844
E	476	ALA	-	expression tag	UNP O75844
E	477	GLU	-	expression tag	UNP O75844
E	478	ASN	-	expression tag	UNP O75844
E	479	LEU	-	expression tag	UNP O75844
E	480	TYR	-	expression tag	UNP O75844
E	481	PHE	-	expression tag	UNP O75844
E	482	GLN	-	expression tag	UNP O75844

- Molecule 2 is a protein called PRELAMIN-A/C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	3	Total	C	N	O	S	0	0	0
			19	11	3	4	1			
2	G	4	Total	C	N	O	S	0	0	0
			28	16	4	6	2			
2	H	4	Total	C	N	O	S	0	0	0
			28	16	4	6	2			
2	I	3	Total	C	N	O	S	0	0	0
			19	11	3	4	1			


- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		
3	E	1	Total	Zn	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CAAX PRENYL PROTEASE 1 HOMOLOG




• Molecule 1: CAAX PRENYL PROTEASE 1 HOMOLOG





• Molecule 1: CAAX PRENYL PROTEASE 1 HOMOLOG





• Molecule 1: CAAX PRENYL PROTEASE 1 HOMOLOG




• Molecule 2: PRELAMIN-A/C



• Molecule 2: PRELAMIN-A/C



• Molecule 2: PRELAMIN-A/C



• Molecule 2: PRELAMIN-A/C



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	61.32Å 95.54Å 132.03Å 76.26° 79.67° 72.30°	Depositor
Resolution (Å)	38.50 – 3.80 38.50 – 3.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (38.50-3.80) 99.2 (38.50-3.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 3.76Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.263 , 0.281 (Not available) , 0.282	Depositor DCC
R_{free} test set	1420 reflections (5.27%)	wwPDB-VP
Wilson B-factor (Å ²)	153.1	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 140.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	12513	wwPDB-VP
Average B, all atoms (Å ²)	173.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.78	0/3247	1.38	2/4447 (0.0%)
1	B	0.77	0/3154	1.40	2/4320 (0.0%)
1	D	0.76	0/3253	1.38	2/4446 (0.0%)
1	E	0.77	0/3136	1.40	2/4299 (0.0%)
2	F	0.87	0/18	1.02	0/23
2	G	0.87	0/27	1.06	0/33
2	H	0.88	0/27	1.09	0/33
2	I	0.69	0/18	0.95	0/23
All	All	0.77	0/12880	1.39	8/17624 (0.0%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	97	GLY	CA-C-N	6.19	126.37	120.43
1	A	97	GLY	C-N-CA	6.19	126.37	120.43
1	D	97	GLY	CA-C-N	6.18	126.36	120.43
1	D	97	GLY	C-N-CA	6.18	126.36	120.43
1	B	97	GLY	CA-C-N	6.05	126.24	120.43
1	B	97	GLY	C-N-CA	6.05	126.24	120.43
1	E	97	GLY	CA-C-N	5.88	126.08	120.43
1	E	97	GLY	C-N-CA	5.88	126.08	120.43

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3153	0	2871	8	0
1	B	3061	0	2709	5	0
1	D	3156	0	2908	10	0
1	E	3045	0	2712	7	0
2	F	19	0	15	1	0
2	G	28	0	24	0	0
2	H	28	0	24	1	0
2	I	19	0	15	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
All	All	12513	0	11278	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:127:LEU:HD13	1:E:385:LEU:HD21	1.79	0.63
1:D:105:SER:HB2	1:D:125:VAL:HG13	1.83	0.60
1:B:105:SER:HB2	1:B:125:VAL:HG13	1.82	0.60
1:A:105:SER:HB2	1:A:125:VAL:HG13	1.85	0.58
1:E:55:LEU:HA	1:E:58:ILE:HD12	1.87	0.57
1:A:265:ASN:HD21	2:F:662:SER:HB2	1.71	0.55
1:D:388:LEU:O	1:D:392:PHE:HB2	2.08	0.54
1:A:388:LEU:O	1:A:392:PHE:HB2	2.09	0.53
1:B:388:LEU:O	1:B:392:PHE:HB2	2.08	0.53
1:E:388:LEU:O	1:E:392:PHE:HB2	2.09	0.53
1:E:265:ASN:HD21	2:I:662:SER:HB2	1.74	0.51
1:A:195:TYR:HB3	1:A:198:ILE:HD12	1.94	0.50
1:E:125:VAL:HA	1:E:128:LEU:HD12	1.94	0.49
1:D:195:TYR:HB3	1:D:198:ILE:HD12	1.95	0.48
1:D:127:LEU:HD13	1:D:385:LEU:HD21	1.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:59:MET:HE2	1:D:63:THR:HG22	1.95	0.47
1:D:265:ASN:HD21	2:H:662:SER:HB2	1.79	0.47
1:D:431:LEU:HD23	1:D:469:LEU:HD21	1.99	0.45
1:B:351:ILE:HG21	1:B:407:LEU:HD21	1.98	0.44
1:A:127:LEU:HD13	1:A:385:LEU:HD21	2.00	0.43
1:D:351:ILE:HG21	1:D:407:LEU:HD21	2.00	0.43
1:A:351:ILE:HG21	1:A:407:LEU:HD21	2.00	0.42
1:A:423:LYS:HA	1:A:428:ALA:HB2	2.02	0.42
1:B:423:LYS:HA	1:B:428:ALA:HB2	2.02	0.42
1:E:423:LYS:HA	1:E:428:ALA:HB2	2.02	0.41
1:D:125:VAL:HA	1:D:128:LEU:HD12	2.01	0.41
1:D:423:LYS:HA	1:D:428:ALA:HB2	2.02	0.41
1:A:171:PHE:O	1:A:175:GLN:HG2	2.20	0.41
1:B:125:VAL:HA	1:B:128:LEU:HD12	2.02	0.40
1:E:105:SER:HB2	1:E:125:VAL:HG13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	417/482 (86%)	405 (97%)	12 (3%)	0	100	100
1	B	409/482 (85%)	396 (97%)	13 (3%)	0	100	100
1	D	408/482 (85%)	396 (97%)	12 (3%)	0	100	100
1	E	407/482 (84%)	396 (97%)	11 (3%)	0	100	100
2	F	1/4 (25%)	0	1 (100%)	0	100	100
2	G	2/4 (50%)	1 (50%)	1 (50%)	0	100	100
2	H	2/4 (50%)	1 (50%)	1 (50%)	0	100	100
2	I	1/4 (25%)	0	1 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1647/1944 (85%)	1595 (97%)	52 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	295/424 (70%)	287 (97%)	8 (3%)	39	59
1	B	277/424 (65%)	271 (98%)	6 (2%)	45	63
1	D	303/424 (72%)	295 (97%)	8 (3%)	40	60
1	E	282/424 (66%)	276 (98%)	6 (2%)	47	64
2	F	2/4 (50%)	2 (100%)	0	100	100
2	G	3/4 (75%)	3 (100%)	0	100	100
2	H	3/4 (75%)	3 (100%)	0	100	100
2	I	2/4 (50%)	2 (100%)	0	100	100
All	All	1167/1712 (68%)	1139 (98%)	28 (2%)	43	62

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	LEU
1	A	41	GLN
1	A	119	GLU
1	A	124	LEU
1	A	166	ASP
1	A	372	LEU
1	A	454	MET
1	A	463	LEU
1	B	25	LEU
1	B	41	GLN
1	B	119	GLU
1	B	124	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	160	LEU
1	B	166	ASP
1	D	25	LEU
1	D	41	GLN
1	D	50	HIS
1	D	61	SER
1	D	119	GLU
1	D	124	LEU
1	D	160	LEU
1	D	166	ASP
1	E	41	GLN
1	E	119	GLU
1	E	160	LEU
1	E	166	ASP
1	E	169	LYS
1	E	437	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	265	ASN
1	A	353	GLN
1	A	467	GLN
1	B	265	ASN
1	B	353	GLN
1	B	355	ASN
1	B	439	ASN
1	D	50	HIS
1	D	353	GLN
1	D	355	ASN
1	D	467	GLN
1	E	41	GLN
1	E	265	ASN
1	E	353	GLN
1	E	355	ASN
1	E	439	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	424/482 (87%)	0.23	17 (4%)	42	30	62, 170, 240, 260	1 (0%)
1	B	416/482 (86%)	0.44	29 (6%)	22	19	63, 178, 252, 261	1 (0%)
1	D	415/482 (86%)	0.25	25 (6%)	27	22	70, 167, 222, 242	1 (0%)
1	E	414/482 (85%)	0.26	13 (3%)	51	35	66, 198, 239, 257	1 (0%)
2	F	3/4 (75%)	1.33	1 (33%)	1	2	175, 175, 180, 181	0
2	G	4/4 (100%)	2.15	1 (25%)	2	3	206, 208, 211, 214	0
2	H	4/4 (100%)	1.03	1 (25%)	2	3	174, 174, 178, 181	0
2	I	3/4 (75%)	1.63	1 (33%)	1	2	217, 217, 220, 235	0
All	All	1683/1944 (86%)	0.31	88 (5%)	33	25	62, 176, 237, 261	4 (0%)

All (88) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	347	LYS	7.4
1	B	327	GLU	7.2
1	B	346	VAL	7.1
2	G	661	CYS	5.7
1	D	406	CYS	4.6
1	E	51	VAL	4.5
1	D	22	ALA	4.5
1	B	348	ASN	4.4
1	E	195	TYR	4.3
1	A	26	PHE	4.2
1	D	345	THR	4.1
1	A	212	VAL	3.9
2	I	661	CYS	3.8
1	D	343	GLY	3.8
1	B	430	ASP	3.8
1	E	194	ASP	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	346	VAL	3.7
1	D	88	GLU	3.7
1	B	18	ARG	3.6
1	A	346	VAL	3.6
1	A	73	ASP	3.6
2	F	661	CYS	3.5
1	B	168	ILE	3.4
1	D	232	GLY	3.3
1	D	430	ASP	3.3
1	D	215	TYR	3.3
1	A	200	ALA	3.3
1	E	420	ALA	3.3
1	E	50	HIS	3.2
1	D	460	PRO	3.2
2	H	661	CYS	3.1
1	A	398	PRO	3.1
1	A	173	VAL	3.1
1	D	21	GLY	3.1
1	B	324	CYS	3.0
1	B	32	LEU	3.0
1	B	52	PRO	2.9
1	D	229	LEU	2.9
1	B	73	ASP	2.9
1	B	54	GLU	2.8
1	B	19	ILE	2.7
1	A	172	VAL	2.7
1	B	434	ALA	2.7
1	D	24	LEU	2.6
1	B	232	GLY	2.6
1	B	55	LEU	2.6
1	E	238	ILE	2.5
1	A	211	LEU	2.5
1	D	95	PHE	2.5
1	A	104	LEU	2.5
1	B	51	VAL	2.5
1	D	324	CYS	2.4
1	D	168	ILE	2.4
1	A	366	LEU	2.4
1	E	192	GLY	2.4
1	B	35	THR	2.4
1	D	87	THR	2.4
1	A	337	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	379	TYR	2.4
1	E	11	TRP	2.3
1	B	76	THR	2.3
1	E	461	PRO	2.3
1	D	402	VAL	2.3
1	D	211	LEU	2.3
1	E	17	LYS	2.3
1	B	101	LEU	2.2
1	B	336	ALA	2.2
1	B	385	LEU	2.2
1	A	348	ASN	2.2
1	B	343	GLY	2.2
1	E	212	VAL	2.2
1	D	461	PRO	2.2
1	D	153	HIS	2.2
1	B	133[A]	PHE	2.1
1	B	220	ALA	2.1
1	A	168	ILE	2.1
1	D	350	ILE	2.1
1	B	345	THR	2.1
1	A	277	ILE	2.1
1	B	326	ASN	2.1
1	E	12	GLU	2.1
1	D	349	ILE	2.1
1	D	441	ASP	2.0
1	D	235	LYS	2.0
1	B	266	ALA	2.0
1	B	173	VAL	2.0
1	A	100	TYR	2.0
1	A	94	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ZN	B	600	1/1	0.96	0.09	154,154,154,154	0
3	ZN	E	600	1/1	0.96	0.05	187,187,187,187	0
3	ZN	A	600	1/1	0.98	0.07	141,141,141,141	0
3	ZN	D	600	1/1	0.99	0.05	130,130,130,130	0

6.5 Other polymers [i](#)

There are no such residues in this entry.