



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:18 AM UTC

PDB ID : 4YPE / pdb_00004ype
Title : ASH1L SET domain H2193F mutant in complex with S-adenosyl methionine (SAM)
Authors : Rogawski, D.S.; Ndoj, J.; Cho, H.J.; Maillard, I.; Grembecka, J.; Cierpicki, T.
Deposited on : 2015-03-12
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	226	9% 74% 20% • •
1	B	226	8% 68% 19% • • 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ZN	B	2302	-	-	X	-
2	ZN	B	2303	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3572 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase ASH1L.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	2	0
			1760	1096	310	330	24			
1	B	202	Total	C	N	O	S	0	1	0
			1645	1026	291	307	21			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2063	GLY	-	expression tag	UNP Q9NR48
A	2064	ALA	-	expression tag	UNP Q9NR48
A	2065	MET	-	expression tag	UNP Q9NR48
A	2066	ALA	-	expression tag	UNP Q9NR48
A	2067	GLY	-	expression tag	UNP Q9NR48
A	2068	SER	-	expression tag	UNP Q9NR48
A	2193	PHE	HIS	engineered mutation	UNP Q9NR48
B	2063	GLY	-	expression tag	UNP Q9NR48
B	2064	ALA	-	expression tag	UNP Q9NR48
B	2065	MET	-	expression tag	UNP Q9NR48
B	2066	ALA	-	expression tag	UNP Q9NR48
B	2067	GLY	-	expression tag	UNP Q9NR48
B	2068	SER	-	expression tag	UNP Q9NR48
B	2193	PHE	HIS	engineered mutation	UNP Q9NR48

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Zn	0	0
			3	3		
2	B	3	Total	Zn	0	0
			3	3		

- Molecule 3 is S-ADENOSYLMETHIONINE (CCD ID: SAM) (formula: C₁₅H₂₂N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
3	B	1	Total	C	N	O	S	0	0
			27	15	6	5	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	43	Total	O	0	0
			43	43		
4	B	64	Total	O	0	0
			64	64		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Histone-lysine N-methyltransferase ASH1L



• Molecule 1: Histone-lysine N-methyltransferase ASH1L



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	58.84Å 58.84Å 232.15Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.56 – 2.20 42.56 – 2.20	Depositor EDS
% Data completeness (in resolution range)	85.9 (42.56-2.20) 86.0 (42.56-2.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.96 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.216 , 0.263 0.213 , 0.261	Depositor DCC
R_{free} test set	1094 reflections (4.42%)	wwPDB-VP
Wilson B-factor (Å ²)	30.6	Xtriage
Anisotropy	0.213	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.050 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3572	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAM, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.19	1/1804 (0.1%)	1.19	8/2425 (0.3%)
1	B	1.26	3/1683 (0.2%)	1.21	9/2262 (0.4%)
All	All	1.22	4/3487 (0.1%)	1.20	17/4687 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2262	VAL	C-O	8.24	1.32	1.24
1	B	2262	VAL	CA-CB	6.21	1.61	1.53
1	B	2236	ARG	CD-NE	-5.71	1.38	1.46
1	B	2184	ILE	C-O	-5.27	1.18	1.24

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2262	VAL	CB-CA-C	-11.12	97.73	110.41
1	B	2148	GLU	CB-CA-C	-7.96	100.01	111.95
1	A	2262	VAL	N-CA-C	-7.44	94.31	107.18
1	B	2157	LYS	N-CA-C	-6.30	105.23	113.17
1	B	2148	GLU	N-CA-C	6.07	119.55	111.30
1	B	2236	ARG	NE-CZ-NH2	-5.96	113.83	119.20
1	B	2190	HIS	N-CA-C	5.60	117.48	107.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	2148	GLU	N-CA-CB	-5.55	103.68	111.62
1	B	2118	SER	N-CA-C	-5.37	102.90	110.31
1	A	2097	ASP	N-CA-C	5.26	117.02	111.28
1	A	2197	ASN	N-CA-C	-5.18	102.61	110.28
1	A	2166	ILE	N-CA-C	5.08	116.25	111.48
1	A	2118	SER	CA-C-N	5.05	125.33	119.47
1	A	2118	SER	C-N-CA	5.05	125.33	119.47
1	A	2105	VAL	N-CA-C	5.04	115.32	110.74
1	B	2090	THR	N-CA-C	5.03	118.50	111.56
1	B	2265	GLN	N-CA-C	5.02	117.31	107.57

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	2147	ALA	Peptide
1	B	2260	PHE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1760	0	1672	39	0
1	B	1645	0	1563	34	0
2	A	3	0	0	0	0
2	B	3	0	0	5	0
3	A	27	0	22	1	0
3	B	27	0	22	3	0
4	A	43	0	0	0	0
4	B	64	0	0	2	0
All	All	3572	0	3279	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2104:CYS:SG	2:B:2303:ZN:ZN	1.25	1.22
1:A:2245:MET:HE1	1:A:2251:LEU:HD23	1.41	1.02
1:A:2195[B]:CYS:SG	1:A:2203:VAL:HG13	2.09	0.92
1:A:2156:THR:HG21	1:A:2160:LEU:HD11	1.49	0.92
1:A:2245:MET:HE1	1:A:2251:LEU:CD2	2.01	0.90
1:B:2091:CYS:HG	2:B:2303:ZN:ZN	0.64	0.90
1:A:2156:THR:HG21	1:A:2160:LEU:CD1	2.10	0.81
1:A:2195[B]:CYS:SG	1:A:2203:VAL:CG1	2.70	0.80
1:A:2199:ASP:OD2	1:A:2228:LYS:NZ	2.16	0.78
1:A:2099:ASP:HA	1:A:2127:GLN:NE2	1.98	0.78
1:B:2073:ARG:NH2	4:B:2401:HOH:O	2.06	0.73
1:B:2179:PHE:CE2	1:B:2183:MET:HE3	2.30	0.66
1:B:2192:ASP:O	3:B:2304:SAM:H5'1	1.96	0.66
1:B:2144:ARG:HD2	1:B:2215:PHE:CD2	2.34	0.63
1:B:2128:CYS:SG	2:B:2302:ZN:ZN	1.88	0.60
1:A:2098:ASP:OD1	1:A:2100:THR:OG1	2.18	0.60
1:A:2108:CYS:O	1:A:2112:MET:HG2	2.02	0.60
1:A:2197:ASN:HD21	1:A:2260:PHE:HD1	1.48	0.59
1:B:2255:TYR:O	1:B:2259:SER:HB2	2.04	0.58
1:A:2148:GLU:HG3	1:A:2149:GLU:H	1.68	0.57
1:B:2090:THR:CG2	1:B:2121:THR:HB	2.35	0.56
1:A:2142:LEU:CD2	1:A:2156:THR:HG22	2.35	0.56
1:B:2265:GLN:OE1	1:B:2278:ILE:HD12	2.07	0.55
1:A:2106:ASP:O	1:A:2112:MET:CE	2.54	0.55
1:B:2090:THR:O	1:B:2091:CYS:C	2.50	0.54
1:A:2106:ASP:O	1:A:2112:MET:HE3	2.08	0.54
1:B:2179:PHE:CZ	1:B:2183:MET:CE	2.91	0.54
1:B:2218:HIS:HB2	1:B:2255:TYR:CD2	2.43	0.54
1:B:2211:ASN:HB2	4:B:2420:HOH:O	2.08	0.53
1:A:2268:CYS:HB2	1:A:2279:ILE:HG13	1.92	0.51
1:B:2111:ARG:NH2	1:B:2129:CYS:O	2.44	0.51
1:B:2128:CYS:HG	2:B:2302:ZN:ZN	1.25	0.51
1:A:2228:LYS:HD3	1:A:2235:TYR:CD2	2.47	0.50
1:A:2150:LYS:O	3:A:2304:SAM:N	2.45	0.49
1:B:2265:GLN:NE2	1:B:2280:GLY:HA3	2.27	0.49
1:B:2168:GLU:OE2	1:B:2236:ARG:HD3	2.12	0.49
1:B:2214:ARG:O	3:B:2304:SAM:HE2	2.13	0.49
1:A:2139:VAL:CG1	1:A:2141:CYS:SG	3.00	0.49
1:A:2195[B]:CYS:SG	1:A:2203:VAL:HG11	2.50	0.48
1:A:2139:VAL:HG12	1:A:2141:CYS:SG	2.53	0.48
1:B:2190:HIS:ND1	1:B:2192:ASP:N	2.58	0.48
1:A:2132:ARG:HH11	1:A:2132:ARG:HG3	1.78	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2090:THR:HG22	1:B:2121:THR:HB	1.95	0.48
1:A:2183:MET:HG3	1:A:2189:ASN:HA	1.95	0.48
1:A:2260:PHE:CG	1:A:2261:ASN:N	2.82	0.48
1:A:2141:CYS:SG	1:A:2158:GLU:HG2	2.53	0.48
1:A:2156:THR:CG2	1:A:2160:LEU:HD11	2.31	0.47
1:A:2240:TYR:N	1:A:2240:TYR:CD1	2.82	0.47
1:A:2087:GLU:O	1:A:2089:THR:HG23	2.15	0.47
1:B:2211:ASN:O	1:B:2214:ARG:HG2	2.15	0.47
1:A:2246:PRO:HD2	1:A:2249:THR:OG1	2.14	0.47
1:A:2176:GLU:HB2	1:A:2201:GLY:O	2.16	0.46
1:A:2245:MET:HE3	1:A:2249:THR:CG2	2.46	0.46
1:A:2218:HIS:HB2	1:A:2255:TYR:CG	2.51	0.45
1:B:2082:PRO:HG3	1:B:2198:LEU:HD13	1.98	0.45
1:B:2218:HIS:HB2	1:B:2255:TYR:CG	2.50	0.45
1:A:2084:SER:HA	1:A:2235:TYR:CE1	2.52	0.45
1:B:2144:ARG:HD3	1:B:2212:GLU:OE1	2.18	0.44
1:B:2190:HIS:CD2	1:B:2264:LYS:HB3	2.52	0.44
1:A:2142:LEU:HD23	1:A:2156:THR:HG22	1.99	0.43
1:A:2195[A]:CYS:HA	1:A:2204:ILE:O	2.18	0.43
1:A:2082:PRO:HA	1:A:2202:MET:HE1	2.00	0.43
1:A:2255:TYR:O	1:A:2259:SER:HB2	2.19	0.43
1:B:2191:SER:OG	1:B:2262:VAL:HG21	2.18	0.43
1:B:2091:CYS:SG	1:B:2104:CYS:SG	3.01	0.42
1:B:2091:CYS:SG	2:B:2303:ZN:ZN	1.81	0.42
1:B:2108:CYS:O	1:B:2112:MET:HG2	2.19	0.41
1:B:2110:ASN:HB3	1:B:2115:ALA:O	2.19	0.41
1:A:2195[B]:CYS:HA	1:A:2204:ILE:O	2.20	0.41
1:A:2199:ASP:O	1:A:2202:MET:HE3	2.21	0.41
1:B:2150:LYS:O	3:B:2304:SAM:N	2.54	0.41
1:B:2204:ILE:HD13	1:B:2204:ILE:HG21	1.93	0.40
1:B:2195:CYS:HA	1:B:2204:ILE:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/226 (96%)	198 (91%)	19 (9%)	0	100	100
1	B	199/226 (88%)	186 (94%)	12 (6%)	1 (0%)	24	27
All	All	416/452 (92%)	384 (92%)	31 (8%)	1 (0%)	43	51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	2125	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/198 (98%)	190 (98%)	4 (2%)	47	63
1	B	181/198 (91%)	175 (97%)	6 (3%)	33	45
All	All	375/396 (95%)	365 (97%)	10 (3%)	39	53

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2105	VAL
1	A	2149	GLU
1	A	2203	VAL
1	A	2262	VAL
1	B	2089	THR
1	B	2090	THR
1	B	2127	GLN
1	B	2139	VAL
1	B	2262	VAL
1	B	2266	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	2127	GLN
1	A	2134	GLN
1	A	2136	HIS
1	A	2190	HIS
1	A	2197	ASN
1	B	2110	ASN
1	B	2177	GLN
1	B	2186	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SAM	A	2304	-	27,29,29	1.32	5 (18%)	34,42,42	2.29	11 (32%)
3	SAM	B	2304	-	27,29,29	1.25	4 (14%)	34,42,42	2.15	7 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SAM	A	2304	-	-	3/17/33/33	0/3/3/3
3	SAM	B	2304	-	-	5/17/33/33	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2304	SAM	C2-N1	3.17	1.39	1.33
3	A	2304	SAM	C2-N3	2.83	1.39	1.33
3	B	2304	SAM	C2-N3	2.71	1.38	1.33
3	A	2304	SAM	C2-N1	2.65	1.38	1.33
3	A	2304	SAM	OXT-C	-2.60	1.22	1.30
3	A	2304	SAM	C8-N7	2.59	1.36	1.31
3	B	2304	SAM	OXT-C	-2.50	1.22	1.30
3	A	2304	SAM	C5-N7	-2.43	1.34	1.39
3	B	2304	SAM	C5-N7	-2.37	1.34	1.39

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2304	SAM	N3-C2-N1	-5.71	119.94	128.58
3	B	2304	SAM	N3-C2-N1	-5.33	120.52	128.58
3	A	2304	SAM	C5-C4-N3	-5.29	119.43	126.72
3	A	2304	SAM	OXT-C-O	-4.54	113.78	124.08
3	B	2304	SAM	N9-C8-N7	-4.39	107.71	113.94
3	B	2304	SAM	C5-C4-N3	-4.36	120.71	126.72
3	A	2304	SAM	C5-N7-C8	4.31	110.22	103.45
3	B	2304	SAM	C5-N7-C8	4.22	110.09	103.45
3	A	2304	SAM	N9-C8-N7	-3.76	108.60	113.94
3	B	2304	SAM	C2-N3-C4	3.70	120.86	111.83
3	A	2304	SAM	C2-N3-C4	3.66	120.77	111.83
3	A	2304	SAM	C4-C5-N7	-3.52	106.56	110.58
3	B	2304	SAM	C4-C5-N7	-3.40	106.70	110.58
3	A	2304	SAM	N3-C4-N9	2.94	132.17	127.17
3	B	2304	SAM	N3-C4-N9	2.59	131.57	127.17
3	A	2304	SAM	C5-C4-N9	2.36	108.38	105.81
3	A	2304	SAM	O4'-C1'-N9	2.25	112.40	108.09
3	A	2304	SAM	C6-C5-C4	2.08	120.02	117.18

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	2304	SAM	CB-CG-SD-C5'
3	B	2304	SAM	CB-CG-SD-CE
3	B	2304	SAM	CB-CG-SD-C5'
3	A	2304	SAM	CB-CG-SD-CE
3	B	2304	SAM	O-C-CA-CB
3	B	2304	SAM	C-CA-CB-CG
3	A	2304	SAM	O-C-CA-CB
3	B	2304	SAM	OXT-C-CA-CB


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2304	SAM	1	0
3	B	2304	SAM	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	217/226 (96%)	0.13	20 (9%)	14 12	14, 31, 61, 74	2 (0%)
1	B	202/226 (89%)	0.18	19 (9%)	14 11	11, 27, 65, 81	1 (0%)
All	All	419/452 (92%)	0.15	39 (9%)	14 12	11, 29, 64, 81	3 (0%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2262	VAL	6.0
1	B	2104	CYS	4.4
1	A	2262	VAL	4.0
1	B	2105	VAL	3.9
1	A	2257[A]	PHE	3.8
1	A	2107	ASP	3.8
1	A	2085	GLY	3.6
1	B	2260	PHE	3.5
1	A	2064	ALA	3.5
1	B	2091	CYS	3.2
1	B	2259	SER	3.1
1	B	2264	LYS	3.1
1	A	2149	GLU	2.8
1	B	2090	THR	2.7
1	A	2263	GLU	2.6
1	B	2073	ARG	2.6
1	B	2280	GLY	2.6
1	B	2106	ASP	2.5
1	B	2261	ASN	2.5
1	B	2122	CYS	2.4
1	A	2088	ALA	2.4
1	A	2260	PHE	2.3
1	B	2089	THR	2.3
1	B	2258	HIS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	2267	LEU	2.3
1	A	2146	ARG	2.2
1	A	2099	ASP	2.2
1	B	2107	ASP	2.2
1	A	2089	THR	2.2
1	A	2100	THR	2.1
1	B	2109	LEU	2.1
1	A	2148	GLU	2.1
1	A	2189	ASN	2.1
1	A	2190	HIS	2.1
1	B	2086	TYR	2.1
1	A	2065	MET	2.0
1	A	2097	ASP	2.0
1	A	2105	VAL	2.0
1	A	2191	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	B	2302	1/1	0.90	0.12	25,25,25,25	1
3	SAM	A	2304	27/27	0.94	0.07	19,25,32,33	0
2	ZN	B	2303	1/1	0.95	0.09	36,36,36,36	1
3	SAM	B	2304	27/27	0.95	0.07	25,28,34,35	0
2	ZN	A	2303	1/1	0.98	0.07	20,20,20,20	1
2	ZN	A	2301	1/1	0.99	0.04	23,23,23,23	1
2	ZN	B	2301	1/1	0.99	0.04	19,19,19,19	1
2	ZN	A	2302	1/1	0.99	0.04	19,19,19,19	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around SAM A 2304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around SAM B 2304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.