



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 09:36 PM UTC

PDB ID : 1YQ3 / pdb_00001yq3
Title : Avian respiratory complex ii with oxaloacetate and ubiquinone
Authors : Huang, L.; Cobessi, D.; Tung, E.Y.; Berry, E.A.
Deposited on : 2005-02-01
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	621	% 64% 30% ..
2	B	252	3% 68% 20% 7% .
3	C	141	% 57% 34% 8% .
4	D	103	2% 64% 30% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	TEO	A	1002	X	-	X	-

2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 9576 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinate dehydrogenase flavoprotein subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	614	Total	C	N	O	S	0	0	0
			4736	2962	845	900	29			

- Molecule 2 is a protein called succinate dehydrogenase Ip subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	242	Total	C	N	O	S	0	0	0
			1930	1220	327	361	22			


- Molecule 3 is a protein called SUCCINATE DEHYDROGENASE CYTOCHROME B, LARGE SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	140	Total	C	N	O	S	0	0	1
			1071	703	179	185	4			

- Molecule 4 is a protein called SUCCINATE DEHYDROGENASE CYTOCHROME B, SMALL SUBUNIT.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	102	Total	C	N	O	S	0	0	0
			770	508	122	137	3			

- Molecule 5 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 6 is MALATE LIKE INTERMEDIATE (CCD ID: TEO) (formula: C₄H₄O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			9	4	5		

- Molecule 7 is UNKNOWN LIGAND (CCD ID: UNL) (formula:).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	15	Total	O	0	0
			15	15		
7	B	6	Total	O	0	0
			6	6		
7	C	4	Total	O	0	0
			4	4		
7	D	5	Total	O	0	0
			5	5		

- Molecule 8 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 9 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 10 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe_3S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 11 is Coenzyme Q10, (2Z,6E,10Z,14E,18E,22E,26Z)-isomer (CCD ID: UQ) (formula: $\text{C}_{59}\text{H}_{90}\text{O}_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	B	1	Total	C	O	0	0
			14	10	4		

- Molecule 12 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 13 is hexyl beta-D-glucopyranoside (CCD ID: JZR) (formula: $C_{12}H_{24}O_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
13	C	1	Total	C	O		0	0
			18	12	6			

- Molecule 14 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
14	C	1	Total	C	Fe	N	O	0	0
			41	32	1	4	4		

- Molecule 15 is 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (CCD ID: PEE) (formula: $C_{41}H_{78}NO_8P$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
15	C	1	Total	C	O	0	0
			21	19	2		
15	D	1	Total	C		0	0
			24	24			


- Molecule 16 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	440	Total	O	0	0
			440	440		
16	B	213	Total	O	0	0
			213	213		
16	C	96	Total	O	0	0
			96	96		
16	D	85	Total	O	0	0
			85	85		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Succinate dehydrogenase flavoprotein subunit





● Molecule 4: SUCCINATE DEHYDROGENASE CYTOCHROME B, SMALL SUBUNIT



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.01Å 84.40Å 289.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.66 – 2.20 38.66 – 2.20	Depositor EDS
% Data completeness (in resolution range)	89.2 (38.66-2.20) 89.3 (38.66-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 2.20Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.175 , 0.223 0.168 , 0.214	Depositor DCC
R_{free} test set	3842 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	39.5	Xtriage
Anisotropy	0.642	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 60.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9576	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, FES, JZR, TEO, F3S, GOL, PEE, FAD, UQ, HEM, UNL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.57	40/4837 (0.8%)	1.53	64/6550 (1.0%)
2	B	1.71	23/1972 (1.2%)	1.54	28/2661 (1.1%)
3	C	1.70	12/1100 (1.1%)	1.50	16/1497 (1.1%)
4	D	1.66	10/793 (1.3%)	1.42	6/1089 (0.6%)
All	All	1.63	85/8702 (1.0%)	1.52	114/11797 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
3	C	0	1
4	D	0	1
All	All	0	7

The worst 5 of 85 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	191	MET	SD-CE	-19.22	1.31	1.79
3	C	36	MET	SD-CE	-17.70	1.35	1.79
2	B	241	MET	SD-CE	-11.90	1.49	1.79
1	A	284	MET	SD-CE	-11.47	1.50	1.79
3	C	126	VAL	CA-CB	11.16	1.68	1.54

The worst 5 of 114 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	255	THR	N-CA-C	14.20	128.56	108.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	11	GLN	N-CA-C	-10.20	100.84	113.18
1	A	219	THR	N-CA-C	-10.07	100.99	113.18
2	B	210	PHE	N-CA-C	9.44	125.32	113.43
1	A	51	PHE	N-CA-C	-9.36	97.94	109.83

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	113	TYR	Sidechain
1	A	172	TYR	Sidechain
1	A	322	TYR	Sidechain
1	A	365	TYR	Sidechain
1	A	98	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4736	0	4616	116	0
2	B	1930	0	1916	55	0
3	C	1071	0	1106	44	0
4	D	770	0	763	32	0
5	A	53	0	29	5	0
6	A	9	0	2	4	0
7	A	15	0	0	0	0
7	B	6	0	0	0	0
7	C	4	0	0	0	0
7	D	5	0	0	1	0
8	B	4	0	0	0	0
9	B	8	0	0	0	0
10	B	7	0	0	0	0
11	B	14	0	9	13	0
12	B	6	0	8	0	0
13	C	18	0	24	1	0
14	C	41	0	24	2	0
15	C	21	0	35	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
15	D	24	0	40	4	0
16	A	440	0	0	18	0
16	B	213	0	0	8	0
16	C	96	0	0	7	0
16	D	85	0	0	2	0
All	All	9576	0	8572	233	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 233 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:123:LYS:CE	2:B:123:LYS:NZ	1.68	1.51
1:A:476:MET:CE	1:A:476:MET:SD	2.01	1.48
2:B:185:MET:SD	2:B:185:MET:CE	2.01	1.47
1:A:284:MET:HE1	1:A:303:SER:HB2	1.40	0.99
2:B:123:LYS:HE3	3:C:8:GLU:OE1	1.62	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	612/621 (99%)	583 (95%)	28 (5%)	1 (0%)	43	51
2	B	240/252 (95%)	227 (95%)	12 (5%)	1 (0%)	30	34
3	C	138/141 (98%)	132 (96%)	5 (4%)	1 (1%)	18	19
4	D	100/103 (97%)	97 (97%)	3 (3%)	0	100	100
All	All	1090/1117 (98%)	1039 (95%)	48 (4%)	3 (0%)	36	42

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	8	THR
1	A	9	SER
3	C	78	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	498/506 (98%)	483 (97%)	15 (3%)	36	49
2	B	214/219 (98%)	206 (96%)	8 (4%)	30	41
3	C	116/119 (98%)	113 (97%)	3 (3%)	40	55
4	D	78/79 (99%)	74 (95%)	4 (5%)	21	27
All	All	906/923 (98%)	876 (97%)	30 (3%)	33	45

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	590	LYS
4	D	19	LEU
2	B	98	LYS
4	D	66	LYS
3	C	40	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	121	GLN
3	C	26	HIS
2	B	144	GLN
3	C	42	HIS
1	A	320	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 41 ligands modelled in this entry, 30 are unknown - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
13	JZR	C	142	-	18,18,18	2.21	7 (38%)	23,23,23	1.50	4 (17%)
14	HEM	C	143	3,4	48,48,50	3.14	24 (50%)	66,80,82	3.20	30 (45%)
8	FES	B	1002	2	0,4,4	-	-	-		
10	F3S	B	1004	2	0,9,9	-	-	-		
9	SF4	B	1003	2	0,12,12	-	-	-		
6	TEO	A	1002	-	5,8,8	1.73	2 (40%)	4,10,10	2.24	2 (50%)
11	UQ	B	1005	-	14,14,63	2.29	8 (57%)	20,20,79	1.04	0
15	PEE	C	144	-	19,19,50	2.10	4 (21%)	19,19,55	0.87	2 (10%)
5	FAD	A	1001	1	58,58,58	2.83	27 (46%)	85,89,89	1.99	24 (28%)
12	GOL	B	1012	-	5,5,5	1.78	2 (40%)	5,5,5	0.90	0
15	PEE	D	104	-	22,22,50	1.63	4 (18%)	20,20,55	1.25	4 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
13	JZR	C	142	-	-	2/9/29/29	0/1/1/1
14	HEM	C	143	3,4	-	1/10/50/54	-
8	FES	B	1002	2	-	-	0/1/1/1
10	F3S	B	1004	2	-	-	0/3/3/3
9	SF4	B	1003	2	-	-	0/6/5/5
6	TEO	A	1002	-	1/1/3/4	4/6/8/8	-
11	UQ	B	1005	-	-	1/4/28/87	0/1/1/1
15	PEE	C	144	-	-	12/17/17/54	-
5	FAD	A	1001	1	-	9/34/50/50	0/6/6/6
12	GOL	B	1012	-	-	3/4/4/4	-
15	PEE	D	104	-	-	8/18/18/54	-

The worst 5 of 78 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1001	FAD	PA-O3P	-12.03	1.46	1.59
14	C	143	HEM	CMC-C2C	8.83	1.68	1.50
14	C	143	HEM	C1B-NB	8.48	1.54	1.40
14	C	143	HEM	CMB-C2B	7.74	1.66	1.50
5	A	1001	FAD	C10-N10	6.82	1.52	1.37

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	C	143	HEM	C4D-ND-C1D	-9.37	94.11	105.21
14	C	143	HEM	C1B-NB-C4B	-8.79	94.79	105.21
14	C	143	HEM	CHC-C1C-NC	-8.44	115.27	124.45
14	C	143	HEM	C2B-C1B-NB	7.44	118.39	109.84
5	A	1001	FAD	N3A-C2A-N1A	-6.75	118.37	128.58

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	A	1002	TEO	C2

5 of 40 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1001	FAD	N10-C1'-C2'-O2'
5	A	1001	FAD	N10-C1'-C2'-C3'
5	A	1001	FAD	O4'-C4'-C5'-O5'

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	1001	FAD	C5'-O5'-P-O2P
5	A	1001	FAD	C5'-O5'-P-O3P


There are no ring outliers.

7 monomers are involved in 32 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
13	C	142	JZR	1	0
14	C	143	HEM	2	0
6	A	1002	TEO	4	0
11	B	1005	UQ	13	0
15	C	144	PEE	3	0
5	A	1001	FAD	5	0
15	D	104	PEE	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand JZR C 142






Ligand HEM C 143



Ligand UQ B 1005





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	614/621 (98%)	-0.42	4 (0%) 84 82	27, 42, 64, 130	0
2	B	242/252 (96%)	-0.49	7 (2%) 53 50	25, 37, 66, 105	0
3	C	140/141 (99%)	-0.30	1 (0%) 84 82	28, 42, 69, 80	0
4	D	102/103 (99%)	-0.41	2 (1%) 65 62	33, 40, 61, 72	0
All	All	1098/1117 (98%)	-0.42	14 (1%) 75 73	25, 41, 66, 130	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	138	GLY	5.2
2	B	6	ALA	4.0
2	B	7	ALA	3.9
1	A	8	ILE	3.5
1	A	9	SER	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	UNL	B	1007	1/-	0.71	0.30	63,63,63,63	0
12	GOL	B	1012	6/6	0.77	0.18	112,113,114,114	0
7	UNL	A	1007	1/-	0.79	0.12	66,66,66,66	0
7	UNL	D	205	1/-	0.80	0.12	59,59,59,59	0
7	UNL	A	1015	1/-	0.82	0.12	76,76,76,76	0
7	UNL	B	1010	1/-	0.82	0.21	75,75,75,75	0
7	UNL	D	228	1/-	0.83	0.12	59,59,59,59	0
15	PEE	D	104	24/51	0.84	0.22	47,62,73,77	0
11	UQ	B	1005	14/63	0.85	0.23	59,72,82,86	0
7	UNL	A	1013	1/-	0.85	0.17	77,77,77,77	0
7	UNL	A	1004	1/-	0.85	0.14	63,63,63,63	0
15	PEE	C	144	21/51	0.86	0.17	56,64,74,77	0
7	UNL	A	1016	1/-	0.87	0.12	63,63,63,63	0
7	UNL	A	1017	1/-	0.87	0.12	66,66,66,66	0
7	UNL	A	1009	1/-	0.89	0.07	65,65,65,65	0
7	UNL	C	203	1/-	0.89	0.22	61,61,61,61	0
13	JZR	C	142	18/18	0.91	0.12	41,47,63,67	0
7	UNL	C	212	1/-	0.92	0.28	76,76,76,76	0
7	UNL	C	226	1/-	0.92	0.09	56,56,56,56	0
7	UNL	D	204	1/-	0.92	0.17	65,65,65,65	0
7	UNL	A	1010	1/-	0.92	0.11	58,58,58,58	0
7	UNL	B	1008	1/-	0.92	0.15	59,59,59,59	0
7	UNL	B	1006	1/-	0.93	0.09	57,57,57,57	0
7	UNL	A	1008	1/-	0.93	0.07	58,58,58,58	0
7	UNL	A	1005	1/-	0.93	0.14	64,64,64,64	0
7	UNL	A	1003	1/-	0.93	0.20	65,65,65,65	0
7	UNL	A	1012	1/-	0.93	0.28	63,63,63,63	0
7	UNL	B	1009	1/-	0.94	0.21	60,60,60,60	0
7	UNL	A	1006	1/-	0.94	0.10	47,47,47,47	0
7	UNL	D	209	1/-	0.94	0.16	61,61,61,61	0
7	UNL	D	218	1/-	0.94	0.10	47,47,47,47	0
7	UNL	B	1011	1/-	0.94	0.10	50,50,50,50	0
7	UNL	C	211	1/-	0.95	0.10	68,68,68,68	0
7	UNL	A	1014	1/-	0.96	0.07	59,59,59,59	0
6	TEO	A	1002	9/9	0.96	0.06	35,37,38,42	0
7	UNL	A	1011	1/-	0.96	0.11	54,54,54,54	0
5	FAD	A	1001	53/53	0.98	0.05	23,32,37,37	0
14	HEM	C	143	41/43	0.98	0.06	32,36,44,49	0
9	SF4	B	1003	8/8	0.99	0.02	31,32,35,36	0
8	FES	B	1002	4/4	1.00	0.02	31,33,34,34	0
10	F3S	B	1004	7/7	1.00	0.03	28,29,31,31	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around UQ B 1005:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PEE C 144:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around JZR C 142:**


$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FAD A 1001:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.