



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:30 PM UTC

PDB ID : 5YQ1 / pdb_00005yq1
Title : Crystal structure of E.coli aminopeptidase N in complex with O-Methyl-L-tyrosine
Authors : Marapaka, A.K.; Ganji, R.J.; Reddi, R.; Addlagatta, A.
Deposited on : 2017-11-04
Resolution : 1.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	891	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MLI	A	904	-	X	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MLI	A	905	-	X	-	-
5	GOL	A	908	-	X	-	-
5	GOL	A	910	-	X	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7661 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminopeptidase N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	866	6982	4422	1208	1324	28	0	8	0


There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP P04825
A	-19	GLY	-	expression tag	UNP P04825
A	-18	SER	-	expression tag	UNP P04825
A	-17	SER	-	expression tag	UNP P04825
A	-16	HIS	-	expression tag	UNP P04825
A	-15	HIS	-	expression tag	UNP P04825
A	-14	HIS	-	expression tag	UNP P04825
A	-13	HIS	-	expression tag	UNP P04825
A	-12	HIS	-	expression tag	UNP P04825
A	-11	HIS	-	expression tag	UNP P04825
A	-10	SER	-	expression tag	UNP P04825
A	-9	SER	-	expression tag	UNP P04825
A	-8	GLY	-	expression tag	UNP P04825
A	-7	LEU	-	expression tag	UNP P04825
A	-6	VAL	-	expression tag	UNP P04825
A	-5	PRO	-	expression tag	UNP P04825
A	-4	ARG	-	expression tag	UNP P04825
A	-3	GLY	-	expression tag	UNP P04825
A	-2	SER	-	expression tag	UNP P04825
A	-1	HIS	-	expression tag	UNP P04825
A	0	MET	-	expression tag	UNP P04825

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is O-methyl-L-tyrosine (CCD ID: 0A1) (formula: $C_{10}H_{13}NO_3$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	10	1	3		

- Molecule 4 is MALONATE ION (CCD ID: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	3	4		
4	A	1	Total	C	O	0	0
			7	3	4		
4	A	1	Total	C	O	0	0
			7	3	4		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	607	Total	O	0	0
			607	607		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aminopeptidase N



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	120.41Å 120.41Å 169.78Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 1.58 50.00 – 1.58	Depositor EDS
% Data completeness (in resolution range)	98.9 (50.00-1.58) 98.9 (50.00-1.58)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 1.58Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.164 , 0.182 0.177 , 0.193	Depositor DCC
R_{free} test set	9687 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 35.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.014 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7661	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MLI, ZN, 0A1, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.56	41/7161 (0.6%)	1.39	27/9722 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (41) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	198	ARG	CA-C	-8.10	1.43	1.52
1	A	500	GLN	CA-C	-7.75	1.43	1.52
1	A	560	ASN	CG-OD1	7.56	1.38	1.23
1	A	865	ILE	N-CA	7.46	1.55	1.46
1	A	349	ARG	CD-NE	-7.19	1.36	1.46
1	A	809	VAL	CA-CB	6.93	1.62	1.54
1	A	357	ALA	CA-CB	-6.89	1.42	1.53
1	A	796	ALA	C-O	6.73	1.32	1.24
1	A	650	THR	C-O	6.68	1.31	1.24
1	A	354	ALA	N-CA	6.65	1.54	1.46
1	A	671	GLU	N-CA	-6.64	1.37	1.46
1	A	258	PHE	N-CA	6.51	1.53	1.46
1	A	357	ALA	N-CA	6.33	1.54	1.46
1	A	685	LEU	N-CA	-6.03	1.39	1.46
1	A	50	ALA	CA-C	5.99	1.60	1.52
1	A	499	LEU	CA-C	-5.96	1.45	1.52
1	A	29	LEU	CB-CG	-5.89	1.41	1.53
1	A	658	ALA	CA-CB	-5.81	1.44	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	525	TYR	CB-CG	5.74	1.64	1.51
1	A	9	TYR	CA-C	-5.73	1.45	1.52
1	A	264	GLU	CA-C	5.73	1.60	1.52
1	A	275	TYR	N-CA	-5.72	1.39	1.46
1	A	715	ALA	CA-CB	-5.68	1.44	1.53
1	A	650	THR	N-CA	5.67	1.53	1.46
1	A	204	ARG	N-CA	-5.58	1.39	1.46
1	A	407	TYR	C-O	-5.37	1.17	1.24
1	A	591	LEU	N-CA	5.36	1.53	1.46
1	A	393	THR	C-O	5.32	1.30	1.24
1	A	196	VAL	N-CA	-5.30	1.39	1.46
1	A	151	LYS	N-CA	-5.26	1.39	1.46
1	A	286	LYS	N-CA	5.25	1.52	1.46
1	A	529	VAL	CA-CB	-5.16	1.48	1.54
1	A	94	ILE	C-O	5.11	1.29	1.24
1	A	590	SER	C-O	-5.10	1.17	1.24
1	A	287	ASP	CA-CB	-5.09	1.45	1.53
1	A	748	LYS	C-O	5.08	1.30	1.24
1	A	423	GLN	CG-CD	-5.08	1.39	1.52
1	A	830	LEU	CB-CG	-5.06	1.43	1.53
1	A	625	MET	SD-CE	-5.06	1.66	1.79
1	A	560	ASN	CB-CG	5.06	1.64	1.52
1	A	507	ASN	CA-C	5.01	1.59	1.52

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	293	ARG	NE-CZ-NH2	-13.74	106.84	119.20
1	A	293	ARG	NE-CZ-NH1	13.50	135.00	121.50
1	A	293	ARG	CD-NE-CZ	13.16	142.82	124.40
1	A	795	ALA	N-CA-C	8.71	120.77	111.28
1	A	675	ILE	O-C-N	7.20	128.85	121.87
1	A	457	GLU	CA-CB-CG	6.75	127.61	114.10
1	A	339	VAL	N-CA-CB	6.26	117.87	110.55
1	A	126	ARG	NE-CZ-NH2	6.12	124.71	119.20
1	A	624	GLU	OE1-CD-OE2	5.99	137.28	122.90
1	A	220	LEU	N-CA-C	5.88	118.47	111.71
1	A	761	VAL	N-CA-C	5.88	116.53	110.36
1	A	675	ILE	CA-C-O	-5.78	114.94	120.95
1	A	793	ASN	CA-C-N	-5.64	113.81	119.56
1	A	793	ASN	C-N-CA	-5.64	113.81	119.56
1	A	681	ARG	NE-CZ-NH2	5.62	124.26	119.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	556	ARG	NE-CZ-NH2	5.51	124.16	119.20
1	A	800	GLU	N-CA-C	5.50	118.04	111.71
1	A	627	GLU	CA-CB-CG	-5.31	103.48	114.10
1	A	378	LEU	O-C-N	-5.26	116.54	122.12
1	A	445	THR	CA-CB-OG1	5.26	117.48	109.60
1	A	838	ALA	N-CA-C	5.21	116.96	111.28
1	A	293	ARG	CB-CG-CD	5.19	123.24	111.30
1	A	111	GLN	CB-CG-CD	-5.18	103.80	112.60
1	A	621	SER	CA-C-O	-5.12	116.16	121.99
1	A	410	ARG	N-CA-C	5.05	116.87	111.36
1	A	223	ALA	O-C-N	-5.05	115.51	121.32
1	A	77	GLU	CA-CB-CG	-5.02	104.06	114.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	293	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6982	0	6836	26	0
2	A	1	0	0	0	0
3	A	14	0	12	0	0
4	A	21	0	6	0	0
5	A	36	0	46	6	0
6	A	607	0	0	9	0
All	All	7661	0	6900	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:627:GLU:HG3	6:A:1322:HOH:O	1.67	0.92
1:A:346[A]:ARG:NH1	1:A:615:GLU:OE1	2.12	0.82
5:A:910:GOL:H12	6:A:1556:HOH:O	1.84	0.77
1:A:546:TRP:CZ2	5:A:909:GOL:H2	2.23	0.74
5:A:910:GOL:H11	6:A:1474:HOH:O	1.89	0.73
1:A:535[B]:GLU:OE1	6:A:1001:HOH:O	2.13	0.67
1:A:608:ILE:HD11	1:A:612:LEU:HD23	1.75	0.66
1:A:640:VAL:HG22	5:A:908:GOL:H11	1.81	0.60
1:A:488:GLU:OE2	1:A:559:ARG:NH2	2.36	0.58
1:A:343:ASN:OD1	1:A:346[B]:ARG:NH2	2.37	0.56
1:A:627:GLU:HG2	6:A:1056:HOH:O	2.08	0.53
1:A:862:TYR:O	1:A:866:THR:HG23	2.09	0.52
1:A:825:ARG:NH1	6:A:1011:HOH:O	2.44	0.51
1:A:714:LEU:HD22	1:A:748:LYS:HD3	1.92	0.50
1:A:535[B]:GLU:OE2	1:A:560:ASN:CG	2.54	0.50
1:A:608:ILE:HD13	1:A:609:ASP:N	2.27	0.50
1:A:346[B]:ARG:HH11	1:A:346[B]:ARG:HD3	1.34	0.47
1:A:840:ARG:HG2	6:A:1201:HOH:O	2.16	0.46
1:A:576:ILE:HG23	1:A:625:MET:HE3	1.99	0.45
1:A:87:PRO:HG2	1:A:90:PHE:HB3	1.99	0.45
1:A:492:ASN:HB3	1:A:526:PHE:CE2	2.52	0.43
5:A:910:GOL:C1	6:A:1474:HOH:O	2.54	0.43
1:A:222:ARG:HG2	1:A:279:ARG:HB2	2.02	0.42
1:A:223:ALA:N	1:A:224:PRO:CD	2.83	0.41
1:A:681:ARG:NH1	6:A:1017:HOH:O	2.46	0.41
1:A:44:ARG:HH11	1:A:44:ARG:HD3	1.65	0.41
1:A:346[A]:ARG:NH2	1:A:565:TRP:HE1	2.18	0.41
1:A:605:ASP:O	1:A:608:ILE:HG22	2.20	0.41
1:A:643:ALA:HB2	5:A:908:GOL:H32	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	872/891 (98%)	859 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	749/763 (98%)	738 (98%)	11 (2%)	57	31

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	220	LEU
1	A	293	ARG
1	A	327	ASP
1	A	356	ASP
1	A	517	GLN
1	A	589	LEU
1	A	608	ILE
1	A	695	LEU
1	A	699	LEU
1	A	826	LEU
1	A	830	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	HIS
1	A	175	GLN
1	A	340	ASN
1	A	511	ASN
1	A	569	GLN
1	A	623	ASN
1	A	663	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	GOL	A	906	-	5,5,5	0.46	0	5,5,5	1.03	0
5	GOL	A	909	-	5,5,5	1.09	0	5,5,5	1.56	2 (40%)
4	MLI	A	903	-	6,6,6	1.55	0	7,7,7	1.35	1 (14%)
4	MLI	A	904	-	6,6,6	2.12	3 (50%)	7,7,7	2.57	4 (57%)
4	MLI	A	905	-	6,6,6	2.05	3 (50%)	7,7,7	1.77	4 (57%)
5	GOL	A	907	-	5,5,5	0.21	0	5,5,5	0.77	0
5	GOL	A	911	-	5,5,5	0.51	0	5,5,5	1.08	0
3	0A1	A	902	2	13,14,14	1.18	1 (7%)	14,18,18	2.03	5 (35%)
5	GOL	A	910	-	5,5,5	2.20	1 (20%)	5,5,5	3.15	4 (80%)
5	GOL	A	908	-	5,5,5	1.51	1 (20%)	5,5,5	2.95	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	906	-	-	2/4/4/4	-
5	GOL	A	909	-	-	2/4/4/4	-
4	MLI	A	903	-	-	0/4/4/4	-
4	MLI	A	904	-	-	2/4/4/4	-
4	MLI	A	905	-	-	0/4/4/4	-
5	GOL	A	907	-	-	2/4/4/4	-
5	GOL	A	911	-	-	2/4/4/4	-
3	0A1	A	902	2	-	0/10/10/10	0/1/1/1
5	GOL	A	910	-	-	1/4/4/4	-
5	GOL	A	908	-	-	3/4/4/4	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	910	GOL	O2-C2	4.38	1.56	1.43
4	A	904	MLI	C1-C2	3.51	1.56	1.51
4	A	905	MLI	C1-C3	3.43	1.56	1.51
4	A	905	MLI	O9-C3	-2.49	1.22	1.30
5	A	908	GOL	O1-C1	2.44	1.52	1.42
3	A	902	0A1	OH-CZ	2.36	1.42	1.37
4	A	904	MLI	O8-C3	2.30	1.29	1.22
4	A	904	MLI	C1-C3	2.09	1.54	1.51
4	A	905	MLI	C1-C2	2.05	1.54	1.51

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	908	GOL	C3-C2-C1	5.87	133.32	111.80
5	A	910	GOL	C3-C2-C1	-5.37	92.10	111.80
4	A	904	MLI	O7-C2-O6	-3.85	113.43	123.33
3	A	902	0A1	CG-CB-CA	3.83	121.97	114.13
4	A	904	MLI	C3-C1-C2	-3.49	100.62	112.95
5	A	910	GOL	O1-C1-C2	3.19	124.75	110.38
3	A	902	0A1	CD1-CE1-CZ	-3.10	116.19	119.73
4	A	904	MLI	O7-C2-C1	3.08	124.05	114.51
3	A	902	0A1	CB-CG-CD2	-2.81	115.67	120.90
3	A	902	0A1	CM-OH-CZ	-2.50	112.13	117.50
4	A	903	MLI	C3-C1-C2	-2.47	104.22	112.95
4	A	904	MLI	O9-C3-O8	-2.46	117.01	123.33
5	A	910	GOL	O2-C2-C3	2.45	119.31	109.18
5	A	909	GOL	O2-C2-C3	2.43	119.23	109.18
5	A	909	GOL	O2-C2-C1	2.42	119.19	109.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	902	0A1	CB-CG-CD1	2.40	125.37	120.90
4	A	905	MLI	O7-C2-C1	2.29	121.61	114.51
4	A	905	MLI	C3-C1-C2	-2.20	105.18	112.95
5	A	908	GOL	O1-C1-C2	2.19	120.23	110.38
5	A	910	GOL	O2-C2-C1	2.10	117.86	109.18
4	A	905	MLI	O9-C3-C1	2.07	120.92	114.51
5	A	908	GOL	O2-C2-C3	-2.01	100.84	109.18
4	A	905	MLI	O6-C2-C1	-2.01	116.39	122.11

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	907	GOL	C1-C2-C3-O3
5	A	907	GOL	O2-C2-C3-O3
5	A	908	GOL	O1-C1-C2-C3
5	A	911	GOL	O1-C1-C2-C3
5	A	908	GOL	O1-C1-C2-O2
5	A	909	GOL	O1-C1-C2-O2
5	A	909	GOL	O2-C2-C3-O3
5	A	911	GOL	O1-C1-C2-O2
5	A	908	GOL	O2-C2-C3-O3
5	A	910	GOL	O2-C2-C3-O3
5	A	906	GOL	C1-C2-C3-O3
4	A	904	MLI	C2-C1-C3-O8
4	A	904	MLI	C2-C1-C3-O9
5	A	906	GOL	O2-C2-C3-O3


There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	909	GOL	1	0
5	A	910	GOL	3	0
5	A	908	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	866/891 (97%)	0.13	18 (2%) 63 73	14, 25, 38, 63	8 (0%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	PRO	3.7
1	A	169	ASN	3.1
1	A	694	HIS	3.0
1	A	504	HIS	3.0
1	A	759	ALA	2.5
1	A	492	ASN	2.4
1	A	795	ALA	2.4
1	A	503	GLY	2.4
1	A	457	GLU	2.4
1	A	525	TYR	2.3
1	A	794	PRO	2.2
1	A	607	LYS	2.2
1	A	666	SER	2.2
1	A	560	ASN	2.2
1	A	606	GLU	2.2
1	A	695	LEU	2.0
1	A	855	GLU	2.0
1	A	502	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	A	910	6/6	0.62	0.28	43,54,56,61	0
5	GOL	A	907	6/6	0.81	0.16	55,60,68,80	0
4	MLI	A	905	7/7	0.82	0.14	44,48,51,55	0
4	MLI	A	904	7/7	0.84	0.17	49,57,64,74	0
5	GOL	A	906	6/6	0.89	0.15	54,62,63,72	0
5	GOL	A	909	6/6	0.90	0.13	33,39,45,62	0
5	GOL	A	908	6/6	0.90	0.13	31,36,45,52	0
5	GOL	A	911	6/6	0.90	0.11	43,51,56,59	0
4	MLI	A	903	7/7	0.95	0.07	30,33,38,41	0
3	0A1	A	902	14/14	0.96	0.07	20,22,29,36	0
2	ZN	A	901	1/1	1.00	0.02	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 0A1 A 902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.