



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 11:21 AM UTC

PDB ID : 1YS9 / pdb_00001ys9
Title : Crystal structure of phosphatase SPy1043 from Streptococcus pyogenes
Authors : Fedorov, A.A.; Fedorov, E.V.; Almo, S.C.; Burley, S.K.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2005-02-07
Resolution : 2.56 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1853 (2.58-2.54)
Clashscore	190562	1897 (2.58-2.54)
Ramachandran outliers	187476	1875 (2.58-2.54)
Sidechain outliers	187428	1875 (2.58-2.54)
RSRZ outliers	180081	1853 (2.58-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	254	7% 71% 22% 6%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1942 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein SPy1043.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	253	Total	C	N	O	S	0	0	0
			1942	1234	323	375	10			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	VAL	MET	engineered mutation	UNP Q99ZW4

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Protein SPY1043



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	119.92Å 120.14Å 58.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.39 – 2.56 24.39 – 2.56	Depositor EDS
% Data completeness (in resolution range)	96.4 (24.39-2.56) 96.2 (24.39-2.56)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.00 (at 2.54Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.226 , 0.252 0.227 , 0.252	Depositor DCC
R_{free} test set	705 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	25.5	Xtriage
Anisotropy	0.653	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 44.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.036 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	1942	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	1/1975 (0.1%)	1.04	11/2687 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	151	ILE	CA-CB	5.25	1.58	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	184	ALA	N-CA-C	9.25	120.97	111.07
1	A	247	SER	N-CA-C	-6.74	104.88	113.23
1	A	227	THR	N-CA-C	-6.71	99.95	110.36
1	A	8	ILE	N-CA-C	6.58	117.32	108.11
1	A	118	VAL	N-CA-C	5.97	116.78	107.28
1	A	95	THR	N-CA-C	5.67	117.46	111.28
1	A	228	GLY	N-CA-C	5.41	120.92	111.78
1	A	185	ILE	N-CA-C	5.32	115.94	110.36
1	A	91	VAL	N-CA-C	5.14	115.88	108.48
1	A	60	PHE	N-CA-C	5.04	118.97	111.87
1	A	85	ARG	N-CA-C	-5.00	101.98	109.79

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1942	0	1973	43	0
All	All	1942	0	1973	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:ILE:H	1:A:66:ILE:HD12	1.34	0.90
1:A:188:ASN:ND2	1:A:198:ARG:HH22	1.73	0.86
1:A:38:TYR:OH	1:A:64:THR:HG22	1.79	0.82
1:A:43:ASN:HD21	1:A:181:LYS:H	1.35	0.72
1:A:189:LYS:O	1:A:193:VAL:HG12	1.91	0.71
1:A:56:LEU:HB3	1:A:62:VAL:HG23	1.75	0.69
1:A:56:LEU:HB3	1:A:62:VAL:CG2	2.23	0.68
1:A:47:ARG:HG2	1:A:51:MET:HE2	1.76	0.65
1:A:225:VAL:HG11	1:A:249:LEU:HG	1.78	0.65
1:A:14:ILE:HD11	1:A:56:LEU:HD22	1.80	0.64
1:A:232:PRO:O	1:A:235:VAL:HG13	1.99	0.62
1:A:120:LEU:HD22	1:A:121:ASP:N	2.15	0.61
1:A:5:GLY:N	1:A:196:ILE:HD13	2.16	0.61
1:A:66:ILE:H	1:A:66:ILE:CD1	2.10	0.59
1:A:235:VAL:HA	1:A:238:LEU:HG	1.84	0.59
1:A:64:THR:HB	1:A:68:THR:OG1	2.04	0.58
1:A:133:THR:O	1:A:137:GLN:HG3	2.03	0.58
1:A:67:GLU:H	1:A:67:GLU:CD	2.10	0.58
1:A:208:TYR:CE2	1:A:213:MET:HE2	2.39	0.58
1:A:208:TYR:CZ	1:A:213:MET:HE2	2.39	0.57
1:A:253:ARG:HH11	1:A:253:ARG:HG3	1.72	0.55
1:A:29:LYS:O	1:A:33:GLU:HG3	2.07	0.54
1:A:196:ILE:HD11	1:A:201:ALA:HA	1.90	0.53
1:A:118:VAL:HG21	1:A:168:LEU:CD1	2.39	0.53
1:A:28:ILE:HD12	1:A:28:ILE:C	2.34	0.52
1:A:14:ILE:HD11	1:A:56:LEU:CD2	2.39	0.52
1:A:197:GLN:HB2	1:A:200:GLU:HG2	1.91	0.51
1:A:120:LEU:HD22	1:A:121:ASP:H	1.76	0.51
1:A:188:ASN:HD22	1:A:198:ARG:HH22	1.56	0.50
1:A:60:PHE:O	1:A:62:VAL:HG13	2.13	0.49
1:A:142:PHE:HD2	1:A:174:VAL:HG22	1.78	0.48
1:A:65:SER:HB3	1:A:67:GLU:OE1	2.13	0.48
1:A:167:LEU:C	1:A:167:LEU:CD1	2.87	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:26:ARG:NH1	1:A:250:ASP:OD2	2.47	0.47
1:A:109:GLU:OE2	1:A:111:GLU:N	2.47	0.47
1:A:226:THR:HB	1:A:232:PRO:HD3	1.96	0.47
1:A:239:PRO:O	1:A:240:ILE:HD13	2.16	0.46
1:A:172:THR:OG1	1:A:174:VAL:HG13	2.16	0.45
1:A:56:LEU:HD22	1:A:62:VAL:HG21	2.01	0.42
1:A:188:ASN:HD21	1:A:198:ARG:HH22	1.57	0.42
1:A:253:ARG:HG3	1:A:253:ARG:NH1	2.35	0.41
1:A:235:VAL:HG22	1:A:236:PRO:HD3	2.03	0.41
1:A:43:ASN:HD21	1:A:181:LYS:N	2.12	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	251/254 (99%)	240 (96%)	8 (3%)	3 (1%)	10 14

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	229	PHE
1	A	86	GLY
1	A	226	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	210/211 (100%)	190 (90%)	20 (10%)	8 10

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	ILE
1	A	28	ILE
1	A	64	THR
1	A	66	ILE
1	A	67	GLU
1	A	77	VAL
1	A	95	THR
1	A	120	LEU
1	A	134	LEU
1	A	149	LEU
1	A	167	LEU
1	A	173	ARG
1	A	174	VAL
1	A	182	PRO
1	A	191	LEU
1	A	193	VAL
1	A	209	LEU
1	A	225	VAL
1	A	235	VAL
1	A	254	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	43	ASN
1	A	44	ASN
1	A	53	GLN
1	A	58	ASN
1	A	188	ASN
1	A	197	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/254 (99%)	0.14	17 (6%) 24 23	10, 23, 47, 57	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	PRO	7.8
1	A	112	ASN	3.7
1	A	66	ILE	3.7
1	A	28	ILE	3.0
1	A	229	PHE	3.0
1	A	154	GLU	2.9
1	A	86	GLY	2.9
1	A	16	GLN	2.9
1	A	175	LYS	2.8
1	A	59	GLN	2.5
1	A	228	GLY	2.4
1	A	155	ARG	2.3
1	A	65	SER	2.2
1	A	253	ARG	2.2
1	A	231	ARG	2.1
1	A	227	THR	2.0
1	A	18	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.